



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 10, 2017 – 07:44 AM EDT

PDB ID : 2FIX  
Title : Structure of human liver FBPase complexed with potent benzoxazole allosteric inhibitors  
Authors : Abad-Zapatero, C.  
Deposited on : unknown  
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1322 (3.60-3.40)
Ramachandran outliers	110173	1283 (3.60-3.40)
Sidechain outliers	110143	1284 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	338	
1	D	338	
1	H	338	
1	L	338	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fructose-1,6-bisphosphatase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	1
			2432	1548	408	459	17			
1	D	319	Total	C	N	O	S	0	0	1
			2432	1548	408	459	17			
1	H	319	Total	C	N	O	S	0	0	1
			2432	1548	408	459	17			
1	L	319	Total	C	N	O	S	0	0	1
			2432	1548	408	459	17			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	217	LYS	ARG	VARIANT	GB 15277851
D	217	LYS	ARG	VARIANT	GB 15277851
H	217	LYS	ARG	VARIANT	GB 15277851
L	217	LYS	ARG	VARIANT	GB 15277851

- Molecule 2 is N-[7-(3-AMINOPHENYL)-5-METHOXY-1,3-BENZOXAZOL-2-YL]-2,5-DICHLOROBENZENESULFONAMIDE (three-letter code: 870) (formula: C<sub>20</sub>H<sub>15</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>4</sub>S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 30	C 20	Cl 2	N 3	O 4	S 1	0	0
2	D	1	Total 30	C 20	Cl 2	N 3	O 4	S 1	0	0
2	H	1	Total 30	C 20	Cl 2	N 3	O 4	S 1	0	0
2	L	1	Total 30	C 20	Cl 2	N 3	O 4	S 1	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

#### • Molecule 1: Fructose-1,6-bisphosphatase 1




#### • Molecule 1: Fructose-1,6-bisphosphatase 1






• Molecule 1: Fructose-1,6-bisphosphatase 1

Chain H: 20% 60% 14% 6%



• Molecule 1: Fructose-1,6-bisphosphatase 1

Chain L: 20% 54% 20% 6%



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.40 Å   108.67 Å   196.40 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	19.88 – 3.50	Depositor
% Data completeness (in resolution range)	61.6 (19.88-3.50)	Depositor
$R_{merge}$	0.24	Depositor
$R_{sym}$	0.24	Depositor
Refinement program	CNX 2002	Depositor
R, $R_{free}$	0.252 , 0.355	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	9848	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:  
870

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/2475	0.83	4/3343 (0.1%)
1	D	0.49	0/2475	0.78	1/3343 (0.0%)
1	H	0.48	0/2475	0.81	1/3343 (0.0%)
1	L	0.49	0/2475	0.82	2/3343 (0.1%)
All	All	0.50	0/9900	0.81	8/13372 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	SER	CB-CA-C	-8.14	94.64	110.10
1	A	25	ARG	O-C-N	-7.35	110.70	123.20
1	H	29	GLU	CB-CA-C	6.30	123.00	110.40
1	L	149	GLU	CB-CA-C	-6.27	97.86	110.40
1	A	132	VAL	N-CA-CB	-5.78	98.78	111.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	25	ARG	Mainchain



## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	0	2474	367	0
1	D	2432	0	2474	392	0
1	H	2432	0	2474	359	0
1	L	2432	0	2474	377	0
2	A	30	0	15	7	0
2	D	30	0	15	1	0
2	H	30	0	15	7	0
2	L	30	0	15	3	0
All	All	9848	0	9956	1406	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 71.

The worst 5 of 1406 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:53:ILE:HD13	1:L:185:MET:HG2	1.25	1.17
1:D:91:THR:HB	1:D:94:LEU:HD21	1.38	1.05
1:H:194:ILE:HG12	1:L:54:ALA:HB2	1.31	1.05
1:D:27:THR:HA	2:H:901:870:HN22	1.18	1.03
1:A:191:GLY:HA3	1:L:191:GLY:HA3	1.37	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/338 (93%)	207 (66%)	72 (23%)	36 (11%)	0	6
1	D	315/338 (93%)	192 (61%)	74 (24%)	49 (16%)	0	3
1	H	315/338 (93%)	211 (67%)	62 (20%)	42 (13%)	0	4
1	L	315/338 (93%)	196 (62%)	73 (23%)	46 (15%)	0	3
All	All	1260/1352 (93%)	806 (64%)	281 (22%)	173 (14%)	0	4

5 of 173 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	23	LYS
1	A	51	ALA
1	A	123	SER
1	A	199	ASP
1	A	222	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/281 (94%)	238 (90%)	27 (10%)	8	36
1	D	265/281 (94%)	225 (85%)	40 (15%)	3	19
1	H	265/281 (94%)	243 (92%)	22 (8%)	13	46
1	L	265/281 (94%)	226 (85%)	39 (15%)	3	20
All	All	1060/1124 (94%)	932 (88%)	128 (12%)	6	27

5 of 128 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	272	ASN
1	H	35	ASN
1	L	264	TYR
1	D	279	TYR
1	D	328	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	32	GLN
1	D	125	ASN
1	H	282	ASN
1	A	334	HIS
1	L	35	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	870	A	701	-	29,33,33	1.97	10 (34%)	41,49,49	1.23	3 (7%)
2	870	D	801	-	29,33,33	2.06	11 (37%)	41,49,49	1.36	5 (12%)
2	870	H	901	-	29,33,33	2.22	11 (37%)	41,49,49	1.50	8 (19%)
2	870	L	1001	-	29,33,33	1.98	9 (31%)	41,49,49	1.59	6 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	870	A	701	-	-	0/15/17/17	0/3/4/4
2	870	D	801	-	-	0/15/17/17	0/3/4/4
2	870	H	901	-	-	0/15/17/17	0/3/4/4
2	870	L	1001	-	-	0/15/17/17	0/3/4/4

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	870	C8-C7	-2.58	1.38	1.41
2	H	901	870	C8-C7	-2.36	1.38	1.41
2	D	801	870	C8-C7	-2.27	1.38	1.41
2	H	901	870	C2-CL2	-2.21	1.69	1.74
2	D	801	870	C3-C2	2.02	1.41	1.38

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	1001	870	O2-S-O1	-4.95	113.22	119.55
2	A	701	870	O4-C9-C8	-3.60	115.04	124.49
2	D	801	870	O2-S-O1	-3.50	115.08	119.55
2	L	1001	870	O4-C9-C8	-2.67	117.48	124.49
2	D	801	870	O4-C9-C8	-2.64	117.58	124.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	870	7	0
2	D	801	870	1	0
2	H	901	870	7	0
2	L	1001	870	3	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.