



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2017 – 07:37 AM EDT

PDB ID : 3FJO  
Title : Structure of chimeric YH CPR  
Authors : Morera, S.; Aigrain, L.; Truan, G.  
Deposited on : unknown  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	637	<div> <div>4%</div> <div>74%</div> <div>18%</div> <div>• 5%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NADPH-cytochrome P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	603	Total	C	N	O	S	0	0	0
			4769	3017	820	911	21			

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	21	MET	-	EXPRESSION TAG	UNP P16603
A	22	GLY	-	EXPRESSION TAG	UNP P16603
A	23	SER	-	EXPRESSION TAG	UNP P16603
A	24	SER	-	EXPRESSION TAG	UNP P16603
A	25	HIS	-	EXPRESSION TAG	UNP P16603
A	26	HIS	-	EXPRESSION TAG	UNP P16603
A	27	HIS	-	EXPRESSION TAG	UNP P16603
A	28	HIS	-	EXPRESSION TAG	UNP P16603
A	29	HIS	-	EXPRESSION TAG	UNP P16603
A	30	HIS	-	EXPRESSION TAG	UNP P16603
A	31	SER	-	EXPRESSION TAG	UNP P16603
A	32	SER	-	EXPRESSION TAG	UNP P16603
A	33	GLY	-	EXPRESSION TAG	UNP P16603
A	34	LEU	-	EXPRESSION TAG	UNP P16603
A	35	VAL	-	EXPRESSION TAG	UNP P16603
A	36	PRO	-	EXPRESSION TAG	UNP P16603
A	37	ARG	-	EXPRESSION TAG	UNP P16603
A	38	GLY	-	EXPRESSION TAG	UNP P16603
A	39	SER	-	EXPRESSION TAG	UNP P16603
A	40	HIS	-	EXPRESSION TAG	UNP P16603
A	41	MET	-	EXPRESSION TAG	UNP P16603
A	42	LEU	-	EXPRESSION TAG	UNP P16603
A	43	GLU	-	EXPRESSION TAG	UNP P16603

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	123	Total 123	O 123	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NADPH-cytochrome P450 reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	146.80 Å   60.70 Å   78.86 Å 90.00°   103.36°   90.00°	Depositor
Resolution (Å)	19.86 – 2.50 33.05 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.86-2.50) 94.8 (33.05-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.61 (at 2.20 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.222   ,   0.290 0.224   ,   0.295	Depositor DCC
$R_{free}$ test set	1156 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	35.9	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 45.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4976	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/4869	0.56	0/6593

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4769	0	4668	76	0
2	A	31	0	19	0	0
3	A	53	0	31	4	0
4	A	123	0	0	1	0
All	All	4976	0	4718	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:750:FAD:N10	3:A:750:FAD:C1'	1.67	1.56
1:A:212:GLY:O	1:A:213:VAL:HG22	1.71	0.90
1:A:217:GLY:O	1:A:218:GLU:HG2	1.75	0.85
3:A:750:FAD:C1'	3:A:750:FAD:C10	2.56	0.82
1:A:229:VAL:HG12	1:A:231:THR:HG23	1.63	0.80
1:A:318:LEU:HD23	1:A:421:ILE:HD12	1.64	0.79
1:A:50:ILE:O	1:A:54:VAL:HG23	1.85	0.76
3:A:750:FAD:C9A	3:A:750:FAD:C1'	2.63	0.75
1:A:411:LEU:HD23	1:A:417:LEU:HD12	1.70	0.72
1:A:125:ASP:O	1:A:128:VAL:HG23	1.96	0.66
1:A:574:ALA:HB2	1:A:588:LEU:HD11	1.77	0.66
1:A:110:ILE:HD12	1:A:211:LEU:HD21	1.81	0.63
1:A:574:ALA:HB2	1:A:588:LEU:CD1	2.28	0.62
1:A:639:VAL:HG23	4:A:10:HOH:O	2.00	0.62
1:A:115:ILE:HG21	1:A:149:MET:HE1	1.81	0.60
1:A:69:THR:O	1:A:69:THR:CG2	2.49	0.60
1:A:223:ARG:O	1:A:331:LEU:HD12	2.03	0.59
1:A:543:TYR:OH	1:A:596:LEU:HD11	2.03	0.59
1:A:444:VAL:CG2	1:A:445:HIS:CD2	2.87	0.57
1:A:60:ASN:HB3	1:A:89:LEU:HD22	1.86	0.57
1:A:624:TYR:CE1	1:A:639:VAL:HA	2.40	0.57
1:A:540:THR:O	1:A:568:LEU:HD12	2.05	0.57
1:A:105:ASN:HD21	1:A:144:ASN:H	1.51	0.56
1:A:115:ILE:HG21	1:A:149:MET:CE	2.37	0.54
1:A:398:SER:HA	1:A:402:GLU:HB2	1.90	0.54
1:A:434:ARG:HD2	1:A:470:ALA:HB2	1.88	0.54
1:A:613:ARG:CZ	1:A:655:VAL:HG21	2.37	0.54
1:A:67:SER:HB2	1:A:72:ALA:HB3	1.90	0.53
1:A:229:VAL:HG12	1:A:231:THR:CG2	2.37	0.53
1:A:303:TYR:CE1	1:A:433:ALA:HB2	2.44	0.53
1:A:594:GLU:HA	1:A:630:LEU:HD21	1.91	0.53
1:A:302:VAL:HG22	1:A:489:VAL:CG1	2.38	0.52
1:A:417:LEU:HD13	1:A:419:PRO:HD3	1.92	0.52
1:A:444:VAL:HG23	1:A:445:HIS:CD2	2.44	0.51
1:A:179:LEU:HD13	1:A:210:GLU:HG3	1.92	0.51
1:A:351:LEU:HD21	1:A:421:ILE:HD11	1.93	0.51
1:A:624:TYR:CZ	1:A:639:VAL:HG12	2.45	0.51
1:A:518:ALA:HB3	1:A:519:PRO:HD3	1.93	0.50
1:A:96:VAL:HG13	1:A:130:PHE:HB2	1.94	0.50
1:A:624:TYR:CE2	1:A:639:VAL:HG12	2.47	0.50
1:A:531:ARG:HG3	1:A:537:VAL:HG21	1.94	0.49
1:A:518:ALA:HB3	1:A:519:PRO:CD	2.43	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:541:LEU:HD12	1:A:541:LEU:N	2.28	0.49
1:A:473:TRP:CH2	1:A:489:VAL:HG13	2.48	0.49
1:A:224:GLN:HA	1:A:224:GLN:HE21	1.78	0.48
1:A:330:ASN:HD22	1:A:331:LEU:N	2.12	0.48
1:A:642:ILE:O	1:A:646:MET:HG3	2.14	0.48
1:A:242:GLU:HA	1:A:249:TYR:CZ	2.48	0.48
1:A:229:VAL:CG1	1:A:231:THR:HG23	2.38	0.48
1:A:473:TRP:HH2	1:A:489:VAL:HG13	1.78	0.47
1:A:60:ASN:HD22	1:A:90:ASN:H	1.62	0.47
1:A:151:GLY:C	1:A:152:LEU:HD12	2.35	0.47
1:A:212:GLY:O	1:A:213:VAL:CG2	2.55	0.46
1:A:369:LEU:HD22	1:A:419:PRO:HG3	1.97	0.46
1:A:294:TYR:CE1	1:A:438:ILE:HG21	2.51	0.45
1:A:242:GLU:OE1	1:A:246:LEU:HD23	2.17	0.45
1:A:152:LEU:N	1:A:152:LEU:HD12	2.32	0.45
1:A:573:VAL:HG11	1:A:575:PHE:CE1	2.51	0.45
1:A:69:THR:O	1:A:69:THR:HG22	2.16	0.45
1:A:369:LEU:HD22	1:A:419:PRO:CG	2.47	0.45
1:A:100:ASP:HB3	1:A:102:GLU:OE1	2.17	0.44
1:A:61:TYR:CE2	1:A:83:LEU:HD21	2.53	0.44
1:A:381:LEU:HD13	1:A:395:LEU:HD21	2.00	0.44
1:A:129:ASN:N	1:A:129:ASN:HD22	2.15	0.43
1:A:280:LEU:HD23	1:A:454:VAL:HB	1.99	0.43
1:A:642:ILE:HD13	1:A:645:LEU:HD12	1.99	0.43
1:A:656:TRP:HB3	3:A:750:FAD:C9	2.48	0.43
1:A:150:PHE:CZ	1:A:185:ALA:HB2	2.53	0.43
1:A:310:LEU:HD23	1:A:408:LEU:HD22	2.01	0.43
1:A:185:ALA:HB1	1:A:192:THR:HG23	2.01	0.43
1:A:120:GLU:O	1:A:120:GLU:CG	2.67	0.42
1:A:613:ARG:NH1	1:A:655:VAL:CG2	2.82	0.42
1:A:553:TYR:CG	1:A:556:ARG:HG3	2.54	0.42
1:A:301:ALA:HB2	1:A:435:TYR:CE2	2.56	0.41
1:A:369:LEU:O	1:A:417:LEU:HD21	2.21	0.41
1:A:590:LYS:HG2	1:A:622:THR:HG23	2.03	0.40
1:A:147:TYR:CE2	1:A:171:LEU:HD13	2.56	0.40
1:A:351:LEU:CD1	1:A:355:LEU:HD12	2.51	0.40
1:A:97:GLU:HG2	1:A:126:GLY:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	597/637 (94%)	566 (95%)	28 (5%)	3 (0%)	32	53

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	213	VAL
1	A	218	GLU
1	A	96	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	508/536 (95%)	463 (91%)	45 (9%)	11	22

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ASN
1	A	55	THR
1	A	68	GLN
1	A	69	THR
1	A	74	ASP
1	A	83	LEU
1	A	96	VAL
1	A	104	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	120	GLU
1	A	156	THR
1	A	224	GLN
1	A	234	ASP
1	A	246	LEU
1	A	247	LYS
1	A	259	LYS
1	A	267	THR
1	A	273	ASN
1	A	274	GLN
1	A	302	VAL
1	A	310	LEU
1	A	312	ASN
1	A	314	LEU
1	A	318	LEU
1	A	322	LEU
1	A	326	MET
1	A	330	ASN
1	A	334	GLU
1	A	337	LYS
1	A	351	LEU
1	A	364	ASN
1	A	380	GLU
1	A	381	LEU
1	A	388	SER
1	A	421	ILE
1	A	428	LEU
1	A	430	ARG
1	A	434	ARG
1	A	444	VAL
1	A	455	VAL
1	A	494	ARG
1	A	552	ASP
1	A	553	TYR
1	A	571	LEU
1	A	582	LYS
1	A	599	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	60	ASN
1	A	105	ASN
1	A	129	ASN
1	A	224	GLN
1	A	230	HIS
1	A	251	ASN
1	A	260	ASN
1	A	273	ASN
1	A	274	GLN
1	A	282	HIS
1	A	330	ASN
1	A	432	GLN
1	A	445	HIS
1	A	466	ASN
1	A	614	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FAD	A	750	-	51,58,58	2.90	7 (13%)	54,89,89	2.15	6 (11%)
2	FMN	A	751	-	31,33,33	1.54	6 (19%)	38,50,50	1.80	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	750	-	-	0/28/50/50	0/6/6/6
2	FMN	A	751	-	-	0/16/18/18	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	751	FMN	C7M-C7	-2.28	1.46	1.51
3	A	750	FAD	C5X-N5	2.13	1.38	1.35
2	A	751	FMN	C5A-N5	2.19	1.38	1.35
3	A	750	FAD	C2A-N1A	2.86	1.39	1.33
2	A	751	FMN	C4-N3	2.88	1.38	1.33
3	A	750	FAD	C4-N3	2.95	1.38	1.33
2	A	751	FMN	C1'-N10	3.18	1.51	1.48
2	A	751	FMN	C4A-N5	3.50	1.38	1.33
3	A	750	FAD	C4X-N5	3.65	1.38	1.33
3	A	750	FAD	C10-N1	3.87	1.38	1.33
3	A	750	FAD	C2A-N3A	4.03	1.38	1.32
2	A	751	FMN	C10-N1	4.37	1.39	1.33
3	A	750	FAD	C1'-N10	18.53	1.67	1.48

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	750	FAD	N3A-C2A-N1A	-11.96	118.44	128.86
3	A	750	FAD	C1'-N10-C10	-4.80	113.58	118.50
2	A	751	FMN	C8M-C8-C9	-3.24	112.23	120.34
3	A	750	FAD	C4X-C4-N3	-2.74	119.59	123.48
2	A	751	FMN	C4A-C4-N3	-2.65	119.71	123.48
2	A	751	FMN	C4-C4A-N5	2.10	120.98	118.68
3	A	750	FAD	C4X-N5-C5X	2.93	119.85	116.76
2	A	751	FMN	C1'-N10-C9A	3.01	121.11	118.35
3	A	750	FAD	C5X-C9A-N10	3.03	119.91	117.66
2	A	751	FMN	C8M-C8-C7	3.20	127.43	120.72

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	751	FMN	C4A-N5-C5A	3.56	120.53	116.76
3	A	750	FAD	C4-N3-C2	6.31	120.68	115.16
2	A	751	FMN	C4-N3-C2	7.28	121.53	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	750	FAD	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	603/637 (94%)	0.12	27 (4%) 34 36	19, 34, 53, 67	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	581	HIS	4.6
1	A	550	ASP	4.4
1	A	334	GLU	3.9
1	A	120	GLU	3.8
1	A	333	GLU	3.5
1	A	577	ARG	3.3
1	A	634	GLU	2.9
1	A	582	LYS	2.9
1	A	220	SER	2.7
1	A	214	GLU	2.6
1	A	222	ILE	2.6
1	A	584	TYR	2.6
1	A	549	SER	2.6
1	A	216	THR	2.6
1	A	231	THR	2.5
1	A	47	ASN	2.5
1	A	336	ASN	2.5
1	A	218	GLU	2.4
1	A	546	CYS	2.4
1	A	337	LYS	2.4
1	A	547	ARG	2.4
1	A	576	SER	2.3
1	A	575	PHE	2.2
1	A	213	VAL	2.2
1	A	234	ASP	2.1
1	A	583	VAL	2.1
1	A	331	LEU	2.1



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FMN	A	751	31/31	0.95	0.13	-0.72	19,23,23,23	0
3	FAD	A	750	53/53	0.96	0.10	-1.20	14,17,22,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.