



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 6, 2018 – 12:05 AM EST

PDB ID : 4FJU
Title : Crystal structure of ureidoglycolate dehydrogenase in ternary complex with NADH and glyoxylate
Authors : Kim, M.I.; Rhee, S.
Deposited on : 2012-06-12
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	7172 (1.80-1.76)
Clashscore	112137	8247 (1.80-1.76)
Ramachandran outliers	110173	8154 (1.80-1.76)
Sidechain outliers	110143	8153 (1.80-1.76)
RSRZ outliers	101464	7262 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	<div> <div>14%</div> <div> <div></div> <div>83%</div> <div>13%</div> <div>• •</div> </div> </div>
1	B	351	<div> <div>16%</div> <div> <div></div> <div>75%</div> <div>13%</div> <div>• 11%</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5749 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ureidoglycolate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	0	0
			2578	1625	452	485	16			
1	B	312	Total	C	N	O	S	0	0	0
			2371	1490	423	443	15			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP B1XGB5
A	0	HIS	-	EXPRESSION TAG	UNP B1XGB5
B	-1	GLY	-	EXPRESSION TAG	UNP B1XGB5
B	0	HIS	-	EXPRESSION TAG	UNP B1XGB5

- Molecule 2 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: C₂₁H₂₉N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is GLYOXYLIC ACID (three-letter code: GLV) (formula: $C_2H_2O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			5	2	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	330	Total	O	0	0
			330	330		
4	B	377	Total	O	0	0
			377	377		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ureidoglycolate dehydrogenase



• Molecule 1: Ureidoglycolate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	162.76 Å 162.76 Å 61.45 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.39 – 1.77 36.39 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (36.39-1.77) 91.6 (36.39-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.51 (at 1.81 Å)	Xtriage
Refinement program	CNS, PHENIX 1.7.2_869	Depositor
R, R_{free}	0.229 , 0.248 0.231 , 0.224	Depositor DCC
R_{free} test set	7080 reflections (10.07%)	DCC
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5749	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAI, GLV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2633	0.52	0/3569
1	B	0.37	0/2423	0.56	0/3283
All	All	0.36	0/5056	0.54	0/6852

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2578	0	2534	34	0
1	B	2371	0	2326	67	1
2	A	44	0	27	1	0
2	B	44	0	27	1	0
3	A	5	0	1	0	0
4	A	330	0	0	19	0
4	B	377	0	0	50	2
All	All	5749	0	4915	99	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:ARG:HD3	4:A:798:HOH:O	1.28	1.28
1:B:16:LEU:HA	4:B:773:HOH:O	1.42	1.15
1:B:45:GLY:C	4:B:842:HOH:O	1.94	1.05
1:A:189:ARG:NH2	4:A:645:HOH:O	1.88	1.04
1:B:38:ALA:CB	4:B:852:HOH:O	2.05	1.02
1:B:38:ALA:HB3	4:B:849:HOH:O	1.57	1.02
1:B:19:ALA:HB1	4:B:801:HOH:O	1.59	1.01
1:B:39:ARG:N	4:B:849:HOH:O	1.96	0.99
1:B:21:LEU:N	4:B:794:HOH:O	1.96	0.98
1:B:204:ASN:ND2	4:B:856:HOH:O	1.98	0.96
1:B:23:ARG:NH2	4:B:867:HOH:O	1.99	0.96
1:B:45:GLY:O	4:B:842:HOH:O	1.84	0.95
1:B:5:ARG:HB3	4:B:775:HOH:O	1.66	0.95
1:A:189:ARG:HA	4:A:762:HOH:O	1.65	0.94
1:B:19:ALA:HB3	4:B:773:HOH:O	1.68	0.93
1:B:204:ASN:CG	4:B:856:HOH:O	2.09	0.91
1:B:129:ARG:NH2	4:B:644:HOH:O	2.04	0.89
1:B:56:ILE:HG21	4:B:794:HOH:O	1.72	0.89
1:A:129:ARG:NH1	4:A:798:HOH:O	1.86	0.88
1:B:38:ALA:CB	4:B:849:HOH:O	2.18	0.86
1:B:20:GLY:CA	4:B:794:HOH:O	2.24	0.85
1:B:46:ALA:HA	4:B:842:HOH:O	1.77	0.85
1:B:5:ARG:CA	4:B:775:HOH:O	2.24	0.84
1:A:188:ALA:O	4:A:762:HOH:O	1.96	0.83
1:B:5:ARG:CB	4:B:775:HOH:O	2.23	0.83
1:A:194:SER:OG	4:A:820:HOH:O	1.89	0.83
1:B:20:GLY:N	4:B:794:HOH:O	2.13	0.82
1:B:19:ALA:O	4:B:801:HOH:O	1.97	0.81
1:B:49:VAL:HB	4:B:785:HOH:O	1.84	0.78
1:B:5:ARG:C	4:B:775:HOH:O	2.20	0.78
1:B:297:PRO:O	4:B:667:HOH:O	2.01	0.77
1:B:38:ALA:C	4:B:849:HOH:O	2.23	0.76
1:B:15:LYS:HG2	4:B:647:HOH:O	1.87	0.73
1:A:7:THR:HG21	4:A:766:HOH:O	1.88	0.73
1:B:5:ARG:O	4:B:775:HOH:O	2.08	0.71
1:A:251:MET:SD	4:A:704:HOH:O	2.48	0.71
1:B:46:ALA:N	4:B:842:HOH:O	2.19	0.69
1:A:102:GLN:OE1	4:A:602:HOH:O	2.09	0.69
1:B:16:LEU:O	4:B:773:HOH:O	2.10	0.69
1:B:37:ASP:OD1	4:B:840:HOH:O	2.11	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:SER:HB3	4:B:801:HOH:O	1.92	0.68
1:B:46:ALA:N	4:B:840:HOH:O	2.22	0.66
1:B:189:ARG:HG3	1:B:211:PRO:HB2	1.77	0.65
1:A:14:ASN:OD1	1:A:23:ARG:NH2	2.28	0.65
1:B:38:ALA:HB3	4:B:852:HOH:O	1.84	0.65
1:B:39:ARG:HG3	4:B:849:HOH:O	1.96	0.65
1:B:37:ASP:OD2	4:B:840:HOH:O	2.13	0.65
1:B:312:LYS:NZ	4:B:844:HOH:O	2.29	0.64
1:B:16:LEU:CA	4:B:773:HOH:O	2.15	0.64
1:B:158:PRO:HD3	2:B:401:NAI:H42N	1.81	0.63
1:A:42:HIS:NE2	4:A:719:HOH:O	2.30	0.61
1:B:46:ALA:CA	4:B:842:HOH:O	2.38	0.60
1:B:37:ASP:CG	4:B:840:HOH:O	2.40	0.59
1:B:46:ALA:HA	1:B:49:VAL:HG23	1.86	0.58
1:A:158:PRO:HD3	2:A:401:NAI:H42N	1.86	0.57
1:B:16:LEU:C	4:B:773:HOH:O	2.36	0.55
1:B:197:ASP:O	4:B:765:HOH:O	2.18	0.55
1:B:33:LEU:HD13	1:B:49:VAL:HG22	1.89	0.54
1:A:113:ARG:NE	4:A:738:HOH:O	2.04	0.52
1:A:129:ARG:NH1	4:A:593:HOH:O	2.11	0.52
1:B:34:VAL:CB	4:B:857:HOH:O	2.57	0.52
1:B:141:ASP:OD2	1:B:251:MET:HG3	2.10	0.52
1:B:43:SER:HB3	1:B:183:GLY:HA3	1.91	0.52
1:A:31:GLU:HA	1:A:34:VAL:HG22	1.93	0.51
1:A:24:GLU:HG2	4:A:701:HOH:O	2.10	0.51
1:A:191:ARG:HG3	1:A:193:MET:HG3	1.92	0.51
1:B:180:GLN:HG3	1:B:184:LYS:HG3	1.93	0.51
1:A:21:LEU:HG	1:A:56:ILE:HD13	1.94	0.50
1:B:204:ASN:CB	4:B:856:HOH:O	2.58	0.49
1:B:44:HIS:HD2	1:B:116:HIS:HE1	1.60	0.49
1:B:34:VAL:HG11	4:B:857:HOH:O	2.13	0.49
1:B:44:HIS:HD2	1:B:116:HIS:CE1	2.30	0.48
1:A:31:GLU:OE2	1:A:91:LYS:NZ	2.46	0.48
1:B:34:VAL:HB	4:B:857:HOH:O	2.12	0.48
1:A:5:ARG:NH1	4:A:680:HOH:O	2.47	0.47
1:A:44:HIS:HD2	1:A:116:HIS:CE1	2.33	0.47
1:A:18:GLN:HG3	1:A:336:LEU:HD21	1.97	0.46
1:A:187:ASP:O	1:A:191:ARG:HG2	2.16	0.45
1:A:338:ASN:C	4:A:815:HOH:O	2.54	0.45
1:B:50:GLU:N	4:B:621:HOH:O	2.49	0.45
1:B:38:ALA:CA	4:B:849:HOH:O	2.48	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:303:TYR:HB3	1:B:307:GLN:HB3	2.00	0.44
1:B:33:LEU:HD22	4:B:842:HOH:O	2.17	0.43
1:B:15:LYS:O	4:B:647:HOH:O	2.21	0.43
1:A:293:ILE:HG23	1:B:151:GLU:HB2	2.00	0.43
1:A:316:ALA:HB2	4:A:819:HOH:O	2.20	0.42
1:A:24:GLU:HB2	4:A:558:HOH:O	2.19	0.42
1:A:32:VAL:HB	1:A:87:GLN:HG2	2.01	0.42
1:B:7:THR:HG22	1:B:8:LEU:N	2.34	0.42
1:A:227:GLY:HA3	1:B:173:PHE:CE1	2.54	0.41
1:B:203:LYS:H	1:B:203:LYS:HG2	1.64	0.41
1:A:78:ILE:HD13	1:A:240:LEU:HD22	2.01	0.41
1:B:4:SER:O	1:B:7:THR:HB	2.21	0.41
1:A:73:GLY:HA2	4:A:802:HOH:O	2.20	0.41
1:B:33:LEU:HD21	1:B:52:TYR:HE2	1.85	0.41
1:A:180:GLN:HG3	1:A:199:TRP:CE2	2.55	0.40
1:B:211:PRO:HA	1:B:214:VAL:HG23	2.04	0.40
1:A:69:LEU:HD21	1:A:96:HIS:CE1	2.57	0.40
1:A:7:THR:CG2	4:A:766:HOH:O	2.58	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:520:HOH:O	4:B:558:HOH:O[8_554]	2.12	0.08
4:B:745:HOH:O	4:B:827:HOH:O[7_555]	2.12	0.08
1:B:7:THR:OG1	1:B:7:THR:OG1[7_555]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	336/351 (96%)	325 (97%)	10 (3%)	1 (0%)	44 27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	310/351 (88%)	298 (96%)	11 (4%)	1 (0%)	44	27
All	All	646/702 (92%)	623 (96%)	21 (3%)	2 (0%)	44	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	143	MET
1	A	157	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/278 (96%)	258 (97%)	9 (3%)	42	23
1	B	245/278 (88%)	241 (98%)	4 (2%)	68	55
All	All	512/556 (92%)	499 (98%)	13 (2%)	53	35

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	LEU
1	A	23	ARG
1	A	78	ILE
1	A	141	ASP
1	A	203	LYS
1	A	246	ARG
1	A	301	GLN
1	A	325	ASP
1	A	332	ILE
1	B	6	GLU
1	B	8	LEU
1	B	63	ARG
1	B	141	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	HIS
1	B	44	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAI	A	401	-	40,48,48	1.96	7 (17%)	41,73,73	1.66	3 (7%)
3	GLV	A	402	-	1,4,4	4.03	1 (100%)	0,4,4	0.00	-
2	NAI	B	401	-	40,48,48	1.93	7 (17%)	41,73,73	1.68	6 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAI	A	401	-	-	0/25/72/72	0/5/5/5
3	GLV	A	402	-	-	0/0/2/2	0/0/0/0
2	NAI	B	401	-	-	0/25/72/72	0/5/5/5

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAI	C2D-C3D	-5.07	1.40	1.53
2	A	401	NAI	C2D-C3D	-4.95	1.40	1.53
2	A	401	NAI	C2B-C3B	-4.41	1.41	1.53
2	A	401	NAI	C4N-C5N	-4.31	1.39	1.49
2	B	401	NAI	C2B-C3B	-4.27	1.42	1.53
2	B	401	NAI	C4N-C5N	-4.03	1.40	1.49
2	B	401	NAI	O4D-C4D	-2.61	1.39	1.45
2	A	401	NAI	C2B-C1B	-2.56	1.49	1.53
2	A	401	NAI	O4D-C4D	-2.56	1.39	1.45
2	B	401	NAI	C2B-C1B	-2.34	1.49	1.53
2	B	401	NAI	C6N-C5N	2.97	1.38	1.33
2	A	401	NAI	C6N-C5N	3.13	1.39	1.33
3	A	402	GLV	O1-C1	4.03	1.35	1.21
2	B	401	NAI	C7N-N7N	5.53	1.48	1.33
2	A	401	NAI	C7N-N7N	5.69	1.49	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAI	N3A-C2A-N1A	-8.58	121.39	128.86
2	B	401	NAI	N3A-C2A-N1A	-7.91	121.97	128.86
2	B	401	NAI	C4A-C5A-N7A	-2.63	106.87	109.41
2	A	401	NAI	C4A-C5A-N7A	-2.54	106.95	109.41
2	B	401	NAI	C3N-C2N-N1N	-2.49	119.47	123.08
2	A	401	NAI	C3N-C2N-N1N	-2.44	119.54	123.08
2	B	401	NAI	O5B-C5B-C4B	2.27	117.07	109.00
2	B	401	NAI	O5D-C5D-C4D	2.46	117.72	109.00
2	B	401	NAI	O4D-C1D-N1N	2.61	113.33	108.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NAI	1	0
2	B	401	NAI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	338/351 (96%)	0.96	49 (14%) 3 3	11, 27, 55, 73	0
1	B	312/351 (88%)	1.08	57 (18%) 1 1	11, 25, 54, 78	0
All	All	650/702 (92%)	1.02	106 (16%) 2 2	11, 26, 54, 78	0

All (106) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	193	MET	8.3
1	B	212	PHE	7.9
1	A	193	MET	7.0
1	B	190	SER	6.6
1	B	188	ALA	6.1
1	A	318	VAL	6.1
1	A	190	SER	5.9
1	A	212	PHE	5.8
1	A	194	SER	5.0
1	A	6	GLU	5.0
1	B	186	LEU	5.0
1	B	45	GLY	4.8
1	B	11	LEU	4.6
1	A	335	ALA	4.5
1	B	18	GLN	4.5
1	A	187	ASP	4.3
1	B	194	SER	4.2
1	B	7	THR	4.2
1	B	5	ARG	4.1
1	B	14	ASN	3.9
1	B	12	ILE	3.9
1	B	6	GLU	3.8
1	B	17	CYS	3.8
1	B	215	HIS	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	191	ARG	3.6
1	A	7	THR	3.6
1	B	264	LEU	3.6
1	A	192	ASN	3.6
1	B	192	ASN	3.5
1	B	47	VAL	3.5
1	B	10	GLN	3.5
1	B	49	VAL	3.4
1	B	191	ARG	3.4
1	B	315	LYS	3.4
1	B	232	ILE	3.3
1	B	57	SER	3.3
1	A	188	ALA	3.3
1	A	211	PRO	3.3
1	A	266	ILE	3.1
1	A	195	ILE	3.1
1	A	228	LEU	3.1
1	A	328	TYR	3.0
1	A	189	ARG	3.0
1	A	183	GLY	3.0
1	B	189	ARG	3.0
1	B	209	THR	2.9
1	B	214	VAL	2.9
1	B	9	HIS	2.9
1	A	332	ILE	2.9
1	B	41	ILE	2.9
1	A	186	LEU	2.9
1	A	252	TYR	2.9
1	B	46	ALA	2.9
1	A	173	PHE	2.8
1	A	232	ILE	2.8
1	A	323	ILE	2.8
1	B	53	ALA	2.8
1	A	3	ILE	2.8
1	B	196	PRO	2.7
1	A	294	THR	2.7
1	B	159	LEU	2.7
1	A	24	GLU	2.7
1	A	135	ILE	2.7
1	B	24	GLU	2.7
1	A	160	ALA	2.6
1	B	32	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	120	ILE	2.6
1	B	33	LEU	2.6
1	A	336	LEU	2.5
1	B	208	THR	2.5
1	A	159	LEU	2.5
1	B	211	PRO	2.4
1	B	30	ALA	2.4
1	A	300	ASN	2.4
1	B	228	LEU	2.4
1	B	181	ALA	2.4
1	A	338	ASN	2.4
1	B	160	ALA	2.4
1	A	324	VAL	2.3
1	A	17	CYS	2.3
1	A	64	GLU	2.3
1	A	264	LEU	2.3
1	B	201	VAL	2.3
1	B	235	LEU	2.3
1	B	195	ILE	2.3
1	B	23	ARG	2.2
1	B	314	ARG	2.2
1	A	29	VAL	2.2
1	A	210	ASP	2.2
1	B	210	ASP	2.2
1	B	44	HIS	2.2
1	A	118	GLY	2.2
1	A	326	ASP	2.2
1	B	135	ILE	2.2
1	A	14	ASN	2.1
1	A	2	LYS	2.1
1	A	161	PHE	2.1
1	B	34	VAL	2.1
1	A	12	ILE	2.1
1	B	52	TYR	2.0
1	B	50	GLU	2.0
1	A	209	THR	2.0
1	B	37	ASP	2.0
1	B	35	TYR	2.0
1	A	33	LEU	2.0
1	A	325	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GLV	A	402	5/5	0.90	0.13	0.09	36,37,40,42	0
2	NAI	B	401	44/44	0.94	0.12	-0.53	17,25,30,33	0
2	NAI	A	401	44/44	0.94	0.11	-0.68	18,24,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.