



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 12, 2017 – 06:05 pm GMT

PDB ID : 1FMH
Title : NMR SOLUTION STRUCTURE OF A DESIGNED HETERODIMERIC
LEUCINE ZIPPER
Authors : Marti, D.N.; Jelesarov, I.; Bosshard, H.R.
Deposited on : 2000-08-17

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 48%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	33	 73% 9% 18%
2	B	33	 67% 21% 12%

2 Ensemble composition and analysis

This entry contains 25 models. Model 14 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:27, B:1-B:29 (56)	0.24	14

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 7, 8, 9, 10, 12, 13, 14, 15, 17, 18, 20, 23, 24, 25
2	16, 21, 22
3	11, 19
Single-model clusters	6

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1005 atoms, of which 505 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4.

Mol	Chain	Residues	Atoms						Trace
1	A	33	Total	C	H	N	O	S	1
			475	150	226	41	57	1	

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLU	ARG	ENGINEERED	UNP P03069
A	2	VAL	MET	ENGINEERED	UNP P03069
A	3	ALA	LYS	ENGINEERED	UNP P03069
A	7	LYS	ASP	ENGINEERED	UNP P03069
A	8	GLU	LYS	ENGINEERED	UNP P03069
A	10	ALA	GLU	ENGINEERED	UNP P03069
A	11	GLN	GLU	ENGINEERED	UNP P03069
A	12	ALA	LEU	ENGINEERED	UNP P03069
A	13	GLU	LEU	ENGINEERED	UNP P03069
A	14	ALA	SER	ENGINEERED	UNP P03069
A	15	GLU	LYS	ENGINEERED	UNP P03069
A	18	GLN	HIS	ENGINEERED	UNP P03069
A	21	GLN	ASN	ENGINEERED	UNP P03069
A	25	GLN	ARG	ENGINEERED	UNP P03069
A	27	GLU	LYS	ENGINEERED	UNP P03069
A	28	HIS	LYS	ENGINEERED	UNP P03069
A	29	GLU	LEU	ENGINEERED	UNP P03069
A	30	CYS	VAL	ENGINEERED	UNP P03069

- Molecule 2 is a protein called GENERAL CONTROL PROTEIN GCN4.

Mol	Chain	Residues	Atoms						Trace
2	B	33	Total	C	H	N	O	S	1
			530	156	279	54	40	1	

There are 23 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	GLU	ARG	ENGINEERED	UNP P03069
B	2	VAL	MET	ENGINEERED	UNP P03069

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	3	GLN	LYS	ENGINEERED	UNP P03069
B	4	ALA	GLN	ENGINEERED	UNP P03069
B	6	LYS	GLU	ENGINEERED	UNP P03069
B	7	LYS	ASP	ENGINEERED	UNP P03069
B	8	ARG	LYS	ENGINEERED	UNP P03069
B	10	GLN	GLU	ENGINEERED	UNP P03069
B	11	ALA	GLU	ENGINEERED	UNP P03069
B	13	LYS	LEU	ENGINEERED	UNP P03069
B	14	ALA	SER	ENGINEERED	UNP P03069
B	15	ARG	LYS	ENGINEERED	UNP P03069
B	18	ALA	HIS	ENGINEERED	UNP P03069
B	19	ALA	LEU	ENGINEERED	UNP P03069
B	20	LYS	GLU	ENGINEERED	UNP P03069
B	21	GLN	ASN	ENGINEERED	UNP P03069
B	22	LYS	GLU	ENGINEERED	UNP P03069
B	24	GLN	ALA	ENGINEERED	UNP P03069
B	25	ALA	ARG	ENGINEERED	UNP P03069
B	27	ARG	LYS	ENGINEERED	UNP P03069
B	28	HIS	LYS	ENGINEERED	UNP P03069
B	29	LYS	LEU	ENGINEERED	UNP P03069
B	30	CYS	VAL	ENGINEERED	UNP P03069

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: GENERAL CONTROL PROTEIN GCN4




- Molecule 2: GENERAL CONTROL PROTEIN GCN4




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 14. Colouring as in section 4.1 above.

- Molecule 1: GENERAL CONTROL PROTEIN GCN4



- Molecule 2: GENERAL CONTROL PROTEIN GCN4



5 Refinement protocol and experimental data overview

The models were refined using the following method: 1. *DISTANCE GEOMETRY/REGULARIZATION*, 2. *SIMULATED ANNEALING*, 3. *MOLECULAR DYNAMICS SIMULATION INCLUDING 8 ANGSTROMS WATER SHELL*.

Of the 50 calculated structures, 25 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
X-PLOR	refinement	3.851

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 4872
Number of chemical shift lists	2
Total number of shifts	456
Number of shifts mapped to atoms	456
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	48%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	216	201	201	4±2
2	B	237	267	267	5±2
All	All	11325	11700	11700	188

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 55 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:1:GLU:HB3	1:A:4:GLN:HG3	0.87	1.46	1	1
1:A:1:GLU:HB3	1:A:4:GLN:HG2	0.78	1.56	12	9
2:B:1:GLU:HB3	2:B:4:ALA:HB3	0.75	1.56	17	2
2:B:25:ALA:HA	2:B:28:HIS:CD2	0.69	2.22	17	3
1:A:15:GLU:O	1:A:19:LEU:HG	0.69	1.87	21	25

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	27/33 (82%)	27±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
2	B	29/33 (88%)	29±0 (100±1%)	0±0 (0±1%)	0±0 (0±0%)	100	100
All	All	1400/1650 (85%)	1397 (100%)	3 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	22/25 (88%)	22±0 (99±2%)	0±0 (1±2%)	83	97
2	B	23/24 (96%)	22±1 (94±4%)	1±1 (6±4%)	25	71
All	All	1125/1225 (92%)	1083 (96%)	42 (4%)	43	86

5 of 12 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	27	ARG	9
2	B	7	LYS	7
2	B	22	LYS	5
2	B	3	GLN	4
2	B	17	TYR	4

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 48% for the well-defined parts and 48% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 4872

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	216
Number of shifts mapped to atoms	216
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 22%, i.e. 161 atoms were assigned a chemical shift out of a possible 736. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	54/280 (19%)	54/112 (48%)	0/112 (0%)	0/56 (0%)
Sidechain	103/433 (24%)	103/252 (41%)	0/154 (0%)	0/27 (0%)
Aromatic	4/23 (17%)	4/12 (33%)	0/10 (0%)	0/1 (0%)
Overall	161/736 (22%)	161/376 (43%)	0/276 (0%)	0/84 (0%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: BMRB entry 4872

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	240
Number of shifts mapped to atoms	240
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 26%, i.e. 195 atoms were assigned a chemical shift out of a possible 736. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	58/280 (21%)	58/112 (52%)	0/112 (0%)	0/56 (0%)
Sidechain	131/433 (30%)	131/252 (52%)	0/154 (0%)	0/27 (0%)
Aromatic	6/23 (26%)	6/12 (50%)	0/10 (0%)	0/1 (0%)
Overall	195/736 (26%)	195/376 (52%)	0/276 (0%)	0/84 (0%)

7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B:

