



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 11:31 pm GMT

PDB ID : 4FNQ  
Title : Crystal structure of GH36 alpha-galactosidase AgaB from *Geobacillus stearothermophilus*  
Authors : Merceron, R.; Foucault, M.; Haser, R.; Mattes, R.; Watzlawick, H.; Gouet, P.  
Deposited on : 2012-06-20  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	729	<div> <div>9%</div> <div>87%</div> <div>11%</div> <div>..</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6232 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-galactosidase AgaB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	723	Total	C	N	O	S	0	0	0
			5832	3704	1025	1078	25			

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	396	Total	O	0	0
			396	396		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Alpha-galactosidase AgaB



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.30Å 113.10Å 161.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.47 – 1.80 19.47 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.47-1.80) 98.8 (19.47-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.78 (at 1.80Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.181 , 0.209 0.180 , 0.208	Depositor DCC
$R_{free}$ test set	3658 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.6	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 55.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6232	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/5982	0.59	2/8109 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	585	ARG	N-CA-CB	-5.98	99.84	110.60
1	A	374	ASP	N-CA-CB	-5.38	100.92	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5832	0	5658	55	0
2	A	4	0	6	0	0
3	A	396	0	0	2	0
All	All	6232	0	5664	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:GLU:HB3	1:A:629:ARG:HD3	1.50	0.93
1:A:1:MET:SD	3:A:1293:HOH:O	2.31	0.88
1:A:416:MET:SD	1:A:445:GLN:NE2	2.61	0.72
1:A:361:GLU:OE1	1:A:629:ARG:NH2	2.22	0.71
1:A:594:HIS:HA	1:A:721:MET:HE1	1.73	0.68
1:A:1:MET:SD	1:A:154:GLU:HG2	2.35	0.67
1:A:457:ASP:OD1	1:A:509:ARG:NH2	2.24	0.66
1:A:413:GLU:HG2	1:A:416:MET:HG3	1.80	0.63
1:A:376:ASP:HB3	1:A:443:ARG:HH21	1.65	0.61
1:A:373:ARG:O	1:A:373:ARG:HG3	2.01	0.60
1:A:351:ILE:HG23	1:A:607:ILE:HD11	1.84	0.60
1:A:337:GLU:H	1:A:337:GLU:CD	2.05	0.59
1:A:375:ASP:OD2	1:A:377:ARG:NH1	2.36	0.59
1:A:692:ILE:HG12	1:A:724:LEU:HG	1.85	0.59
1:A:587:ALA:O	1:A:592:ARG:NH1	2.36	0.58
1:A:1:MET:N	1:A:15:ARG:O	2.30	0.57
1:A:336:TRP:HZ3	1:A:367:ASP:HB3	1.69	0.57
1:A:422:GLU:OE2	1:A:425:ARG:NH2	2.38	0.57
1:A:551:ASP:OD1	1:A:585:ARG:HB3	2.04	0.57
1:A:1:MET:HB3	1:A:154:GLU:CA	2.38	0.54
1:A:551:ASP:OD2	1:A:585:ARG:NH1	2.34	0.53
1:A:552:ALA:HB2	1:A:592:ARG:HG2	1.90	0.53
1:A:339:THR:HB	1:A:342:ASP:O	2.10	0.52
1:A:383:TRP:HB2	1:A:462:THR:HG22	1.92	0.51
1:A:662:LEU:HD11	1:A:721:MET:HG2	1.93	0.50
1:A:367:ASP:OD1	1:A:368:GLY:N	2.45	0.49
1:A:1:MET:O	1:A:131:GLU:HB2	2.13	0.49
1:A:343:PHE:HZ	1:A:369:TRP:HA	1.77	0.49
1:A:592:ARG:HG3	1:A:592:ARG:HH11	1.77	0.49
1:A:346:GLU:HG3	1:A:347:LYS:N	2.27	0.48
1:A:400:GLN:HA	1:A:403:GLU:HG2	1.95	0.48
1:A:578:VAL:HG11	1:A:586:VAL:HA	1.95	0.48
1:A:413:GLU:HG3	1:A:478:ASP:O	2.14	0.48
1:A:1:MET:HB3	1:A:154:GLU:HA	1.95	0.48
1:A:578:VAL:HA	1:A:579:PRO:C	2.37	0.46
1:A:338:ALA:HA	1:A:606:ASP:CG	2.36	0.45
1:A:582:GLN:OE1	3:A:1097:HOH:O	2.20	0.45
1:A:70:ASP:HB2	1:A:82:THR:O	2.16	0.45
1:A:591:ALA:HB1	1:A:717:PHE:CE2	2.53	0.44
1:A:421:SER:O	1:A:425:ARG:HG3	2.18	0.44
1:A:551:ASP:OD2	1:A:585:ARG:HD2	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:594:HIS:HA	1:A:721:MET:CE	2.44	0.43
1:A:336:TRP:CZ3	1:A:367:ASP:HB3	2.52	0.43
1:A:341:PHE:CE2	1:A:375:ASP:HA	2.53	0.43
1:A:362:LEU:HD21	1:A:474:TYR:CG	2.55	0.42
1:A:693:GLU:HG2	1:A:723:ARG:H	1.83	0.42
1:A:590:LYS:HE2	1:A:594:HIS:NE2	2.34	0.42
1:A:1:MET:HA	1:A:152:ALA:O	2.19	0.42
1:A:442:GLY:O	1:A:445:GLN:HG2	2.19	0.42
1:A:411:TRP:HD1	1:A:476:LYS:HE3	1.85	0.42
1:A:560:TYR:CE2	1:A:639:ARG:HD2	2.56	0.41
1:A:1:MET:HB3	1:A:154:GLU:CB	2.50	0.41
1:A:1:MET:HB3	1:A:154:GLU:HB2	2.02	0.41
1:A:345:GLU:HA	1:A:348:LEU:HD12	2.02	0.40
1:A:434:VAL:HG23	1:A:437:ARG:HB2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	719/729 (99%)	685 (95%)	31 (4%)	3 (0%)	38 23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	342	ASP
1	A	225	SER
1	A	579	PRO



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	614/620 (99%)	601 (98%)	13 (2%)	59 46

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	51	ARG
1	A	215	GLN
1	A	334	ASN
1	A	350	ASN
1	A	380	LEU
1	A	387	ARG
1	A	410	LEU
1	A	416	MET
1	A	420	ASN
1	A	426	LYS
1	A	533	PHE
1	A	612	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	420	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	801	-	3,3,3	0.59	0	2,2,2	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	801	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	723/729 (99%)	0.30	69 (9%) 9 7	8, 16, 56, 87	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	340	TYR	14.3
1	A	586	VAL	13.8
1	A	583	VAL	11.6
1	A	374	ASP	11.6
1	A	581	HIS	10.0
1	A	1	MET	9.2
1	A	584	GLY	9.1
1	A	338	ALA	9.0
1	A	579	PRO	8.6
1	A	580	ASN	8.5
1	A	336	TRP	8.2
1	A	372	LYS	7.9
1	A	420	ASN	7.3
1	A	694	GLY	7.2
1	A	376	ASP	7.0
1	A	578	VAL	7.0
1	A	585	ARG	6.9
1	A	341	PHE	6.6
1	A	582	GLN	6.3
1	A	387	ARG	6.3
1	A	350	ASN	6.3
1	A	377	ARG	6.0
1	A	343	PHE	5.7
1	A	371	GLY	5.6
1	A	342	ASP	5.5
1	A	339	THR	5.5
1	A	388	ARG	5.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	347	LYS	5.2
1	A	337	GLU	5.1
1	A	392	ASN	5.1
1	A	391	PRO	5.0
1	A	386	ASN	4.7
1	A	695	LEU	4.3
1	A	375	ASP	4.2
1	A	384	ILE	4.2
1	A	587	ALA	4.2
1	A	378	ARG	4.1
1	A	712	TYR	4.0
1	A	385	VAL	3.9
1	A	346	GLU	3.8
1	A	44	ARG	3.8
1	A	373	ARG	3.6
1	A	353	LYS	3.6
1	A	403	GLU	3.6
1	A	608	THR	3.6
1	A	612	GLU	3.5
1	A	425	ARG	3.4
1	A	613	THR	3.4
1	A	50	SER	3.4
1	A	390	LEU	3.3
1	A	607	ILE	3.3
1	A	393	GLY	3.0
1	A	588	SER	3.0
1	A	63	SER	3.0
1	A	65	ARG	2.9
1	A	422	GLU	2.8
1	A	443	ARG	2.7
1	A	577	ALA	2.6
1	A	382	ASP	2.6
1	A	367	ASP	2.6
1	A	438	PRO	2.5
1	A	357	GLU	2.5
1	A	370	PHE	2.4
1	A	51	ARG	2.3
1	A	383	TRP	2.2
1	A	354	THR	2.1
1	A	713	ARG	2.1
1	A	369	TRP	2.1
1	A	444	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	EDO	A	801	4/4	0.97	0.10	0.88	11,11,12,13	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.