



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2018 – 08:50 AM EST

PDB ID : 4FRA
Title : Crystal Structure of ABBA+UDP+Gal at pH 5.0 with MPD as the cryoprotectant
Authors : Johal, A.R.; Alfaro, J.A.; Blackler, R.J.; Schuman, B.; Borisova, S.N.; Evans, S.V.
Deposited on : 2012-06-26
Resolution : 1.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1367 (1.46-1.42)
Clashscore	112137	1425 (1.46-1.42)
Ramachandran outliers	110173	1405 (1.46-1.42)
Sidechain outliers	110143	1405 (1.46-1.42)
RSRZ outliers	101464	1372 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UDP	A	401	-	-	-	X
3	GAL	A	402	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4942 atoms, of which 2323 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histo-blood group ABO system transferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	272	4553	1470	2294	387	388	14	674	7	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	62	PHE	-	EXPRESSION TAG	UNP H6A2X0
A	63	MET	-	EXPRESSION TAG	UNP H6A2X0
A	176	ARG	GLY	ENGINEERED MUTATION	UNP H6A2X0
A	268	GLY	ALA	ENGINEERED MUTATION	UNP H6A2X0
A	355	GLU	-	EXPRESSION TAG	UNP H6A2X0

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C₉H₁₄N₂O₁₂P₂).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
2	A	1	36	9	11	2	12	2	2	0

- Molecule 3 is BETA-D-GALACTOSE (three-letter code: GAL) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	4	0
			24	6	12	6		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	1	0
			13	4	6	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	316	Total 316	O 316	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histo-blood group ABO system transferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	52.56 Å 149.67 Å 79.32 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.43 19.81 – 1.43	Depositor EDS
% Data completeness (in resolution range)	94.8 (20.00-1.43) 94.8 (19.81-1.43)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 1.43 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.183 , 0.208 0.186 , 0.209	Depositor DCC
R_{free} test set	2804 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.149	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4942	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, PEG, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.34	7/2348 (0.3%)	1.46	32/3183 (1.0%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	135	GLU	CD-OE1	7.24	1.33	1.25
1	A	282	ARG	CZ-NH1	6.96	1.42	1.33
1	A	332	TRP	CD2-CE2	5.82	1.48	1.41
1	A	110	ARG	CZ-NH2	5.47	1.40	1.33
1	A	126	TYR	CE1-CZ	-5.42	1.31	1.38
1	A	260	GLU	CG-CD	-5.35	1.44	1.51
1	A	213	ASP	CB-CG	-5.11	1.41	1.51

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	ARG	NE-CZ-NH1	15.64	128.12	120.30
1	A	282	ARG	NE-CZ-NH2	-15.13	112.73	120.30
1	A	217	ARG	NE-CZ-NH1	-11.51	114.55	120.30
1	A	213	ASP	CB-CG-OD1	10.65	127.89	118.30
1	A	211	ASP	CB-CG-OD2	-9.92	109.38	118.30
1	A	312	ARG	NE-CZ-NH1	-9.82	115.39	120.30
1	A	217	ARG	NE-CZ-NH2	9.70	125.15	120.30
1	A	259	ASP	CB-CG-OD1	9.10	126.48	118.30
1	A	244	PHE	CB-CG-CD2	8.89	127.03	120.80
1	A	340	ARG	NE-CZ-NH1	8.85	124.72	120.30
1	A	260	GLU	OE1-CD-OE2	-8.45	113.16	123.30
1	A	97	GLU	OE1-CD-OE2	-8.39	113.23	123.30
1	A	168	ARG	NE-CZ-NH2	-8.23	116.18	120.30
1	A	211	ASP	CB-CG-OD1	7.96	125.47	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	154	ASP	CB-CG-OD1	7.90	125.41	118.30
1	A	135	GLU	OE1-CD-OE2	7.70	132.54	123.30
1	A	269	PHE	CB-CG-CD2	6.98	125.69	120.80
1	A	120[A]	VAL	CG1-CB-CG2	-6.42	100.63	110.90
1	A	120[B]	VAL	CG1-CB-CG2	-6.42	100.63	110.90
1	A	63	MET	CG-SD-CE	6.05	109.88	100.20
1	A	244	PHE	CB-CG-CD1	-5.98	116.62	120.80
1	A	255	TYR	CB-CG-CD1	5.88	124.53	121.00
1	A	217	ARG	CB-CG-CD	5.83	126.76	111.60
1	A	213	ASP	CB-CG-OD2	-5.76	113.11	118.30
1	A	326	ASP	CB-CG-OD2	-5.74	113.14	118.30
1	A	189	MET	CG-SD-CE	5.68	109.29	100.20
1	A	264	TYR	CB-CG-CD1	5.53	124.31	121.00
1	A	278	GLN	CB-CG-CD	5.49	125.86	111.60
1	A	329	LEU	CB-CG-CD1	5.46	120.28	111.00
1	A	76	VAL	N-CA-CB	-5.32	99.79	111.50
1	A	154	ASP	OD1-CG-OD2	-5.31	113.21	123.30
1	A	218	ASP	CB-CG-OD2	5.11	122.89	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2259	2294	2266	40	0
2	A	25	11	11	1	0
3	A	12	12	12	0	0
4	A	7	6	10	2	0
5	A	316	0	0	14	2
All	All	2619	2323	2299	42	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:MET:HE2	1:A:324:LEU:HG	1.23	1.12
1:A:266:MET:HE2	1:A:324:LEU:CG	1.89	1.02
1:A:302:ASP:OD2	5:A:722:HOH:O	1.99	0.81
1:A:197:GLU:HA	1:A:278:GLN:OE1	1.80	0.80
1:A:214[A]:MET:HE2	5:A:740:HOH:O	1.83	0.78
1:A:63:MET:HE2	5:A:723:HOH:O	1.86	0.76
2:A:401:UDP:O1A	5:A:816:HOH:O	2.06	0.73
1:A:75:LYS:HB2	1:A:78[A]:THR:HG23	1.73	0.71
1:A:112:GLN:NE2	5:A:745:HOH:O	2.29	0.66
1:A:345:PRO:HD2	5:A:784:HOH:O	1.98	0.64
1:A:266:MET:HE2	1:A:324:LEU:CD2	2.27	0.64
1:A:123:ILE:CD1	1:A:153:THR:HA	2.29	0.62
1:A:120[A]:VAL:HG21	1:A:133:PHE:CZ	2.35	0.62
1:A:101:ASN:C	1:A:101:ASN:HD22	2.06	0.59
1:A:266:MET:CE	1:A:324:LEU:CD2	2.82	0.57
1:A:266:MET:CE	1:A:324:LEU:HD21	2.34	0.57
1:A:193:SER:HA	1:A:278:GLN:NE2	2.20	0.57
1:A:305:HIS:HE1	5:A:550:HOH:O	1.87	0.56
1:A:110:ARG:HH22	1:A:113:ASN:HD22	1.54	0.56
4:A:403:PEG:H22	5:A:811:HOH:O	2.05	0.56
1:A:106:ASN:OD1	1:A:145:HIS:HE1	1.90	0.55
1:A:217:ARG:HD3	1:A:342:THR:OG1	2.07	0.55
1:A:140:HIS:HA	4:A:403:PEG:H12	1.89	0.55
1:A:123:ILE:HD13	1:A:153:THR:HA	1.90	0.54
1:A:285:HIS:HA	1:A:288:MET:HE3	1.90	0.54
1:A:101:ASN:HD21	1:A:103:ASP:HB2	1.74	0.53
1:A:120[A]:VAL:HG11	1:A:212:VAL:HG23	1.91	0.52
1:A:62:PHE:N	5:A:706:HOH:O	2.42	0.52
1:A:266:MET:HE1	1:A:324:LEU:HD21	1.92	0.52
1:A:305:HIS:HD2	5:A:542:HOH:O	1.92	0.51
1:A:120[A]:VAL:CG1	1:A:212:VAL:HB	2.44	0.48
1:A:242:GLU:CD	1:A:242:GLU:H	2.17	0.47
1:A:120[A]:VAL:HG21	1:A:133:PHE:CE2	2.50	0.46
1:A:80:CYS:SG	5:A:566:HOH:O	2.23	0.46
1:A:145:HIS:HD2	5:A:512:HOH:O	1.98	0.45
1:A:110:ARG:HH22	1:A:113:ASN:ND2	2.13	0.45
1:A:125:LYS:H	1:A:125:LYS:HE2	1.84	0.43
1:A:297:GLU:OE2	1:A:301:HIS:HD2	2.02	0.43
1:A:285:HIS:HD2	5:A:711:HOH:O	2.01	0.43
1:A:123:ILE:HD12	1:A:154:ASP:H	1.83	0.42
1:A:80:CYS:HB3	1:A:98:GLY:O	2.19	0.41
1:A:214[A]:MET:CE	5:A:740:HOH:O	2.54	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:579:HOH:O	5:A:647:HOH:O[3_455]	2.09	0.11
5:A:647:HOH:O	5:A:647:HOH:O[3_455]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/294 (93%)	267 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/263 (95%)	241 (96%)	9 (4%)	40	7

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78[A]	THR
1	A	78[B]	THR
1	A	101	ASN
1	A	113	ASN
1	A	187	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	269	PHE
1	A	278	GLN
1	A	324	LEU
1	A	329	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
1	A	113	ASN
1	A	145	HIS
1	A	275	GLN
1	A	285	HIS
1	A	286	GLN
1	A	294	ASN
1	A	301	HIS
1	A	305	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	UDP	A	401	-	20,26,26	2.53	6 (30%)	19,40,40	2.12	8 (42%)
3	GAL	A	402	-	12,12,12	1.57	3 (25%)	17,17,17	1.67	5 (29%)
4	PEG	A	403	-	6,6,6	1.00	0	5,5,5	3.14	4 (80%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	401	-	-	0/12/32/32	0/2/2/2
3	GAL	A	402	-	-	0/2/22/22	0/1/1/1
4	PEG	A	403	-	-	0/4/4/4	0/0/0/0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	UDP	C3'-C2'	-2.53	1.46	1.53
3	A	402	GAL	O5-C5	-2.19	1.39	1.44
3	A	402	GAL	O3-C3	-2.14	1.38	1.43
2	A	401	UDP	O3'-C3'	2.59	1.48	1.43
2	A	401	UDP	PB-O1B	3.21	1.61	1.50
2	A	401	UDP	C4-N3	3.60	1.41	1.36
3	A	402	GAL	O5-C1	3.72	1.49	1.43
2	A	401	UDP	C6-N1	5.86	1.43	1.35
2	A	401	UDP	O4'-C1'	6.13	1.49	1.41

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	UDP	O2'-C2'-C3'	-4.84	96.31	111.83
2	A	401	UDP	O4'-C1'-N1	-4.17	99.73	108.08
2	A	401	UDP	O4'-C4'-C3'	-3.09	99.03	105.17
2	A	401	UDP	O3'-C3'-C2'	-2.76	102.97	111.83
3	A	402	GAL	C1-O5-C5	-2.74	108.44	113.39
3	A	402	GAL	O1-C1-O5	-2.44	102.98	110.20
2	A	401	UDP	O2B-PB-O1B	-2.26	101.64	110.50
3	A	402	GAL	O2-C2-C1	-2.04	105.51	109.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	UDP	C4'-O4'-C1'	-2.02	107.62	109.77
2	A	401	UDP	C2'-C3'-C4'	2.02	106.55	102.62
2	A	401	UDP	O2'-C2'-C1'	2.15	118.36	111.61
3	A	402	GAL	O1-C1-C2	2.36	116.09	109.42
4	A	403	PEG	O2-C3-C4	2.41	121.29	110.15
4	A	403	PEG	C3-O2-C2	2.56	124.40	113.30
3	A	402	GAL	O5-C5-C4	3.64	116.36	109.66
4	A	403	PEG	O2-C2-C1	3.90	128.16	110.15
4	A	403	PEG	O1-C1-C2	4.31	136.65	111.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	UDP	1	0
4	A	403	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	272/294 (92%)	0.61	22 (8%) 13 14	10, 17, 34, 69	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	176	ARG	6.3
1	A	345	PRO	6.2
1	A	80	CYS	6.2
1	A	191	MET	4.9
1	A	161	ARG	4.5
1	A	292	GLN	3.7
1	A	187	ARG	3.5
1	A	192	ILE	3.4
1	A	256	ILE	3.2
1	A	198	ARG	3.1
1	A	120[A]	VAL	2.9
1	A	344	VAL	2.9
1	A	62	PHE	2.8
1	A	78[A]	THR	2.6
1	A	64	VAL	2.5
1	A	151	VAL	2.5
1	A	196	CYS	2.4
1	A	202	SER	2.4
1	A	190	GLU	2.3
1	A	294	ASN	2.2
1	A	87[A]	VAL	2.2
1	A	125	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	UDP	A	401	25/25	0.59	0.29	2.80	16,26,36,39	11
3	GAL	A	402	12/12	0.93	0.12	2.32	8,16,21,21	16
4	PEG	A	403	7/7	0.82	0.20	1.90	19,34,42,42	1

6.5 Other polymers [i](#)

There are no such residues in this entry.