



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:28 pm GMT

PDB ID : 3FS8  
Title : Crystal structure of QdtC, the dTDP-3-amino-3,6-dideoxy-D-glucose N-acetyl transferase from *Thermoanaerobacterium thermosaccharolyticum* in complex with Acetyl-CoA  
Authors : Holden, H.M.; Thoden, J.B.  
Deposited on : 2009-01-09  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	
1	B	273	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TDR	A	301	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called QdtC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	2	0
			2058	1320	354	381	3			
1	B	257	Total	C	N	O	S	0	2	0
			2042	1310	350	379	3			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	266	LEU	-	EXPRESSION TAG	UNP Q6TFC6
A	267	GLU	-	EXPRESSION TAG	UNP Q6TFC6
A	268	HIS	-	EXPRESSION TAG	UNP Q6TFC6
A	269	HIS	-	EXPRESSION TAG	UNP Q6TFC6
A	270	HIS	-	EXPRESSION TAG	UNP Q6TFC6
A	271	HIS	-	EXPRESSION TAG	UNP Q6TFC6
A	272	HIS	-	EXPRESSION TAG	UNP Q6TFC6
A	273	HIS	-	EXPRESSION TAG	UNP Q6TFC6
B	266	LEU	-	EXPRESSION TAG	UNP Q6TFC6
B	267	GLU	-	EXPRESSION TAG	UNP Q6TFC6
B	268	HIS	-	EXPRESSION TAG	UNP Q6TFC6
B	269	HIS	-	EXPRESSION TAG	UNP Q6TFC6
B	270	HIS	-	EXPRESSION TAG	UNP Q6TFC6
B	271	HIS	-	EXPRESSION TAG	UNP Q6TFC6
B	272	HIS	-	EXPRESSION TAG	UNP Q6TFC6
B	273	HIS	-	EXPRESSION TAG	UNP Q6TFC6

- Molecule 2 is ACETYL COENZYME \*A (three-letter code: ACO) (formula: C<sub>23</sub>H<sub>38</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		

- Molecule 3 is THYMINE (three-letter code: TDR) (formula: C<sub>5</sub>H<sub>6</sub>N<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	5	2	2		
3	B	1	Total	C	N	O	0	0
			9	5	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	230	Total 230	O 230	0	0
4	B	268	Total 268	O 268	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: QdtC



#### • Molecule 1: QdtC



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.30Å 67.30Å 112.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.70 40.46 – 1.70	Depositor EDS
% Data completeness (in resolution range)	93.9 (30.00-1.70) 94.1 (40.46-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.67 (at 1.70Å)	Xtriage
Refinement program	TNT	Depositor
R, $R_{free}$	0.174 , 0.241 0.174 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	9.8	Xtriage
Anisotropy	0.381	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 86.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for -h,-k,l 0.054 for h,-h-k,-l 0.030 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TDR, ACO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.76	0/2112	1.23	7/2872 (0.2%)
1	B	0.79	0/2093	1.22	5/2846 (0.2%)
All	All	0.78	0/4205	1.23	12/5718 (0.2%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	50	ARG	NE-CZ-NH2	-10.09	115.25	120.30
1	B	50	ARG	NE-CZ-NH1	9.92	125.26	120.30
1	A	59	ASP	CB-CG-OD1	6.66	124.30	118.30
1	A	50	ARG	NE-CZ-NH2	-6.22	117.19	120.30
1	B	25	ASP	CB-CG-OD1	6.06	123.75	118.30
1	A	80	ARG	NE-CZ-NH1	5.83	123.21	120.30
1	A	93	ASP	CB-CG-OD2	5.63	123.37	118.30
1	A	37	ASP	CB-CG-OD1	5.40	123.16	118.30
1	A	63	ASP	CB-CG-OD1	5.37	123.13	118.30
1	A	96	GLN	CB-CA-C	-5.30	99.81	110.40
1	B	247	MET	CG-SD-CE	-5.26	91.79	100.20
1	B	175	LEU	CB-CG-CD1	-5.25	102.08	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2058	0	2079	28	1
1	B	2042	0	2065	41	0
2	A	51	0	34	6	0
2	B	51	0	34	0	0
3	A	9	0	6	0	0
3	B	9	0	6	0	0
4	A	230	0	0	3	3
4	B	268	0	0	4	2
All	All	4718	0	4224	71	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:123:HIS:HD2	1:B:124:HIS:HD2	1.21	0.88
1:A:153:PRO:HG2	2:A:300:ACO:HH31	1.58	0.86
1:B:257:TRP:CZ3	1:B:258:ILE:HD13	2.21	0.76
1:B:232:THR:HG23	1:B:234:GLU:H	1.52	0.73
1:B:123:HIS:CD2	1:B:124:HIS:HD2	2.07	0.71
1:A:153:PRO:HD2	2:A:300:ACO:HH32	1.74	0.70
1:B:7:LYS:NZ	4:B:531:HOH:O	2.24	0.70
1:A:216:ASN:HD22	1:A:216:ASN:C	1.95	0.68
1:B:123:HIS:HD2	1:B:124:HIS:CD2	2.07	0.68
1:B:252:THR:HG21	1:B:256:THR:HG21	1.77	0.65
1:A:153:PRO:HG2	2:A:300:ACO:CH3	2.25	0.64
1:B:62:ASN:ND2	4:B:516:HOH:O	2.30	0.64
1:A:130:TYR:CE2	1:A:248:PRO:HD3	2.33	0.63
1:B:229:ASN:HB3	1:B:232:THR:HG22	1.80	0.62
2:A:300:ACO:O1A	2:A:300:ACO:O5A	2.18	0.62
1:A:172:THR:HB	1:A:190:HIS:HD2	1.65	0.62
1:A:237:TYR:HA	1:A:238:PRO:C	2.21	0.60
1:A:153:PRO:CD	2:A:300:ACO:HH32	2.31	0.60
1:A:117:THR:HG22	1:A:118:LEU:HG	1.83	0.60
1:B:232:THR:HG23	1:B:234:GLU:HG2	1.84	0.59
1:A:42:LYS:HG3	1:A:73:ILE:O	2.02	0.59
1:B:78:LEU:HB3	1:B:96:GLN:HG2	1.84	0.59
1:B:43:LYS:HG2	1:B:75:GLU:HG2	1.85	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:220:GLU:OE2	4:A:433:HOH:O	2.17	0.57
2:A:300:ACO:H51A	2:A:300:ACO:H8A	1.87	0.57
1:B:141:GLU:HG2	1:B:142:LYS:HG3	1.88	0.56
1:B:216:ASN:C	1:B:216:ASN:HD22	2.08	0.55
1:B:237:TYR:HA	1:B:238:PRO:C	2.26	0.55
1:B:224:ILE:HD12	1:B:237:TYR:HB3	1.89	0.54
1:B:247:MET:HG3	1:B:249:TRP:CZ2	2.44	0.53
1:A:194:ASP:O	1:A:224:ILE:HD12	2.09	0.53
1:A:256:THR:HA	1:A:259:LYS:HE3	1.91	0.53
1:B:229:ASN:CB	1:B:232:THR:HG22	2.38	0.53
1:A:130:TYR:CZ	1:A:248:PRO:HD3	2.44	0.52
1:B:141:GLU:O	1:B:142:LYS:HB2	2.11	0.51
1:A:141:GLU:HG2	1:A:142:LYS:HG3	1.92	0.50
1:A:216:ASN:ND2	1:A:216:ASN:C	2.66	0.49
1:A:219:ARG:NH2	4:A:353:HOH:O	2.26	0.49
1:B:232:THR:CG2	1:B:234:GLU:HG2	2.43	0.49
1:B:67:LYS:N	1:B:67:LYS:HD3	2.28	0.49
1:B:229:ASN:O	1:B:233:GLY:N	2.40	0.48
1:A:141:GLU:O	1:A:142:LYS:HB2	2.14	0.48
1:B:257:TRP:O	1:B:258:ILE:C	2.51	0.48
1:B:108:LYS:HE2	1:B:108:LYS:HB3	1.73	0.48
1:B:224:ILE:HG13	1:B:225:ARG:N	2.30	0.47
1:A:114:LYS:HE2	1:A:132:ASN:OD1	2.15	0.47
1:B:258:ILE:HA	1:B:258:ILE:HD12	1.54	0.47
1:A:67:LYS:N	1:A:67:LYS:HD3	2.30	0.46
1:A:219:ARG:NH2	4:A:461:HOH:O	2.32	0.46
1:A:123:HIS:HD2	1:A:124:HIS:HD2	1.65	0.45
1:B:152:PHE:HB3	1:B:153:PRO:HD2	1.97	0.45
1:A:126:TYR:C	1:A:126:TYR:CD2	2.91	0.44
1:B:152:PHE:HB3	1:B:153:PRO:CD	2.48	0.44
1:B:214:VAL:O	1:B:218:ALA:HA	2.18	0.44
1:A:221:ILE:O	1:A:222:CYS:HB3	2.18	0.43
1:B:255:ASP:O	1:B:258:ILE:HG22	2.18	0.43
1:B:213:VAL:HG12	1:B:220:GLU:HA	2.00	0.43
1:A:152:PHE:HB3	1:A:153:PRO:CD	2.49	0.42
1:A:226:LYS:HG2	1:A:226:LYS:O	2.18	0.42
1:A:209:LYS:HG3	1:A:210:GLU:HG3	2.01	0.42
1:B:207:VAL:HA	1:B:208:PRO:HD3	1.86	0.42
1:B:190:HIS:ND1	4:B:446:HOH:O	2.14	0.42
1:B:19:GLU:HG3	1:B:20[A]:ASN:OD1	2.20	0.42
1:B:219:ARG:CG	4:B:519:HOH:O	2.67	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:LYS:HG3	1:A:174:GLU:HA	2.03	0.41
1:B:123:HIS:CD2	1:B:124:HIS:CD2	2.95	0.41
1:B:114:LYS:HD2	1:B:132:ASN:HD21	1.85	0.41
1:B:249:TRP:HB3	1:B:257:TRP:CB	2.51	0.41
1:B:117:THR:HG22	1:B:118:LEU:HG	2.03	0.40
1:B:252:THR:CG2	1:B:256:THR:HG21	2.50	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:276:HOH:O	4:A:419:HOH:O[2_555]	2.05	0.15
1:A:96:GLN:OE1	4:A:494:HOH:O[3_555]	2.11	0.09
4:B:426:HOH:O	4:B:487:HOH:O[2_655]	2.14	0.06
4:A:338:HOH:O	4:B:531:HOH:O[2_655]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/273 (95%)	248 (96%)	11 (4%)	0	100	100
1	B	257/273 (94%)	249 (97%)	8 (3%)	0	100	100
All	All	516/546 (94%)	497 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/242 (95%)	219 (95%)	11 (5%)	30	11
1	B	228/242 (94%)	220 (96%)	8 (4%)	41	19
All	All	458/484 (95%)	439 (96%)	19 (4%)	34	14

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ILE
1	A	17	ILE
1	A	29	ILE
1	A	43	LYS
1	A	67	LYS
1	A	83	ASN
1	A	146	LYS
1	A	216	ASN
1	A	219	ARG
1	A	251	GLU
1	A	261	ILE
1	B	13	GLU
1	B	28	TYR
1	B	83	ASN
1	B	108	LYS
1	B	189	ILE
1	B	216	ASN
1	B	252	THR
1	B	258	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	38	ASN
1	A	94	ASN
1	A	124	HIS
1	A	216	ASN
1	B	62	ASN
1	B	94	ASN
1	B	112	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	123	HIS
1	B	124	HIS
1	B	216	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACO	A	300	-	46,53,53	1.12	3 (6%)	53,79,79	1.92	9 (16%)
3	TDR	A	301	-	7,9,9	1.61	2 (28%)	7,12,12	5.46	5 (71%)
2	ACO	B	300	-	46,53,53	0.99	3 (6%)	53,79,79	1.82	10 (18%)
3	TDR	B	301	-	7,9,9	1.31	1 (14%)	7,12,12	5.33	5 (71%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACO	A	300	-	-	2/47/67/67	0/3/3/3
3	TDR	A	301	-	-	0/0/0/0	0/1/1/1
2	ACO	B	300	-	-	2/47/67/67	0/3/3/3
3	TDR	B	301	-	-	0/0/0/0	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	300	ACO	C8A-N7A	-2.48	1.30	1.34
3	A	301	TDR	C6-C5	-2.01	1.36	1.39
2	B	300	ACO	P3B-O9A	2.22	1.64	1.54
2	A	300	ACO	P3B-O3B	2.23	1.63	1.59
2	B	300	ACO	O4B-C1B	2.31	1.44	1.41
3	B	301	TDR	C4-N3	2.62	1.37	1.33
3	A	301	TDR	C4-N3	2.88	1.38	1.33
2	A	300	ACO	O4B-C1B	2.96	1.45	1.41
2	A	300	ACO	C2B-C1B	3.76	1.59	1.53

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	TDR	N1-C2-N3	-9.43	121.61	128.40
3	A	301	TDR	N1-C2-N3	-9.30	121.71	128.40
2	A	300	ACO	N3A-C2A-N1A	-8.74	121.24	128.86
3	A	301	TDR	C5-C6-N1	-7.99	118.51	125.26
2	B	300	ACO	N3A-C2A-N1A	-7.98	121.91	128.86
3	B	301	TDR	C5-C4-N3	-6.19	118.42	125.24
2	B	300	ACO	C7P-C6P-C5P	-5.35	103.61	112.22
3	B	301	TDR	C5-C6-N1	-5.14	120.92	125.26
3	A	301	TDR	C5-C4-N3	-5.11	119.61	125.24
2	A	300	ACO	C5B-C4B-C3B	-4.55	98.87	114.30
2	A	300	ACO	C7P-C6P-C5P	-3.98	105.82	112.22
2	B	300	ACO	C3P-C2P-S1P	-3.42	101.30	111.23
2	A	300	ACO	C4B-O4B-C1B	-3.19	106.37	109.77
2	A	300	ACO	C2B-C3B-C4B	-2.76	98.30	103.23
2	A	300	ACO	C3P-C2P-S1P	-2.51	103.95	111.23
2	A	300	ACO	C7P-N8P-C9P	-2.39	118.13	122.59
2	A	300	ACO	P2A-O6A-CCP	-2.35	106.73	121.57
2	B	300	ACO	O3B-C3B-C4B	-2.33	101.28	110.04
2	B	300	ACO	O6A-CCP-CBP	-2.25	106.92	110.55
2	B	300	ACO	C4A-C5A-N7A	-2.15	107.33	109.41
2	B	300	ACO	C2B-C3B-C4B	-2.14	99.40	103.23
2	B	300	ACO	O5P-C5P-N4P	-2.11	118.94	122.97

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	300	ACO	C2A-N1A-C6A	2.14	122.52	118.77
3	B	301	TDR	C6-N1-C2	2.52	119.40	115.36
2	A	300	ACO	CDP-CBP-CAP	2.70	113.51	108.82
2	B	300	ACO	CEP-CBP-CCP	3.15	112.99	108.37
3	A	301	TDR	C4-N3-C2	3.83	118.51	115.16
3	A	301	TDR	C6-N1-C2	4.11	121.94	115.36
3	B	301	TDR	C4-N3-C2	6.20	120.58	115.16

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	300	ACO	CH3-C-S1P-C2P
2	B	300	ACO	CH3-C-S1P-C2P
2	B	300	ACO	O-C-S1P-C2P
2	A	300	ACO	O-C-S1P-C2P

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	ACO	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/273 (94%)	-0.35	0 100 100	7, 17, 41, 65	0
1	B	257/273 (94%)	-0.46	3 (1%) 79 83	6, 13, 45, 60	0
All	All	516/546 (94%)	-0.41	3 (0%) 89 91	6, 15, 43, 65	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	258	ILE	2.2
1	B	231	ILE	2.1
1	B	233	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	TDR	A	301	9/9	0.88	0.14	2.78	16,31,100,100	0
3	TDR	B	301	9/9	0.91	0.10	1.33	17,29,52,100	0
2	ACO	A	300	51/51	0.92	0.10	0.32	5,27,100,100	0
2	ACO	B	300	51/51	0.95	0.07	-0.14	10,20,47,100	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.