



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 7, 2017 – 03:10 AM EST

PDB ID : 4FSL  
Title : Crystal structure of beta-site app-cleaving enzyme 1 (BACE-DB-MUT) complex with N-(N-(4- acetamido-3-chloro-5-methylbenzyl)carbamimidoyl)-3-(4-methoxyphenyl)-5-methyl-4-isothiazolecarboxamide  
Authors : Muckelbauer, J.K.  
Deposited on : unknown  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	412	<div> <div>4%</div> <div> <div></div> <div>83%</div> <div>10%</div> <div>6%</div> </div> </div>
1	B	412	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>7%</div> <div>6%</div> </div> </div>
1	D	412	<div> <div>3%</div> <div> <div></div> <div>80%</div> <div>12%</div> <div>6%</div> </div> </div>
1	E	412	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>12%</div> <div>6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	0VB	A	501	-	X	-	-
2	0VB	B	501	-	X	-	-
2	0VB	D	501	-	X	-	-
2	0VB	E	501	-	X	-	-
3	IOD	A	502	-	-	X	-
3	IOD	B	503	-	-	X	-
3	IOD	D	502	-	-	X	-
3	IOD	E	502	-	-	X	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	387	Total	C	N	O	S	0	0	0
			3036	1943	504	575	14			
1	B	387	Total	C	N	O	S	0	0	0
			3038	1945	505	574	14			
1	D	387	Total	C	N	O	S	0	0	0
			3042	1948	505	575	14			
1	E	387	Total	C	N	O	S	0	0	0
			3040	1946	505	575	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	LYS	ARG	CONFLICT	UNP P56817
A	44	LYS	ARG	CONFLICT	UNP P56817
A	441	THR	-	EXPRESSION TAG	UNP P56817
B	43	LYS	ARG	CONFLICT	UNP P56817
B	44	LYS	ARG	CONFLICT	UNP P56817
B	441	THR	-	EXPRESSION TAG	UNP P56817
D	43	LYS	ARG	CONFLICT	UNP P56817
D	44	LYS	ARG	CONFLICT	UNP P56817
D	441	THR	-	EXPRESSION TAG	UNP P56817
E	43	LYS	ARG	CONFLICT	UNP P56817
E	44	LYS	ARG	CONFLICT	UNP P56817
E	441	THR	-	EXPRESSION TAG	UNP P56817

- Molecule 2 is N-{N-[4-(acetylamino)-3-chloro-5-methylbenzyl]carbamimidoyl}-3-(4-methoxyphenyl)-5-methyl-1,2-thiazole-4-carboxamide (three-letter code: 0VB) (formula: C<sub>23</sub>H<sub>24</sub>ClN<sub>5</sub>O<sub>3</sub>S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 33	C 23	Cl 1	N 5	O 3	S 1	0	0
2	B	1	Total 33	C 23	Cl 1	N 5	O 3	S 1	0	0
2	D	1	Total 33	C 23	Cl 1	N 5	O 3	S 1	0	0
2	E	1	Total 33	C 23	Cl 1	N 5	O 3	S 1	0	0

- Molecule 3 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	3	Total 3 I	0	0
3	A	3	Total 3 I	0	0
3	D	3	Total 3 I	0	0
3	E	3	Total 3 I	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	215	Total 215 O	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	211	Total 211	O 211	0	0
4	D	182	Total 182	O 182	0	0
4	E	211	Total 211	O 211	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Beta-secretase 1




#### • Molecule 1: Beta-secretase 1




#### • Molecule 1: Beta-secretase 1



#### • Molecule 1: Beta-secretase 1







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.54Å 131.18Å 90.31Å 90.00° 97.61° 90.00°	Depositor
Resolution (Å)	50.00 – 2.50 39.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.5 (50.00-2.50) 96.6 (39.00-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.48 (at 2.51Å)	Xtriage
Refinement program	CNS, REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.224 , 0.277 0.223 , 0.274	Depositor DCC
$R_{free}$ test set	3377 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.1	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.003 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13119	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.87 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.4340e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, 0VB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/3114	0.52	0/4236
1	B	0.40	2/3116 (0.1%)	0.57	2/4237 (0.0%)
1	D	0.37	0/3120	0.53	0/4243
1	E	0.36	0/3118	0.52	0/4240
All	All	0.37	2/12468 (0.0%)	0.53	2/16956 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	80	ASP	CG-OD2	7.20	1.42	1.25
1	B	80	ASP	CB-CG	-5.01	1.41	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	80	ASP	CB-CG-OD1	12.36	129.42	118.30
1	B	80	ASP	CB-CG-OD2	-7.64	111.42	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3036	0	2933	25	0
1	B	3038	0	2939	17	0
1	D	3042	0	2953	30	0
1	E	3040	0	2944	26	0
2	A	33	0	24	4	0
2	B	33	0	24	3	0
2	D	33	0	24	2	0
2	E	33	0	24	4	0
3	A	3	0	0	2	0
3	B	3	0	0	3	0
3	D	3	0	0	3	0
3	E	3	0	0	3	0
4	A	215	0	0	5	0
4	B	211	0	0	3	0
4	D	182	0	0	2	0
4	E	211	0	0	1	0
All	All	13119	0	11865	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 99 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:118:PRO:HA	1:E:123:LYS:HB3	1.56	0.86
1:A:363:SER:HB3	4:A:667:HOH:O	1.75	0.85
1:E:173:GLU:OE2	1:E:243:ARG:NH1	2.21	0.74
1:D:359:ASP:OD2	1:D:360:VAL:HG13	1.90	0.72
1:A:365:ASP:HB2	4:A:667:HOH:O	1.93	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/412 (93%)	378 (98%)	7 (2%)	0	100	100
1	B	385/412 (93%)	374 (97%)	11 (3%)	0	100	100
1	D	385/412 (93%)	375 (97%)	9 (2%)	1 (0%)	44	66
1	E	385/412 (93%)	377 (98%)	8 (2%)	0	100	100
All	All	1540/1648 (93%)	1504 (98%)	35 (2%)	1 (0%)	55	76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	360	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/353 (93%)	319 (97%)	9 (3%)	50	77
1	B	328/353 (93%)	319 (97%)	9 (3%)	50	77
1	D	330/353 (94%)	309 (94%)	21 (6%)	20	38
1	E	329/353 (93%)	316 (96%)	13 (4%)	36	62
All	All	1315/1412 (93%)	1263 (96%)	52 (4%)	36	62

5 of 52 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	178	ASP
1	D	271	ASP
1	E	348	ILE
1	D	197	LEU
1	D	221	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 12 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	0VB	A	501	-	32,35,35	6.28	25 (78%)	41,49,49	3.87	24 (58%)
2	0VB	B	501	-	32,35,35	6.43	25 (78%)	41,49,49	4.06	23 (56%)
2	0VB	D	501	-	32,35,35	6.28	25 (78%)	41,49,49	3.76	22 (53%)
2	0VB	E	501	-	32,35,35	6.35	25 (78%)	41,49,49	3.70	22 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0VB	A	501	-	-	0/19/23/23	0/2/3/3
2	0VB	B	501	-	-	0/19/23/23	0/2/3/3
2	0VB	D	501	-	-	0/19/23/23	0/2/3/3
2	0VB	E	501	-	-	0/19/23/23	0/2/3/3

The worst 5 of 100 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	0VB	C11-N27	2.26	1.47	1.43
2	D	501	0VB	C11-N27	2.31	1.47	1.43
2	B	501	0VB	C11-N27	2.40	1.48	1.43
2	E	501	0VB	C11-N27	2.46	1.48	1.43
2	D	501	0VB	O31-C12	2.47	1.42	1.37

The worst 5 of 91 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	0VB	C06-C13-CL1	-6.31	108.52	118.50
2	B	501	0VB	O30-C17-N27	-6.10	115.14	123.06
2	B	501	0VB	O29-C16-C08	-6.01	111.59	120.88
2	D	501	0VB	O29-C16-C08	-5.95	111.68	120.88
2	B	501	0VB	C06-C13-CL1	-5.89	109.19	118.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	0VB	4	0
2	B	501	0VB	3	0
2	D	501	0VB	2	0
2	E	501	0VB	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	387/412 (93%)	0.09	15 (3%) 40 42	18, 25, 37, 43	0
1	B	387/412 (93%)	0.05	11 (2%) 53 56	18, 25, 36, 40	0
1	D	387/412 (93%)	0.06	12 (3%) 49 52	18, 26, 38, 42	0
1	E	387/412 (93%)	0.07	10 (2%) 56 59	18, 25, 36, 40	0
All	All	1548/1648 (93%)	0.07	48 (3%) 49 52	18, 25, 37, 43	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	360	VAL	7.5
1	B	362	THR	7.2
1	A	360	VAL	5.9
1	E	362	THR	5.5
1	D	361	ALA	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	0VB	A	501	33/33	0.93	0.20	1.05	21,22,23,24	0
2	0VB	B	501	33/33	0.92	0.20	0.90	20,22,26,27	0
2	0VB	D	501	33/33	0.93	0.18	0.74	18,20,23,24	0
2	0VB	E	501	33/33	0.93	0.20	0.59	20,22,25,26	0
3	IOD	A	504	1/1	0.97	0.05	-2.54	66,66,66,66	0
3	IOD	B	504	1/1	0.94	0.07	-2.55	78,78,78,78	0
3	IOD	E	504	1/1	0.98	0.04	-2.68	71,71,71,71	0
3	IOD	D	504	1/1	0.97	0.06	-2.76	80,80,80,80	0
3	IOD	E	503	1/1	0.97	0.10	-	99,99,99,99	0
3	IOD	E	502	1/1	0.98	0.04	-	69,69,69,69	0
3	IOD	A	503	1/1	0.94	0.14	-	135,135,135,135	0
3	IOD	A	502	1/1	0.95	0.13	-	97,97,97,97	0
3	IOD	B	502	1/1	0.97	0.06	-	76,76,76,76	0
3	IOD	D	502	1/1	0.98	0.07	-	74,74,74,74	0
3	IOD	B	503	1/1	0.80	0.13	-	142,142,142,142	0
3	IOD	D	503	1/1	0.89	0.10	-	111,111,111,111	0

## 6.5 Other polymers

There are no such residues in this entry.