



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2017 – 04:45 pm GMT

PDB ID : 5FUJ  
Title : Crystallization of a dimeric heme peroxxygenase from the fungus *Marasmius rotula*  
Authors : Piontek, K.; Strittmatter, E.; Plattner, D.A.  
Deposited on : 2016-01-27  
Resolution : 1.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	recalc29047
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc29047


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2964 (1.86-1.82)
Clashscore	112137	3197 (1.86-1.82)
Ramachandran outliers	110173	3164 (1.86-1.82)
Sidechain outliers	110143	3165 (1.86-1.82)
RSRZ outliers	101464	2973 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	<div> <div></div> <div>92%</div> <div>7%</div> </div>
1	B	236	<div> <div></div> <div>93%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	351	-	-	-	X
3	ACT	B	1001	-	-	-	X
3	ACT	B	351	-	-	-	X
5	NAG	A	722	-	-	-	X
7	GOL	A	1002	-	-	-	X

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 4376 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MROUPO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	236	Total	C	N	O	S	0	2	0
			1820	1140	323	351	6			
1	B	236	Total	C	N	O	S	0	3	0
			1825	1143	324	352	6			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 6 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	3	Total	C	N	O	0	0
			39	22	2	15		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	263	Total	O	0	0
			263	263		
9	B	256	Total	O	0	0
			256	256		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### ● Molecule 1: MROUPO



#### ● Molecule 1: MROUPO



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.50Å 75.70Å 104.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.77 – 1.83 44.75 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.77-1.83) 99.8 (44.75-1.83)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.155 , 0.190 0.155 , 0.190	Depositor DCC
$R_{free}$ test set	2326 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.9	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4376	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, BMA, NAG, ACT, HEM, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/1859	0.62	0/2528
1	B	0.48	0/1867	0.61	0/2539
All	All	0.50	0/3726	0.62	0/5067

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1820	0	1801	15	4
1	B	1825	0	1809	12	1
2	A	43	0	30	1	0
2	B	43	0	30	0	0
3	A	4	0	3	1	0
3	B	8	0	6	0	1
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	28	0	25	1	0
6	A	39	0	34	2	0
7	A	6	0	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	39	0	34	2	0
9	A	263	0	0	1	3
9	B	256	0	0	2	1
All	All	4376	0	3780	28	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35[A]:ASN:ND2	9:B:2058:HOH:O	2.05	0.71
8:B:732:NAG:O3	8:B:733:MAN:O2	2.08	0.67
1:A:130:GLN:HE22	1:A:155:SER:HA	1.63	0.64
1:B:235:GLU:HA	1:B:235:GLU:OE1	1.98	0.64
1:A:192:LYS:HE3	9:A:2086:HOH:O	2.03	0.56

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34[B]:ARG:NH1	9:A:2160:HOH:O[4_445]	0.93	1.27
1:A:212:GLU:OE1	1:B:212:GLU:OE1[3_554]	1.85	0.35
1:A:34[B]:ARG:CZ	9:A:2160:HOH:O[4_445]	2.12	0.08
9:A:2069:HOH:O	9:B:2153:HOH:O[2_455]	2.14	0.06
1:A:43:LYS:NZ	3:B:1001:ACT:O[2_455]	2.19	0.01

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/236 (100%)	231 (98%)	5 (2%)	0	100	100
1	B	237/236 (100%)	232 (98%)	5 (2%)	0	100	100
All	All	473/472 (100%)	463 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/198 (101%)	196 (98%)	4 (2%)	60	45
1	B	201/198 (102%)	198 (98%)	3 (2%)	70	57
All	All	401/396 (101%)	394 (98%)	7 (2%)	64	52

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	161	PHE
1	B	160	PHE
1	B	1	SER
1	A	58	LEU
1	B	143	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	GLN
1	B	11	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	BMA	A	1001	6	11,11,12	1.15	1 (9%)	13,15,17	0.78	0
5	NAG	A	721	1,5	14,14,15	0.78	0	15,19,21	1.03	0
5	NAG	A	722	5	14,14,15	0.57	0	15,19,21	1.32	3 (20%)
6	NAG	A	731	1,6	14,14,15	0.68	0	15,19,21	0.91	1 (6%)
6	NAG	A	732	6	14,14,15	0.46	0	15,19,21	1.12	2 (13%)
8	NAG	B	731	1,8	14,14,15	0.57	0	15,19,21	1.00	2 (13%)
8	NAG	B	732	8	14,14,15	0.60	0	15,19,21	1.55	2 (13%)
8	MAN	B	733	8	11,11,12	0.48	0	13,15,17	1.16	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BMA	A	1001	6	-	0/2/19/22	0/1/1/1
5	NAG	A	721	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	722	5	-	0/6/23/26	0/1/1/1
6	NAG	A	731	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	732	6	-	0/6/23/26	0/1/1/1
8	NAG	B	731	1,8	-	0/6/23/26	0/1/1/1
8	NAG	B	732	8	-	0/6/23/26	0/1/1/1
8	MAN	B	733	8	-	0/2/19/22	1/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1001	BMA	O5-C1	-3.56	1.37	1.43

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	732	NAG	C1-O5-C5	-2.81	108.29	112.17
6	A	731	NAG	O5-C1-C2	-2.36	108.19	111.47
8	B	731	NAG	C2-N2-C7	-2.11	119.86	122.94
5	A	722	NAG	O6-C6-C5	-2.08	104.35	111.34
5	A	722	NAG	O7-C7-C8	-2.07	118.30	122.06

There are no chirality outliers.

There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	733	MAN	C1-C2-C3-C4-C5-O5

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1001	BMA	2	0
5	A	721	NAG	1	0
6	A	732	NAG	2	0
8	B	732	NAG	2	0
8	B	733	MAN	2	0

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	GOL	A	1002	-	5,5,5	0.28	0	5,5,5	0.22	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	350	1,3,4	28,50,50	0.94	3 (10%)	17,82,82	1.58	2 (11%)
3	ACT	A	351	2	1,3,3	2.76	1 (100%)	0,3,3	0.00	-
3	ACT	B	1001	-	1,3,3	1.59	0	0,3,3	0.00	-
2	HEM	B	350	1,3,4	28,50,50	0.81	1 (3%)	17,82,82	1.48	2 (11%)
3	ACT	B	351	2	1,3,3	2.32	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	1002	-	-	0/4/4/4	0/0/0/0
2	HEM	A	350	1,3,4	-	0/6/54/54	0/0/8/8
3	ACT	A	351	2	-	0/0/0/0	0/0/0/0
3	ACT	B	1001	-	-	0/0/0/0	0/0/0/0
2	HEM	B	350	1,3,4	-	0/6/54/54	0/0/8/8
3	ACT	B	351	2	-	0/0/0/0	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	350	HEM	C3B-C2B	-2.77	1.36	1.40
2	B	350	HEM	C1C-NC	2.08	1.39	1.36
2	A	350	HEM	C1C-NC	2.10	1.39	1.36
2	A	350	HEM	C4A-NA	2.12	1.40	1.36
3	B	351	ACT	CH3-C	2.32	1.51	1.48

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	350	HEM	CAD-CBD-CGD	-3.77	106.22	112.66
2	B	350	HEM	CAD-CBD-CGD	-3.18	107.23	112.66
2	B	350	HEM	C3C-C4C-NC	-2.08	107.01	110.94
2	A	350	HEM	CMC-C2C-C3C	2.59	129.71	124.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	350	HEM	1	0
3	A	351	ACT	1	0
3	B	1001	ACT	0	1

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	236/236 (100%)	-0.40	3 (1%) 77 76	13, 19, 35, 72	0
1	B	236/236 (100%)	-0.44	2 (0%) 86 86	13, 20, 34, 80	0
All	All	472/472 (100%)	-0.42	5 (1%) 80 80	13, 20, 35, 80	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	236	LEU	7.7
1	B	236	LEU	4.9
1	A	1	SER	3.2
1	B	1	SER	2.8
1	A	235	GLU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	A	722	14/15	0.87	0.13	2.07	32,47,53,56	0
5	NAG	A	721	14/15	0.87	0.13	1.73	26,30,41,43	0
6	NAG	A	731	14/15	0.96	0.08	-0.28	19,25,28,29	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
8	NAG	B	731	14/15	0.95	0.08	-	21,27,31,31	0
8	MAN	B	733	11/12	0.68	0.20	-	44,71,79,96	0
8	NAG	B	732	14/15	0.86	0.19	-	39,52,64,65	0
6	NAG	A	732	14/15	0.87	0.18	-	35,48,57,75	0
6	BMA	A	1001	11/12	0.54	0.23	-	54,71,82,83	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	GOL	A	1002	6/6	0.85	0.15	8.85	42,46,58,65	0
3	ACT	B	1001	4/4	0.73	0.21	6.57	43,49,51,57	0
3	ACT	B	351	4/4	0.94	0.13	4.20	17,21,27,27	0
3	ACT	A	351	4/4	0.93	0.13	3.31	19,22,24,26	0
4	MG	A	353	1/1	0.97	0.09	1.22	15,15,15,15	0
2	HEM	B	350	43/43	0.98	0.09	0.70	11,13,16,23	0
2	HEM	A	350	43/43	0.98	0.09	0.62	11,14,18,20	0
4	MG	B	353	1/1	0.99	0.07	-1.76	16,16,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.