



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:38 am GMT

PDB ID : 3FV6
Title : Crystal Structure of the CBS domains from the Bacillus subtilis CcpN repressor
Authors : Chaix, D.; Arold, S.; Hoh, F.; Declerck, N.
Deposited on : 2009-01-15
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	<div> <div>6%</div> <div> <div></div> <div>65%</div> <div>23%</div> <div>• • 9%</div> </div> </div>
1	B	159	<div> <div>3%</div> <div> <div></div> <div>73%</div> <div>16%</div> <div>• 8%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YqzB protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	145	Total	C	N	O	S	0	0	0
			1143	729	193	214	7			
1	B	146	Total	C	N	O	S	0	0	0
			1142	726	195	214	7			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	MET	-	EXPRESSION TAG	UNP O34994
A	213	LEU	-	EXPRESSION TAG	UNP O34994
A	214	GLN	-	EXPRESSION TAG	UNP O34994
A	215	HIS	-	EXPRESSION TAG	UNP O34994
A	216	HIS	-	EXPRESSION TAG	UNP O34994
A	217	HIS	-	EXPRESSION TAG	UNP O34994
A	218	HIS	-	EXPRESSION TAG	UNP O34994
A	219	HIS	-	EXPRESSION TAG	UNP O34994
A	220	HIS	-	EXPRESSION TAG	UNP O34994
B	62	MET	-	EXPRESSION TAG	UNP O34994
B	213	LEU	-	EXPRESSION TAG	UNP O34994
B	214	GLN	-	EXPRESSION TAG	UNP O34994
B	215	HIS	-	EXPRESSION TAG	UNP O34994
B	216	HIS	-	EXPRESSION TAG	UNP O34994
B	217	HIS	-	EXPRESSION TAG	UNP O34994
B	218	HIS	-	EXPRESSION TAG	UNP O34994
B	219	HIS	-	EXPRESSION TAG	UNP O34994
B	220	HIS	-	EXPRESSION TAG	UNP O34994

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	58	Total	O	0	0
			58	58		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	82	Total	O	0	0
			82	82		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: YqzB protein



• Molecule 1: YqzB protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	54.75Å 104.05Å 99.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.56 – 1.95 43.56 – 1.95	Depositor EDS
% Data completeness (in resolution range)	91.9 (43.56-1.95) 91.9 (43.56-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.26 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.4.0062	Depositor
R, R_{free}	0.200 , 0.272 0.218 , 0.281	Depositor DCC
R_{free} test set	642 reflections (3.42%)	DCC
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2425	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.15	5/1155 (0.4%)	1.12	8/1562 (0.5%)
1	B	1.17	2/1154 (0.2%)	1.06	4/1559 (0.3%)
All	All	1.16	7/2309 (0.3%)	1.09	12/3121 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	185	ASP	CG-OD2	9.36	1.46	1.25
1	B	160	GLU	CD-OE2	6.91	1.33	1.25
1	A	77	LYS	CD-CE	6.08	1.66	1.51
1	A	212	LEU	N-CA	5.70	1.57	1.46
1	A	185	ASP	CB-CG	5.67	1.63	1.51
1	A	185	ASP	CG-OD1	5.63	1.38	1.25
1	B	106	GLU	CD-OE2	-5.55	1.19	1.25

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	185	ASP	CB-CG-OD2	-15.04	104.77	118.30
1	B	105	LEU	CB-CG-CD2	-6.98	99.13	111.00
1	A	131	ARG	NE-CZ-NH2	-6.94	116.83	120.30
1	A	212	LEU	N-CA-CB	6.43	123.27	110.40
1	A	212	LEU	CB-CG-CD2	6.42	121.92	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	ARG	NE-CZ-NH1	5.83	123.22	120.30
1	B	193	ARG	NE-CZ-NH1	5.69	123.14	120.30
1	A	185	ASP	CB-CG-OD1	5.63	123.36	118.30
1	B	150	ARG	NE-CZ-NH2	-5.51	117.54	120.30
1	B	103	MET	CG-SD-CE	5.51	109.02	100.20
1	A	124	LEU	CA-CB-CG	5.33	127.57	115.30
1	A	105	LEU	CB-CG-CD2	-5.06	102.40	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	151	MET	Peptide
1	A	211	ILE	Peptide
1	B	207	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1143	0	1210	23	0
1	B	1142	0	1208	17	0
2	A	58	0	0	0	0
2	B	82	0	0	2	0
All	All	2425	0	2418	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:MET:SD	1:A:154:ILE:HD11	2.13	0.88
1:B:195:THR:H	1:B:198:ASN:HD22	1.27	0.83
1:B:86:ILE:O	1:B:193:ARG:NH1	2.15	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:ASP:OD1	2:B:221:HOH:O	2.07	0.72
1:A:81:LYS:HA	1:A:84:GLN:HE21	1.55	0.71
1:A:128:ASP:OD1	1:A:131:ARG:NH2	2.24	0.71
1:A:155:THR:OG1	1:A:173:LYS:HE3	1.95	0.67
1:A:211:ILE:HG22	1:A:212:LEU:HA	1.79	0.63
1:B:182:LYS:HE3	1:B:189:GLU:CD	2.19	0.63
1:B:103:MET:HG3	1:B:108:VAL:O	1.99	0.61
1:A:134:ILE:O	1:A:134:ILE:HG23	2.02	0.59
1:A:195:THR:H	1:A:198:ASN:HD22	1.50	0.58
1:A:155:THR:OG1	1:A:173:LYS:CE	2.53	0.56
1:A:151:MET:SD	1:A:154:ILE:CD1	2.91	0.56
1:B:195:THR:N	1:B:198:ASN:HD22	2.02	0.54
1:A:159:ARG:CZ	1:A:191:ILE:HD12	2.38	0.54
1:A:168:LYS:O	1:A:172:GLU:HG3	2.07	0.53
1:A:151:MET:HA	1:A:154:ILE:HG12	1.91	0.52
1:B:182:LYS:HE3	1:B:189:GLU:OE2	2.11	0.51
1:A:94:VAL:HG13	1:A:98:ASP:HB2	1.92	0.51
1:B:209:ASN:N	1:B:209:ASN:OD1	2.32	0.50
1:B:85:SER:HB2	1:B:193:ARG:HG2	1.93	0.49
1:B:120:LEU:HG	1:B:151:MET:HE1	1.95	0.48
1:A:166:ILE:HG22	1:A:199:MET:CE	2.44	0.47
1:B:155:THR:O	1:B:178:LEU:HD22	2.14	0.47
1:A:117:ASP:OD1	1:A:186:LYS:HE3	2.15	0.47
1:A:138:GLU:O	1:A:141:SER:HB2	2.15	0.47
1:B:117:ASP:O	1:B:118:ALA:HB3	2.16	0.45
1:A:87:PRO:HD3	1:A:192:GLY:HA2	1.98	0.45
1:B:169:HIS:CD2	2:B:291:HOH:O	2.70	0.45
1:B:208:GLU:HG2	1:B:208:GLU:O	2.16	0.44
1:B:151:MET:CE	1:B:154:ILE:HD12	2.49	0.43
1:A:155:THR:OG1	1:A:173:LYS:NZ	2.50	0.43
1:A:195:THR:N	1:A:198:ASN:HD22	2.17	0.42
1:B:208:GLU:CD	1:B:208:GLU:H	2.22	0.42
1:B:182:LYS:HE2	1:B:191:ILE:HG21	2.02	0.42
1:A:151:MET:SD	1:A:154:ILE:CG1	3.09	0.41
1:A:91:HIS:CE1	1:A:93:ASN:HB2	2.55	0.41
1:A:84:GLN:HE22	1:A:159:ARG:HE	1.69	0.40
1:A:115:ASP:HB3	1:A:121:VAL:HG11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/159 (90%)	135 (94%)	6 (4%)	2 (1%)	13	4
1	B	144/159 (91%)	140 (97%)	4 (3%)	0	100	100
All	All	287/318 (90%)	275 (96%)	10 (4%)	2 (1%)	25	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	151	MET
1	A	152	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/146 (92%)	126 (94%)	8 (6%)	22	9
1	B	133/146 (91%)	126 (95%)	7 (5%)	26	12
All	All	267/292 (91%)	252 (94%)	15 (6%)	25	11

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	LEU
1	A	77	LYS
1	A	117	ASP
1	A	136	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	137	GLN
1	A	139	LEU
1	A	140	THR
1	A	203	LEU
1	B	136	GLN
1	B	173	LYS
1	B	182	LYS
1	B	185	ASP
1	B	203	LEU
1	B	208	GLU
1	B	209	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	93	ASN
1	A	198	ASN
1	B	84	GLN
1	B	169	HIS
1	B	198	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	145/159 (91%)	0.77	10 (6%)	18 27	19, 27, 42, 51	0
1	B	146/159 (91%)	0.48	5 (3%)	46 56	21, 27, 36, 51	0
All	All	291/318 (91%)	0.63	15 (5%)	28 38	19, 27, 41, 51	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	136	GLN	4.2
1	B	185	ASP	3.5
1	A	151	MET	3.5
1	B	186	LYS	3.3
1	A	185	ASP	3.1
1	A	186	LYS	2.9
1	A	70	LEU	2.8
1	A	130	LEU	2.5
1	A	134	ILE	2.2
1	A	105	LEU	2.2
1	B	184	THR	2.2
1	B	208	GLU	2.1
1	A	162	TYR	2.0
1	B	66	THR	2.0
1	A	189	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.