

# wwPDB/EMDataBank EM Map/Model Validation Summary Report

Aug 17, 2017 – 09:16 AM EDT

PDB ID : 5FXG  
EMDB ID: : EMD-3352  
Title : GLUN1B-GLUN2B NMDA RECEPTOR IN ACTIVE CONFORMATION  
Authors : Tajima, N.; Karakas, E.; Grant, T.; Simorowski, N.; Diaz-Avalos, R.; Grigorieff, N.; Furukawa, H.  
Deposited on : unknown  
Resolution : 6.80 Å(reported)

This is a wwPDB/EMDataBank EM Map/Model Validation Summary Report  
for a publicly released PDB/EMDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>  
with specific help available everywhere you see the  symbol.

---

MolProbitY : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *ELECTRON MICROSCOPY*


The reported resolution of this entry is 6.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	125131	1336
Ramachandran outliers	121729	1120

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%



## 2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 12648 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-METHYL-D-ASPARTATE RECEPTOR GLUN1.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	A	648	Total	C	N	O	0	0
			3199	1903	648	648		
1	C	648	Total	C	N	O	0	0
			3199	1903	648	648		

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	GLN	ASN	engineered mutation	UNP P35439
A	260	ASP	ASN	engineered mutation	UNP P35439
A	371	GLN	ASN	engineered mutation	UNP P35439
A	492	GLN	ASN	engineered mutation	UNP P35439
A	512	GLN	ASN	engineered mutation	UNP P35439
A	615	GLN	GLU	engineered mutation	UNP P35439
A	616	SER	GLU	engineered mutation	UNP P35439
A	618	SER	GLU	engineered mutation	UNP P35439
A	619	THR	GLU	engineered mutation	UNP P35439
A	792	GLN	ASN	engineered mutation	UNP P35439
A	831	CYS	PHE	engineered mutation	UNP P35439
A	865	ASN	ARG	engineered mutation	UNP P35439
A	866	GLY	ARG	engineered mutation	UNP P35439
A	867	ALA	LYS	engineered mutation	UNP P35439
C	61	GLN	ASN	engineered mutation	UNP P35439
C	260	ASP	ASN	engineered mutation	UNP P35439
C	371	GLN	ASN	engineered mutation	UNP P35439
C	492	GLN	ASN	engineered mutation	UNP P35439
C	512	GLN	ASN	engineered mutation	UNP P35439
C	615	GLN	GLU	engineered mutation	UNP P35439
C	616	SER	GLU	engineered mutation	UNP P35439
C	618	SER	GLU	engineered mutation	UNP P35439
C	619	THR	GLU	engineered mutation	UNP P35439
C	792	GLN	ASN	engineered mutation	UNP P35439
C	831	CYS	PHE	engineered mutation	UNP P35439
C	865	ASN	ARG	engineered mutation	UNP P35439

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	866	GLY	ARG	engineered mutation	UNP P35439
C	867	ALA	LYS	engineered mutation	UNP P35439

- Molecule 2 is a protein called N-METHYL-D-ASPARTATE RECEPTOR GLUN2B.

Mol	Chain	Residues	Atoms	AltConf	Trace
2	B	634	Total C N O 3125 1857 634 634	0	0
2	D	634	Total C N O 3125 1857 634 634	0	0


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	26	GLY	-	expression tag	UNP Q00960
B	348	ASP	ASN	engineered mutation	UNP Q00960
B	557	CYS	ASP	engineered mutation	UNP Q00960
B	588	SER	CYS	engineered mutation	UNP Q00960
B	838	SER	CYS	engineered mutation	UNP Q00960
B	849	SER	CYS	engineered mutation	UNP Q00960
D	26	GLY	-	expression tag	UNP Q00960
D	348	ASP	ASN	engineered mutation	UNP Q00960
D	557	CYS	ASP	engineered mutation	UNP Q00960
D	588	SER	CYS	engineered mutation	UNP Q00960
D	838	SER	CYS	engineered mutation	UNP Q00960
D	849	SER	CYS	engineered mutation	UNP Q00960


### 3 Residue-property plots [\(i\)](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-METHYL-D-ASPARTATE RECEPTOR GLUN1




- #### • Molecule 1: N-METHYL-D-ASPARTATE RECEPTOR GLUN1






- Molecule 2: N-METHYL-D-ASPARTATE RECEPTOR GLUN2B



- Molecule 2: N-METHYL-D-ASPARTATE RECEPTOR GLUN2B

Chain D:



## 4 Experimental information (i)

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	12000	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	Not provided	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	100	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	38168	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 2$	RMSZ	# $ Z  > 2$
1	A	0.27	0/3194	0.45	0/4439
1	C	0.28	0/3194	0.45	0/4439
2	B	0.27	0/3121	0.45	0/4336
2	D	0.27	0/3121	0.45	0/4336
All	All	0.27	0/12630	0.45	0/17550

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3199	0	1427	124	0
1	C	3199	0	1427	127	0
2	B	3125	0	1379	123	0
2	D	3125	0	1379	121	0
All	All	12648	0	5612	493	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 493 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:736:LEU:O	2:B:740:ALA:HB2	1.55	1.06
2:D:736:LEU:O	2:D:740:ALA:HB2	1.55	1.04
1:A:388:TYR:HA	1:A:393:VAL:HA	1.57	0.87
1:C:388:TYR:HA	1:C:393:VAL:HA	1.57	0.86
2:B:343:THR:HA	2:B:349:LEU:H	1.42	0.83

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	638/846 (75%)	563 (88%)	75 (12%)	0	100 100
1	C	638/846 (75%)	563 (88%)	75 (12%)	0	100 100
2	B	626/827 (76%)	554 (88%)	69 (11%)	3 (0%)	32 74
2	D	626/827 (76%)	554 (88%)	69 (11%)	3 (0%)	32 74
All	All	2528/3346 (76%)	2234 (88%)	288 (11%)	6 (0%)	54 84

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	660	VAL
2	D	660	VAL
2	B	523	VAL
2	D	523	VAL
2	B	468	ILE

#### 5.3.2 Protein sidechains [\(i\)](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.