



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2017 – 04:22 AM EDT

PDB ID : 2FZW  
Title : Structure of the binary complex of the E67L mutant of human glutathione-dependent formaldehyde dehydrogenase with NAD(H)  
Authors : Sanghani, P.C.; Robinson, H.  
Deposited on : unknown  
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2964 (1.86-1.82)
Clashscore	112137	3197 (1.86-1.82)
Ramachandran outliers	110173	3164 (1.86-1.82)
Sidechain outliers	110143	3165 (1.86-1.82)
RSRZ outliers	101464	2973 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>12%</div> </div> <div>.</div> </div>
1	B	373	<div> <div>%</div> <div> <div></div> <div>92%</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6791 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alcohol dehydrogenase class III chi chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	373	Total	C	N	O	S	0	8	0
			2841	1812	479	526	24			
1	B	373	Total	C	N	O	S	0	6	0
			2820	1795	473	530	22			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	LEU	GLU	ENGINEERED	UNP P11766
B	67	LEU	GLU	ENGINEERED	UNP P11766

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	K	0	0
			1	1		
4	A	1	Total	K	0	0
			1	1		

- Molecule 5 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
5	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	460	Total	O	0	0
			460	460		
6	B	561	Total	O	0	0
			561	561		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alcohol dehydrogenase class III chi chain



- Molecule 1: Alcohol dehydrogenase class III chi chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.27Å 79.27Å 311.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.84 49.33 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-1.84) 99.8 (49.33-1.84)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.52 (at 1.84Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.177 , 0.204 0.189 , 0.182	Depositor DCC
$R_{free}$ test set	4362 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.4	Xtriage
Anisotropy	0.636	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6791	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, PO4, ZN, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/2896	0.59	2/3916 (0.1%)
1	B	0.29	0/2871	0.60	2/3881 (0.1%)
All	All	0.28	0/5767	0.59	4/7797 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	322	LYS	N-CA-C	-5.52	96.09	111.00
1	B	310	GLY	N-CA-C	5.21	126.12	113.10
1	A	322	LYS	N-CA-C	-5.18	97.02	111.00
1	A	310	GLY	N-CA-C	5.03	125.67	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2841	0	2888	39	0
1	B	2820	0	2873	22	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	5	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	10	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	44	0	26	1	0
5	B	44	0	26	2	0
6	A	460	0	0	9	0
6	B	561	0	0	5	0
All	All	6791	0	5813	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:2377:NAD:H5N	6:B:2853:HOH:O	1.53	1.07
1:A:250:GLN:HE21	1:A:250:GLN:H	1.13	0.90
1:A:274:MET:HE3	1:B:304:PRO:HB3	1.58	0.83
1:A:304:PRO:HB3	1:B:274:MET:HE3	1.64	0.80
1:A:67[B]:LEU:HD21	1:A:173[B]:CYS:SG	2.23	0.78
1:A:274:MET:SD	1:A:290:VAL:HG13	2.30	0.70
1:A:250:GLN:NE2	1:A:250:GLN:H	1.89	0.69
1:A:43[B]:VAL:HG22	6:A:2257:HOH:O	1.93	0.67
1:A:232:LYS:NZ	1:A:238:GLU:HA	2.09	0.67
1:A:173[B]:CYS:SG	1:A:174:GLY:N	2.67	0.66
1:A:366:SER:HB2	6:A:2184:HOH:O	1.98	0.63
1:A:274:MET:CE	1:B:304:PRO:HB3	2.30	0.60
1:A:295:ALA:HB3	1:A:298:GLU:OE2	2.01	0.59
1:A:361:MET:HG3	6:A:2252:HOH:O	2.01	0.59
1:B:112:LYS:HE3	1:B:112:LYS:HA	1.86	0.58
1:B:360:LEU:HD23	1:B:365:LYS:HE2	1.85	0.56
1:A:358:PHE:O	1:A:362[A]:HIS:ND1	2.39	0.55
1:B:293:VAL:HG23	6:B:2877:HOH:O	2.07	0.55
1:A:305[B]:PHE:CZ	1:A:306:GLN:NE2	2.76	0.54
1:A:362[B]:HIS:HB3	6:A:2185:HOH:O	2.07	0.54
1:B:281:CYS:HB3	1:B:286:GLY:HA3	1.89	0.54
1:B:101:PHE:CE1	1:B:111[A]:GLN:HG3	2.42	0.54
1:B:43:VAL:HG22	6:B:2427:HOH:O	2.08	0.53
1:B:274:MET:SD	1:B:290:VAL:HG13	2.49	0.52
1:A:304:PRO:CB	1:B:274:MET:HE3	2.38	0.52
1:A:232:LYS:HZ3	1:A:238:GLU:HA	1.76	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:ALA:O	1:A:233:GLU:HG3	2.11	0.50
1:A:292:GLY:HA2	5:A:1377:NAD:H1D	1.94	0.49
1:B:195:ALA:HB2	1:B:261:VAL:HG11	1.93	0.49
1:B:93:ILE:HD12	1:B:318:PHE:CB	2.42	0.49
1:A:364:GLY:HA2	6:A:2186:HOH:O	2.11	0.49
1:A:274:MET:SD	1:A:290:VAL:CG1	2.98	0.49
1:A:30:LYS:HD2	1:A:128:ARG:NH2	2.27	0.49
1:B:91:LEU:HD13	1:B:323:SER:HB2	1.94	0.48
1:B:114:ARG:HD3	6:B:2768:HOH:O	2.14	0.48
1:A:195:ALA:HB2	1:A:261:VAL:HG11	1.95	0.47
1:A:281:CYS:HB3	1:A:286:GLY:HA3	1.95	0.47
1:A:362[A]:HIS:HB2	6:A:2185:HOH:O	2.15	0.47
1:B:350:SER:OG	1:B:353:GLU:HG3	2.15	0.47
1:B:30:LYS:HD2	1:B:128:ARG:NH2	2.29	0.46
1:B:271:VAL:HA	1:B:274:MET:HE2	1.97	0.46
1:A:274:MET:CE	1:B:304:PRO:CB	2.94	0.46
1:B:93:ILE:HD12	1:B:318:PHE:HB2	1.97	0.45
1:A:241:ASN:HB3	1:A:244:ASP:OD2	2.16	0.45
1:A:91:LEU:HD13	1:A:323:SER:HB2	1.98	0.44
1:B:360:LEU:HD23	1:B:365:LYS:CE	2.47	0.44
1:A:347[B]:HIS:HD2	1:A:349:LEU:HG	1.82	0.44
1:A:271:VAL:HA	1:A:274:MET:HG3	1.99	0.43
1:A:114:ARG:HD3	6:A:2113:HOH:O	2.18	0.43
1:A:91:LEU:HA	6:A:1834:HOH:O	2.19	0.42
1:B:274:MET:SD	1:B:290:VAL:CG1	3.08	0.42
5:B:2377:NAD:H8A	6:B:2838:HOH:O	2.19	0.42
1:A:67[B]:LEU:HD11	1:A:173[B]:CYS:SG	2.60	0.41
1:A:188:LEU:HD11	1:A:213:ALA:HB3	2.02	0.41
1:A:104:ASN:HA	1:A:105:PRO:HD3	1.91	0.41
1:A:232:LYS:HZ1	1:A:238:GLU:HA	1.83	0.41
1:A:161:PRO:HG3	6:A:2139:HOH:O	2.20	0.41
1:A:62:VAL:HG22	1:A:63:ILE:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/373 (102%)	365 (96%)	12 (3%)	2 (0%)	32	16
1	B	377/373 (101%)	367 (97%)	9 (2%)	1 (0%)	44	28
All	All	756/746 (101%)	732 (97%)	21 (3%)	3 (0%)	44	21

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	173[A]	CYS
1	A	173[B]	CYS
1	B	173	CYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/299 (103%)	305 (99%)	2 (1%)	87	82
1	B	305/299 (102%)	302 (99%)	3 (1%)	80	73
All	All	612/598 (102%)	607 (99%)	5 (1%)	85	79

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	250	GLN
1	A	313	TRP
1	B	112	LYS
1	B	188	LEU
1	B	313	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	111	GLN
1	A	184	ASN
1	A	250	GLN
1	B	184	ASN
1	B	243	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NAD	A	1377	-	41,48,48	2.26	8 (19%)	43,73,73	2.12	10 (23%)
3	PO4	A	1803	-	4,4,4	1.36	0	6,6,6	0.39	0
3	PO4	B	1801	-	4,4,4	1.38	0	6,6,6	0.39	0
3	PO4	B	1802	-	4,4,4	1.37	0	6,6,6	0.39	0
5	NAD	B	2377	-	41,48,48	2.45	8 (19%)	43,73,73	2.44	12 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAD	A	1377	-	-	0/22/62/62	0/5/5/5
3	PO4	A	1803	-	-	0/0/0/0	0/0/0/0
3	PO4	B	1801	-	-	0/0/0/0	0/0/0/0
3	PO4	B	1802	-	-	0/0/0/0	0/0/0/0
5	NAD	B	2377	-	-	0/22/62/62	0/5/5/5

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	2377	NAD	C3N-C7N	-12.67	1.30	1.50
5	A	1377	NAD	C3N-C7N	-11.22	1.33	1.50
5	B	2377	NAD	C2N-C3N	-3.16	1.34	1.39
5	B	2377	NAD	C5N-C4N	-3.16	1.32	1.38
5	A	1377	NAD	C5N-C4N	-2.53	1.34	1.38
5	A	1377	NAD	C2N-C3N	-2.41	1.35	1.39
5	A	1377	NAD	C5A-C4A	-2.02	1.36	1.40
5	B	2377	NAD	C6N-N1N	2.04	1.40	1.35
5	B	2377	NAD	C4A-N3A	2.06	1.38	1.35
5	A	1377	NAD	C6N-N1N	2.43	1.41	1.35
5	B	2377	NAD	C8A-N7A	2.76	1.39	1.34
5	A	1377	NAD	C8A-N7A	2.94	1.40	1.34
5	B	2377	NAD	C2A-N1A	3.29	1.40	1.33
5	A	1377	NAD	C2A-N1A	3.39	1.40	1.33
5	A	1377	NAD	C2A-N3A	4.41	1.39	1.32
5	B	2377	NAD	C2A-N3A	4.64	1.39	1.32

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	2377	NAD	N3A-C2A-N1A	-9.29	120.77	128.86
5	A	1377	NAD	N3A-C2A-N1A	-8.36	121.58	128.86
5	B	2377	NAD	C4D-O4D-C1D	-7.27	102.03	109.77
5	A	1377	NAD	C4B-O4B-C1B	-5.33	104.10	109.77
5	A	1377	NAD	C4D-O4D-C1D	-5.05	104.39	109.77
5	B	2377	NAD	C4B-O4B-C1B	-4.18	105.32	109.77
5	B	2377	NAD	C5N-C6N-N1N	-3.74	114.65	120.40
5	B	2377	NAD	O5D-C5D-C4D	-3.28	97.37	109.00
5	A	1377	NAD	C2B-C3B-C4B	-2.61	97.54	102.62
5	A	1377	NAD	O7N-C7N-N7N	-2.52	119.00	122.58
5	A	1377	NAD	C2D-C3D-C4D	-2.49	97.77	102.62

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	2377	NAD	O7N-C7N-N7N	-2.34	119.25	122.58
5	B	2377	NAD	C2D-C3D-C4D	-2.08	98.57	102.62
5	B	2377	NAD	C2B-C3B-C4B	-2.04	98.65	102.62
5	B	2377	NAD	O4B-C4B-C3B	2.33	109.81	105.17
5	A	1377	NAD	C6N-C5N-C4N	2.34	122.97	119.44
5	A	1377	NAD	O7N-C7N-C3N	2.41	122.44	119.62
5	B	2377	NAD	C3N-C7N-N7N	2.52	120.66	117.77
5	A	1377	NAD	O5D-C5D-C4D	2.82	119.00	109.00
5	A	1377	NAD	O4B-C4B-C3B	2.95	111.03	105.17
5	B	2377	NAD	O4D-C4D-C3D	3.51	112.14	105.17
5	B	2377	NAD	C6N-C5N-C4N	4.29	125.90	119.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1377	NAD	1	0
5	B	2377	NAD	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	373/373 (100%)	-0.04	8 (2%) 64 62	10, 17, 32, 47	0
1	B	373/373 (100%)	-0.29	4 (1%) 80 80	10, 15, 26, 43	0
All	All	746/746 (100%)	-0.16	12 (1%) 72 71	10, 16, 31, 47	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	365	LYS	4.9
1	A	362[A]	HIS	4.8
1	A	1	ALA	4.2
1	B	1	ALA	3.7
1	A	274	MET	2.8
1	B	274	MET	2.5
1	B	295	ALA	2.4
1	A	246	SER	2.3
1	B	294	ALA	2.3
1	A	58	GLY	2.2
1	A	373	ILE	2.2
1	A	296	SER	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAD	B	2377	44/44	0.96	0.11	0.21	12,15,39,41	0
5	NAD	A	1377	44/44	0.95	0.10	-0.21	16,20,32,34	0
2	ZN	B	375	1/1	1.00	0.06	-1.17	14,14,14,14	0
2	ZN	A	375	1/1	0.99	0.04	-1.27	16,16,16,16	0
2	ZN	B	376	1/1	1.00	0.06	-1.38	14,14,14,14	0
2	ZN	A	376	1/1	0.99	0.08	-1.88	23,23,23,23	0
4	K	A	1804	1/1	0.99	0.08	-1.97	13,13,13,13	0
4	K	B	1805	1/1	0.99	0.05	-3.02	14,14,14,14	0
3	PO4	A	1803	5/5	0.87	0.18	-	68,68,68,68	0
3	PO4	B	1802	5/5	0.89	0.17	-	71,72,72,72	0
3	PO4	B	1801	5/5	0.81	0.19	-	63,63,64,64	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.