



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 04:37 am GMT

PDB ID : 3G1H  
Title : Crystal structure of orotidine 5'-monophosphate decarboxylase from Methanobacterium thermoautotrophicum complexed with 5,6-dihydrouridine 5'-monophosphate  
Authors : Fedorov, A.A.; Fedorov, E.V.; Chan, K.K.; Gerlt, J.A.; Almo, S.C.  
Deposited on : 2009-01-29  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949

# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:  
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	<div><div>6%</div><div>78%19%.</div></div>
1	B	228	<div><div>8%</div><div>69%24%.5%</div></div>
1	C	228	<div><div>6%</div><div>73%22%..</div></div>
1	D	228	<div><div>9%</div><div>65%28%.6%</div></div>
1	E	228	<div><div>13%</div><div>67%29%5%</div></div>
1	F	228	<div><div>12%</div><div>68%27%5%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	228	<div><div></div><div>13%</div><div>59%</div><div>35%</div><div>5%</div></div>
1	H	228	<div><div></div><div>11%</div><div>65%</div><div>29%</div><div>5%</div></div>
1	I	228	<div><div></div><div>20%</div><div>60%</div><div>29%</div><div>7%</div></div>
1	J	228	<div><div></div><div>7%</div><div>70%</div><div>24%</div><div></div></div>
1	K	228	<div><div></div><div>24%</div><div>60%</div><div>34%</div><div>5%</div></div>
1	L	228	<div><div></div><div>20%</div><div>57%</div><div>36%</div><div>5%</div></div>
1	M	228	<div><div></div><div>18%</div><div>74%</div><div>19%</div><div>5%</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 21877 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Orotidine 5'-phosphate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	0	0
			1679	1054	295	318	12			
1	B	216	Total	C	N	O	S	0	0	0
			1638	1029	287	311	11			
1	C	218	Total	C	N	O	S	0	0	0
			1653	1039	289	313	12			
1	D	215	Total	C	N	O	S	0	0	0
			1630	1025	286	308	11			
1	E	217	Total	C	N	O	S	0	0	0
			1646	1034	288	312	12			
1	F	217	Total	C	N	O	S	0	0	0
			1646	1034	288	312	12			
1	G	217	Total	C	N	O	S	0	0	0
			1646	1034	288	312	12			
1	H	216	Total	C	N	O	S	0	0	0
			1638	1029	287	311	11			
1	I	212	Total	C	N	O	S	0	0	0
			1607	1011	282	304	10			
1	J	218	Total	C	N	O	S	0	0	0
			1653	1039	289	313	12			
1	K	216	Total	C	N	O	S	0	0	0
			1638	1029	287	311	11			
1	L	216	Total	C	N	O	S	0	0	0
			1638	1029	287	311	11			
1	M	217	Total	C	N	O	S	0	0	0
			1646	1034	288	312	12			

There are 13 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	101	PRO	ARG	ENGINEERED	UNP O26232
B	101	PRO	ARG	ENGINEERED	UNP O26232
C	101	PRO	ARG	ENGINEERED	UNP O26232

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	101	PRO	ARG	ENGINEERED	UNP O26232
E	101	PRO	ARG	ENGINEERED	UNP O26232
F	101	PRO	ARG	ENGINEERED	UNP O26232
G	101	PRO	ARG	ENGINEERED	UNP O26232
H	101	PRO	ARG	ENGINEERED	UNP O26232
I	101	PRO	ARG	ENGINEERED	UNP O26232
J	101	PRO	ARG	ENGINEERED	UNP O26232
K	101	PRO	ARG	ENGINEERED	UNP O26232
L	101	PRO	ARG	ENGINEERED	UNP O26232
M	101	PRO	ARG	ENGINEERED	UNP O26232

- Molecule 2 is 5,6-DIHYDROURIDINE-5'-MONOPHOSPHATE (three-letter code: H2U) (formula: C<sub>9</sub>H<sub>15</sub>N<sub>2</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	B	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	C	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	D	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	E	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	F	1	Total	C	N	O	P	0	0
			21	9	2	9	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	G	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	H	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	I	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	J	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	K	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	L	1	Total	C	N	O	P	0	0
			21	9	2	9	1		
2	M	1	Total	C	N	O	P	0	0
			21	9	2	9	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	45	Total	O	0	0
			45	45		
3	B	26	Total	O	0	0
			26	26		
3	C	44	Total	O	0	0
			44	44		
3	D	26	Total	O	0	0
			26	26		
3	E	9	Total	O	0	0
			9	9		
3	F	10	Total	O	0	0
			10	10		
3	G	7	Total	O	0	0
			7	7		
3	H	3	Total	O	0	0
			3	3		
3	I	14	Total	O	0	0
			14	14		
3	J	27	Total	O	0	0
			27	27		
3	K	4	Total	O	0	0
			4	4		
3	L	1	Total	O	0	0
			1	1		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	M	30	Total	O	0	0
			30	30		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Orotidine 5'-phosphate decarboxylase



#### • Molecule 1: Orotidine 5'-phosphate decarboxylase




#### • Molecule 1: Orotidine 5'-phosphate decarboxylase




#### • Molecule 1: Orotidine 5'-phosphate decarboxylase









• Molecule 1: Orotidine 5'-phosphate decarboxylase




• Molecule 1: Orotidine 5'-phosphate decarboxylase




• Molecule 1: Orotidine 5'-phosphate decarboxylase




## ● Molecule 1: Orotidine 5'-phosphate decarboxylase




## ● Molecule 1: Orotidine 5'-phosphate decarboxylase




## ● Molecule 1: Orotidine 5'-phosphate decarboxylase




## ● Molecule 1: Orotidine 5'-phosphate decarboxylase







• Molecule 1: Orotidine 5'-phosphate decarboxylase



• Molecule 1: Orotidine 5'-phosphate decarboxylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	147.90Å 101.80Å 192.83Å 90.00° 91.59° 90.00°	Depositor
Resolution (Å)	24.76 – 2.30 39.92 – 2.09	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.76-2.30) 96.7 (39.92-2.09)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.05 (at 2.10Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.243 , 0.285 0.242 , 0.283	Depositor DCC
$R_{free}$ test set	6367 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	45.5	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	21877	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: H2U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.47	0/1702	0.70	1/2296 (0.0%)
1	B	0.48	1/1661 (0.1%)	0.63	0/2241
1	C	0.45	0/1676	0.65	0/2261
1	D	0.42	0/1653	0.62	0/2230
1	E	0.34	0/1669	0.58	0/2251
1	F	0.38	0/1669	0.60	0/2251
1	G	0.33	0/1669	0.57	0/2251
1	H	0.34	0/1661	0.56	0/2241
1	I	0.34	0/1630	0.59	0/2199
1	J	0.38	0/1676	0.60	0/2261
1	K	0.30	0/1661	0.52	0/2241
1	L	0.30	0/1661	0.52	0/2241
1	M	0.53	1/1669 (0.1%)	1.27	6/2251 (0.3%)
All	All	0.40	2/21657 (0.0%)	0.67	7/29215 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	109	CYS	CB-SG	7.00	1.94	1.82
1	M	11	VAL	CB-CG2	5.01	1.63	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	M	144	ARG	NE-CZ-NH1	-29.35	105.63	120.30
1	M	66	ARG	CA-CB-CG	26.22	171.07	113.40
1	M	144	ARG	NE-CZ-NH2	24.90	132.75	120.30
1	M	118	ARG	CG-CD-NE	18.51	150.67	111.80
1	M	144	ARG	CD-NE-CZ	13.57	142.60	123.60

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1679	0	1695	36	0
1	B	1638	0	1651	54	0
1	C	1653	0	1669	48	0
1	D	1630	0	1647	60	0
1	E	1646	0	1660	53	0
1	F	1646	0	1660	55	0
1	G	1646	0	1660	75	0
1	H	1638	0	1651	53	0
1	I	1607	0	1623	75	0
1	J	1653	0	1669	55	0
1	K	1638	0	1651	67	1
1	L	1638	0	1651	84	0
1	M	1646	0	1660	44	0
2	A	21	0	13	0	0
2	B	21	0	13	0	0
2	C	21	0	13	1	0
2	D	21	0	13	2	0
2	E	21	0	13	0	0
2	F	21	0	13	1	0
2	G	21	0	13	1	0
2	H	21	0	13	0	0
2	I	21	0	13	2	0
2	J	21	0	13	0	0
2	K	21	0	13	0	0
2	L	21	0	13	0	0
2	M	21	0	13	2	0
3	A	45	0	0	0	0
3	B	26	0	0	1	0
3	C	44	0	0	2	0
3	D	26	0	0	1	0
3	E	9	0	0	2	0
3	F	10	0	0	0	0
3	G	7	0	0	0	0
3	H	3	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	I	14	0	0	3	0
3	J	27	0	0	1	0
3	K	4	0	0	0	0
3	L	1	0	0	0	0
3	M	30	0	0	5	0
All	All	21877	0	21716	713	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 713 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:M:116:MET:HE2	3:M:248:HOH:O	1.40	1.18
1:J:58:GLU:HG2	1:J:62:ARG:HD2	1.43	0.99
1:B:23:ASN:HD22	1:B:26:ASP:H	1.05	0.98
1:A:5:ARG:HG3	1:A:6:VAL:H	1.30	0.96
1:K:189:PRO:HB3	1:K:222:ILE:HD11	1.52	0.91

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:118:ARG:NH1	1:K:118:ARG:NH1[2_354]	2.08	0.12

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/228 (96%)	214 (98%)	5 (2%)	0	100	100
1	B	214/228 (94%)	202 (94%)	12 (6%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	216/228 (95%)	206 (95%)	10 (5%)	0	100	100
1	D	213/228 (93%)	201 (94%)	10 (5%)	2 (1%)	20	23
1	E	215/228 (94%)	198 (92%)	17 (8%)	0	100	100
1	F	215/228 (94%)	195 (91%)	16 (7%)	4 (2%)	9	8
1	G	215/228 (94%)	198 (92%)	17 (8%)	0	100	100
1	H	214/228 (94%)	200 (94%)	12 (6%)	2 (1%)	20	23
1	I	210/228 (92%)	183 (87%)	22 (10%)	5 (2%)	7	5
1	J	216/228 (95%)	196 (91%)	18 (8%)	2 (1%)	20	23
1	K	214/228 (94%)	191 (89%)	19 (9%)	4 (2%)	9	8
1	L	214/228 (94%)	194 (91%)	18 (8%)	2 (1%)	20	23
1	M	215/228 (94%)	205 (95%)	10 (5%)	0	100	100
All	All	2790/2964 (94%)	2583 (93%)	186 (7%)	21 (1%)	22	26

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	184	ALA
1	F	14	ARG
1	I	130	GLY
1	K	203	ARG
1	F	39	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/182 (96%)	173 (99%)	2 (1%)	78	89
1	B	170/182 (93%)	164 (96%)	6 (4%)	41	56
1	C	172/182 (94%)	169 (98%)	3 (2%)	66	81
1	D	169/182 (93%)	168 (99%)	1 (1%)	89	95
1	E	171/182 (94%)	169 (99%)	2 (1%)	75	87

Continued on next page...



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	171/182 (94%)	169 (99%)	2 (1%)	75	87
1	G	171/182 (94%)	167 (98%)	4 (2%)	56	73
1	H	170/182 (93%)	169 (99%)	1 (1%)	89	95
1	I	166/182 (91%)	159 (96%)	7 (4%)	34	47
1	J	172/182 (94%)	169 (98%)	3 (2%)	66	81
1	K	170/182 (93%)	167 (98%)	3 (2%)	64	79
1	L	170/182 (93%)	165 (97%)	5 (3%)	48	64
1	M	171/182 (94%)	169 (99%)	2 (1%)	75	87
All	All	2218/2366 (94%)	2177 (98%)	41 (2%)	64	79

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	167	LEU
1	I	63	PHE
1	L	195	PHE
1	G	206	TYR
1	H	210	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	13	ASN
1	G	136	GLN
1	M	98	HIS
1	G	23	ASN
1	H	210	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	H2U	A	229	-	21,22,22	1.77	5 (23%)	28,33,33	1.89	5 (17%)
2	H2U	B	229	-	21,22,22	1.61	4 (19%)	28,33,33	1.91	5 (17%)
2	H2U	C	229	-	21,22,22	1.62	6 (28%)	28,33,33	1.93	5 (17%)
2	H2U	D	229	-	21,22,22	1.67	5 (23%)	28,33,33	1.93	5 (17%)
2	H2U	E	229	-	21,22,22	1.61	5 (23%)	28,33,33	1.92	5 (17%)
2	H2U	F	229	-	21,22,22	1.68	6 (28%)	28,33,33	1.93	5 (17%)
2	H2U	G	229	-	21,22,22	1.60	4 (19%)	28,33,33	1.91	5 (17%)
2	H2U	H	229	-	21,22,22	1.66	5 (23%)	28,33,33	1.94	5 (17%)
2	H2U	I	229	-	21,22,22	1.69	5 (23%)	28,33,33	1.95	5 (17%)
2	H2U	J	229	-	21,22,22	1.72	5 (23%)	28,33,33	1.94	5 (17%)
2	H2U	K	229	-	21,22,22	1.70	5 (23%)	28,33,33	1.94	5 (17%)
2	H2U	L	229	-	21,22,22	1.67	5 (23%)	28,33,33	1.95	5 (17%)
2	H2U	M	229	-	21,22,22	1.70	5 (23%)	28,33,33	1.94	5 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	H2U	A	229	-	-	0/10/39/39	0/2/2/2
2	H2U	B	229	-	-	0/10/39/39	0/2/2/2
2	H2U	C	229	-	-	0/10/39/39	0/2/2/2
2	H2U	D	229	-	-	0/10/39/39	0/2/2/2

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	H2U	E	229	-	-	0/10/39/39	0/2/2/2
2	H2U	F	229	-	-	0/10/39/39	0/2/2/2
2	H2U	G	229	-	-	0/10/39/39	0/2/2/2
2	H2U	H	229	-	-	0/10/39/39	0/2/2/2
2	H2U	I	229	-	-	0/10/39/39	0/2/2/2
2	H2U	J	229	-	-	0/10/39/39	0/2/2/2
2	H2U	K	229	-	-	0/10/39/39	0/2/2/2
2	H2U	L	229	-	-	0/10/39/39	0/2/2/2
2	H2U	M	229	-	-	0/10/39/39	0/2/2/2

The worst 5 of 65 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	229	H2U	C2'-C3'	-2.25	1.47	1.53
2	D	229	H2U	C6-N1	2.00	1.49	1.47
2	F	229	H2U	C6-C5	2.01	1.56	1.52
2	H	229	H2U	C6-N1	2.10	1.49	1.47
2	E	229	H2U	C6-N1	2.11	1.49	1.47

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	229	H2U	C4-N3-C2	-5.79	120.85	125.81
2	C	229	H2U	C4-N3-C2	-5.78	120.86	125.81
2	L	229	H2U	C4-N3-C2	-5.76	120.88	125.81
2	J	229	H2U	C4-N3-C2	-5.76	120.88	125.81
2	K	229	H2U	C4-N3-C2	-5.74	120.89	125.81

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	229	H2U	1	0
2	D	229	H2U	2	0
2	F	229	H2U	1	0
2	G	229	H2U	1	0
2	I	229	H2U	2	0
2	M	229	H2U	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	221/228 (96%)	0.15	14 (6%) 21 27	26, 39, 59, 91	0
1	B	216/228 (94%)	0.33	19 (8%) 11 15	28, 47, 71, 102	0
1	C	218/228 (95%)	0.22	14 (6%) 20 26	29, 42, 71, 90	0
1	D	215/228 (94%)	0.62	21 (9%) 8 11	31, 52, 95, 124	0
1	E	217/228 (95%)	0.67	29 (13%) 4 5	46, 66, 85, 107	0
1	F	217/228 (95%)	0.58	28 (12%) 4 6	37, 62, 95, 114	0
1	G	217/228 (95%)	0.82	29 (13%) 4 5	50, 66, 105, 134	0
1	H	216/228 (94%)	0.73	26 (12%) 5 7	46, 67, 99, 124	0
1	I	212/228 (92%)	1.17	45 (21%) 1 1	36, 66, 139, 152	0
1	J	218/228 (95%)	0.45	17 (7%) 14 19	32, 58, 89, 112	0
1	K	216/228 (94%)	1.19	54 (25%) 1 1	62, 83, 103, 114	0
1	L	216/228 (94%)	1.17	46 (21%) 1 1	56, 90, 113, 130	0
1	M	217/228 (95%)	1.04	40 (18%) 1 2	33, 56, 86, 117	0
All	All	2816/2964 (95%)	0.70	382 (13%) 3 5	26, 61, 103, 152	0

The worst 5 of 382 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	225	LEU	9.6
1	I	159	THR	9.5
1	D	225	LEU	9.1
1	A	225	LEU	8.6
1	M	224	ASP	7.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	H2U	L	229	21/21	0.73	0.25	0.63	118,119,123,124	0
2	H2U	C	229	21/21	0.94	0.21	0.48	37,39,48,51	0
2	H2U	J	229	21/21	0.89	0.20	0.47	64,67,69,70	0
2	H2U	A	229	21/21	0.96	0.18	0.42	31,37,40,43	0
2	H2U	I	229	21/21	0.63	0.33	0.39	137,138,138,139	0
2	H2U	F	229	21/21	0.80	0.17	-0.01	57,64,69,70	0
2	H2U	B	229	21/21	0.94	0.15	-0.05	42,45,52,54	0
2	H2U	G	229	21/21	0.91	0.17	-0.15	64,66,75,76	0
2	H2U	E	229	21/21	0.94	0.16	-0.23	57,59,64,64	0
2	H2U	M	229	21/21	0.89	0.18	-0.37	59,66,68,70	0
2	H2U	K	229	21/21	0.89	0.16	-0.41	72,75,77,78	0
2	H2U	H	229	21/21	0.93	0.13	-0.89	64,68,78,79	0
2	H2U	D	229	21/21	0.92	0.14	-0.90	54,58,65,66	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.