



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2017 – 12:26 AM EDT

PDB ID : 3G8T  
Title : Crystal structure of the G33A mutant Bacillus anthracis glmS ribozyme bound to GlcN6P  
Authors : Strobel, S.A.; Cochrane, J.C.; Lipchock, S.V.; Smith, K.D.  
Deposited on : unknown  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1692 (3.00-3.00)
Clashscore	112137	2037 (3.00-3.00)
Ramachandran outliers	110173	1973 (3.00-3.00)
Sidechain outliers	110143	1976 (3.00-3.00)
RSRZ outliers	101464	1716 (3.00-3.00)
RNA backbone	2435	1007 (3.34-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	98	<div> <div>0.1%</div> <div>79%</div> <div>13%</div> <div>8%</div> </div>
1	B	98	<div> <div>2%</div> <div>81%</div> <div>11%</div> <div>8%</div> </div>
1	C	98	<div> <div>88%</div> <div>•</div> <div>8%</div> </div>
1	D	98	<div> <div>0.1%</div> <div>86%</div> <div>•</div> <div>10%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	E	13	
2	F	13	
2	G	13	
2	H	13	
3	P	141	
3	Q	141	
3	R	141	
3	S	141	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	A2M	F	0	-	-	X	-
2	A2M	G	0	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 16182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called U1 small nuclear ribonucleoprotein A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	90	Total	C	N	O	S	0	0	0
			716	459	125	129	3			
1	B	90	Total	C	N	O	S	0	0	0
			716	459	125	129	3			
1	C	90	Total	C	N	O	S	0	0	0
			716	459	125	129	3			
1	D	88	Total	C	N	O	S	0	0	0
			706	453	123	127	3			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	HIS	TYR	ENGINEERED	UNP P09012
A	36	ARG	GLN	ENGINEERED	UNP P09012
B	31	HIS	TYR	ENGINEERED	UNP P09012
B	36	ARG	GLN	ENGINEERED	UNP P09012
C	31	HIS	TYR	ENGINEERED	UNP P09012
C	36	ARG	GLN	ENGINEERED	UNP P09012
D	31	HIS	TYR	ENGINEERED	UNP P09012
D	36	ARG	GLN	ENGINEERED	UNP P09012


- Molecule 2 is a RNA chain called RNA (5'-R(\*AP\*(A2M)P\*GP\*CP\*GP\*CP\*CP\*AP\*GP\*AP\*AP\*CP\*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	13	Total	C	N	O	P	0	0	0
			277	126	54	85	12			
2	F	13	Total	C	N	O	P	0	0	0
			277	126	54	85	12			
2	G	13	Total	C	N	O	P	0	0	0
			277	126	54	85	12			
2	H	13	Total	C	N	O	P	0	0	0
			277	126	54	85	12			

- Molecule 3 is a RNA chain called glmS glucosamine-6-phosphate activated ribozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	141	Total	C	N	O	P	0	0	0
			3010	1340	532	995	143			
3	Q	141	Total	C	N	O	P	0	0	0
			3018	1344	535	996	143			
3	R	141	Total	C	N	O	P	0	0	0
			3018	1344	535	996	143			
3	S	141	Total	C	N	O	P	0	0	0
			3018	1344	535	996	143			

- Molecule 4 is GLUCOSAMINE 6-PHOSPHATE (three-letter code: GLP) (formula: C<sub>6</sub>H<sub>14</sub>NO<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	E	1	Total	C	N	O	P	0	0
			16	6	1	8	1		
4	Q	1	Total	C	N	O	P	0	0
			16	6	1	8	1		
4	R	1	Total	C	N	O	P	0	0
			16	6	1	8	1		
4	H	1	Total	C	N	O	P	0	0
			16	6	1	8	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	P	2	Total	Mg	0	0
			2	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	R	3	Total 3	Mg 3	0	0
5	Q	3	Total 3	Mg 3	0	0
5	F	2	Total 2	Mg 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total 2	O 2	0	0
6	E	2	Total 2	O 2	0	0
6	P	19	Total 19	O 19	0	0
6	B	5	Total 5	O 5	0	0
6	F	1	Total 1	O 1	0	0
6	Q	12	Total 12	O 12	0	0
6	C	3	Total 3	O 3	0	0
6	G	1	Total 1	O 1	0	0
6	R	14	Total 14	O 14	0	0
6	D	2	Total 2	O 2	0	0
6	H	3	Total 3	O 3	0	0
6	S	18	Total 18	O 18	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: U1 small nuclear ribonucleoprotein A




- Molecule 1: U1 small nuclear ribonucleoprotein A




- Molecule 1: U1 small nuclear ribonucleoprotein A



- Molecule 1: U1 small nuclear ribonucleoprotein A




- Molecule 2: RNA (5'-R(\*AP\*(A2M)P\*GP\*CP\*GP\*CP\*CP\*AP\*GP\*AP\*AP\*CP\*U)-3')




- Molecule 2: RNA (5'-R(\*AP\*(A2M)P\*GP\*CP\*GP\*CP\*CP\*AP\*GP\*AP\*AP\*CP\*U)-3')







- Molecule 2: RNA (5'-R(\*AP\*(A2M)P\*GP\*CP\*GP\*CP\*CP\*AP\*GP\*AP\*AP\*CP\*U)-3')




- Molecule 2: RNA (5'-R(\*AP\*(A2M)P\*GP\*CP\*GP\*CP\*CP\*AP\*GP\*AP\*AP\*CP\*U)-3')




- Molecule 3: glmS glucosamine-6-phosphate activated ribozyme



- Molecule 3: glmS glucosamine-6-phosphate activated ribozyme




- Molecule 3: glmS glucosamine-6-phosphate activated ribozyme



- Molecule 3: glmS glucosamine-6-phosphate activated ribozyme





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.39 Å   232.67 Å   106.58 Å 90.00°   92.21°   90.00°	Depositor
Resolution (Å)	28.23 – 3.00 28.22 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.7 (28.23-3.00) 97.8 (28.22-3.00)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.88 (at 3.00 Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.250 , 0.318 0.264 , 0.312	Depositor DCC
$R_{free}$ test set	2333 reflections (5.36%)	DCC
Wilson B-factor (Å <sup>2</sup> )	64.2	Xtriage
Anisotropy	0.538	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.24 , 48.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.046 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	16182	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GLP, GTP, MG, A2M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/729	0.47	0/979
1	B	0.30	0/729	0.46	0/979
1	C	0.30	0/729	0.46	0/979
1	D	0.31	0/719	0.45	0/965
2	E	1.02	5/284 (1.8%)	1.12	0/441
2	F	0.52	0/284	0.98	0/441
2	G	0.55	0/284	1.01	1/441 (0.2%)
2	H	0.54	0/284	1.01	1/441 (0.2%)
3	P	0.59	0/3330	1.07	3/5188 (0.1%)
3	Q	0.56	0/3339	1.04	0/5202
3	R	0.59	1/3339 (0.0%)	1.07	2/5202 (0.0%)
3	S	0.59	0/3339	1.03	2/5202 (0.0%)
All	All	0.56	6/17389 (0.0%)	0.99	9/26460 (0.0%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	-1	A	C5-C4	-5.94	1.34	1.38
2	E	-1	A	N3-C4	-5.36	1.31	1.34
3	R	17(J)	C	C1'-N1	5.28	1.56	1.48
2	E	-1	A	N7-C5	-5.27	1.36	1.39
2	E	-1	A	N9-C4	-5.24	1.34	1.37

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	S	97	U	C2-N1-C1'	5.81	124.67	117.70
3	R	86	A	P-O3'-C3'	5.78	126.64	119.70
2	H	11	U	C3'-C2'-C1'	5.64	106.01	101.50
2	G	5	C	O4'-C1'-N1	5.47	112.57	108.20
3	P	86	A	C1'-O4'-C4'	-5.34	105.62	109.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	716	0	721	6	0
1	B	716	0	721	7	0
1	C	716	0	721	1	0
1	D	706	0	714	1	0
2	E	277	0	146	26	0
2	F	277	0	146	13	0
2	G	277	0	146	15	0
2	H	277	0	146	10	0
3	P	3010	0	1509	71	0
3	Q	3018	0	1514	103	0
3	R	3018	0	1514	49	0
3	S	3018	0	1514	97	0
4	E	16	0	12	2	0
4	H	16	0	12	0	0
4	Q	16	0	12	0	0
4	R	16	0	12	0	0
5	F	2	0	0	0	0
5	P	2	0	0	0	0
5	Q	3	0	0	0	0
5	R	3	0	0	0	0
6	A	2	0	0	2	0
6	B	5	0	0	3	0
6	C	3	0	0	0	0
6	D	2	0	0	0	0
6	E	2	0	0	0	0
6	F	1	0	0	0	0
6	G	1	0	0	0	0
6	H	3	0	0	0	0
6	P	19	0	0	0	0
6	Q	12	0	0	0	0
6	R	14	0	0	0	0
6	S	18	0	0	0	0
All	All	16182	0	9560	365	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 365 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:Q:85:C:H1'	3:Q:86:A:OP2	1.47	1.14
3:Q:117:G:H2'	3:Q:118:U:H6	1.13	1.13
3:Q:17(D):U:H1'	3:Q:17(E):G:C5	1.85	1.12
3:S:87:A:H4'	3:S:88:G:OP1	1.44	1.10
3:S:83:C:H4'	3:S:86:A:O2'	1.53	1.08

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	88/98 (90%)	85 (97%)	3 (3%)	0	100	100
1	B	88/98 (90%)	87 (99%)	1 (1%)	0	100	100
1	C	88/98 (90%)	88 (100%)	0	0	100	100
1	D	86/98 (88%)	85 (99%)	1 (1%)	0	100	100
All	All	350/392 (89%)	345 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/88 (88%)	74 (96%)	3 (4%)	37	75
1	B	77/88 (88%)	75 (97%)	2 (3%)	51	83
1	C	77/88 (88%)	74 (96%)	3 (4%)	37	75
1	D	77/88 (88%)	75 (97%)	2 (3%)	51	83
All	All	308/352 (88%)	298 (97%)	10 (3%)	44	79

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	47	ARG
1	C	36	ARG
1	C	71	SER
1	B	36	ARG
1	C	47	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	GLN
1	B	67	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	12/13 (92%)	7 (58%)	1 (8%)
2	F	12/13 (92%)	4 (33%)	1 (8%)
2	G	12/13 (92%)	4 (33%)	1 (8%)
2	H	12/13 (92%)	3 (25%)	1 (8%)
3	P	139/141 (98%)	35 (25%)	5 (3%)
3	Q	139/141 (98%)	40 (28%)	5 (3%)
3	R	139/141 (98%)	26 (18%)	3 (2%)
3	S	139/141 (98%)	44 (31%)	4 (2%)
All	All	604/616 (98%)	163 (26%)	21 (3%)

5 of 163 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	E	0	A2M
2	E	1	G

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	E	2	C
2	E	3	G
2	E	6	A

5 of 21 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	Q	49	U
3	Q	85	C
3	S	17(F)	C
3	Q	42	A
3	S	86	A

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	A2M	E	0	3,2	18,25,26	0.64	0	20,36,39	0.60	0
2	A2M	F	0	3,2	18,25,26	0.63	0	20,36,39	0.59	0
2	A2M	G	0	2	18,25,26	0.63	0	20,36,39	0.60	0
2	A2M	H	0	3,2	18,25,26	0.64	0	20,36,39	0.59	0
3	GTP	P	12	3	27,34,34	0.75	0	27,54,54	1.76	4 (14%)
3	GTP	Q	12	3	27,34,34	0.75	0	27,54,54	1.76	4 (14%)
3	GTP	R	12	3	27,34,34	0.76	1 (3%)	27,54,54	1.78	4 (14%)
3	GTP	S	12	3	27,34,34	0.75	0	27,54,54	1.82	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A2M	E	0	3,2	-	0/5/27/28	0/3/3/3
2	A2M	F	0	3,2	-	0/5/27/28	0/3/3/3
2	A2M	G	0	2	-	0/5/27/28	0/3/3/3
2	A2M	H	0	3,2	-	0/5/27/28	0/3/3/3
3	GTP	P	12	3	-	0/18/38/38	0/3/3/3
3	GTP	Q	12	3	-	0/18/38/38	0/3/3/3
3	GTP	R	12	3	-	0/18/38/38	0/3/3/3
3	GTP	S	12	3	-	0/18/38/38	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	R	12	GTP	O4'-C1'	2.09	1.44	1.41

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	S	12	GTP	N3-C2-N1	-4.70	120.60	127.46
3	P	12	GTP	N3-C2-N1	-4.61	120.73	127.46
3	Q	12	GTP	N3-C2-N1	-4.58	120.77	127.46
3	R	12	GTP	N3-C2-N1	-4.53	120.84	127.46
3	Q	12	GTP	C5-C6-N1	-2.91	119.34	123.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	0	A2M	6	0
2	F	0	A2M	9	0
2	G	0	A2M	10	0
2	H	0	A2M	7	0
3	P	12	GTP	1	0
3	Q	12	GTP	1	0
3	R	12	GTP	2	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.



## 5.6 Ligand geometry

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GLP	E	5001	-	16,16,16	0.64	0	22,24,24	0.91	0
4	GLP	H	5004	-	16,16,16	0.61	0	22,24,24	0.98	0
4	GLP	Q	5002	-	16,16,16	0.66	0	22,24,24	1.11	1 (4%)
4	GLP	R	5003	-	16,16,16	0.61	0	22,24,24	1.07	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLP	E	5001	-	-	0/6/26/26	0/1/1/1
4	GLP	H	5004	-	-	0/6/26/26	0/1/1/1
4	GLP	Q	5002	-	-	0/6/26/26	0/1/1/1
4	GLP	R	5003	-	-	0/6/26/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	Q	5002	GLP	C1-O5-C5	2.03	117.05	113.39
4	R	5003	GLP	O5-C5-C6	2.21	111.05	106.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	5001	GLP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	90/98 (91%)	0.30	1 (1%)	80	55	55, 58, 70, 72	0
1	B	90/98 (91%)	0.39	2 (2%)	62	33	50, 57, 67, 69	0
1	C	90/98 (91%)	0.23	0	100	100	51, 57, 68, 70	0
1	D	88/98 (89%)	0.40	1 (1%)	80	55	56, 60, 69, 71	0
2	E	12/13 (92%)	1.07	2 (16%)	2	1	53, 76, 86, 87	0
2	F	12/13 (92%)	0.65	1 (8%)	12	4	32, 46, 59, 69	0
2	G	12/13 (92%)	0.94	2 (16%)	2	1	35, 59, 65, 71	0
2	H	12/13 (92%)	0.96	2 (16%)	2	1	59, 77, 87, 87	0
3	P	140/141 (99%)	0.80	12 (8%)	11	4	57, 78, 106, 116	0
3	Q	140/141 (99%)	0.51	11 (7%)	13	5	31, 57, 105, 123	0
3	R	140/141 (99%)	0.50	8 (5%)	24	9	36, 59, 98, 107	0
3	S	140/141 (99%)	0.81	10 (7%)	17	6	58, 80, 106, 115	0
All	All	966/1008 (95%)	0.55	52 (5%)	26	11	31, 61, 98, 123	0

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	P	72	U	4.6
3	P	73	U	4.1
3	Q	139	U	3.8
3	R	72	U	3.7
2	G	6	A	3.5

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GTP	S	12	32/32	0.78	0.33	1.65	94,95,102,102	0
3	GTP	P	12	32/32	0.76	0.30	0.07	95,96,103,103	0
3	GTP	Q	12	32/32	0.86	0.24	-0.26	43,51,65,65	0
3	GTP	R	12	32/32	0.76	0.26	-0.29	70,72,83,83	0
2	A2M	H	0	23/24	0.77	0.32	-	60,61,63,64	0
2	A2M	G	0	23/24	0.86	0.28	-	43,44,55,55	0
2	A2M	E	0	23/24	0.90	0.26	-	55,56,58,58	0
2	A2M	F	0	23/24	0.87	0.23	-	42,43,54,54	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GLP	R	5003	16/16	0.91	0.28	1.50	39,41,44,44	0
4	GLP	H	5004	16/16	0.90	0.32	1.46	62,62,63,64	0
4	GLP	E	5001	16/16	0.92	0.29	0.79	55,55,56,56	0
4	GLP	Q	5002	16/16	0.94	0.22	-0.32	39,40,41,42	0
5	MG	Q	10	1/1	0.95	0.19	-	35,35,35,35	0
5	MG	F	12	1/1	0.92	0.79	-	51,51,51,51	0
5	MG	F	13	1/1	0.85	0.26	-	35,35,35,35	0
5	MG	Q	1	1/1	0.94	0.43	-	40,40,40,40	0
5	MG	R	8	1/1	0.91	0.34	-	38,38,38,38	0
5	MG	R	3	1/1	0.94	0.44	-	26,26,26,26	0
5	MG	P	6	1/1	0.95	0.28	-	38,38,38,38	0
5	MG	Q	4	1/1	0.95	0.49	-	40,40,40,40	0
5	MG	R	5	1/1	0.96	0.44	-	37,37,37,37	0
5	MG	P	9	1/1	0.88	0.37	-	35,35,35,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.