



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:27 am GMT

PDB ID : 2G9T  
Title : Crystal structure of the SARS coronavirus nsp10 at 2.1Å  
Authors : Su, D.; Lou, Z.; Yang, H.; Sun, F.; Rao, Z.  
Deposited on : 2006-03-07  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	<div> <div>3%</div> <div>72%</div> <div>5%</div> <div>22%</div> </div>
1	B	152	<div> <div>3%</div> <div>65%</div> <div>10%</div> <div>22%</div> </div>
1	C	152	<div> <div>3%</div> <div>60%</div> <div>16%</div> <div>22%</div> </div>
1	D	152	<div> <div>4%</div> <div>55%</div> <div>19%</div> <div>22%</div> </div>
1	E	152	<div> <div>3%</div> <div>56%</div> <div>18%</div> <div>22%</div> </div>
1	F	152	<div> <div>3%</div> <div>61%</div> <div>15%</div> <div>23%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	152	
1	H	152	
1	I	152	
1	J	152	
1	K	152	
1	L	152	
1	M	152	
1	N	152	
1	O	152	
1	P	152	
1	Q	152	
1	R	152	
1	S	152	
1	T	152	
1	U	152	
1	V	152	
1	W	152	
1	X	152	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 22470 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called orfla polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	B	119	Total	C	N	O	S	0	0	0
			880	549	148	168	15			
1	C	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	D	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	E	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	F	117	Total	C	N	O	S	0	0	0
			863	538	144	166	15			
1	G	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	H	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	I	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	J	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	K	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	L	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	M	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	N	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	O	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	P	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	R	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	S	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	T	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	U	116	Total	C	N	O	S	0	0	0
			856	534	143	164	15			
1	V	116	Total	C	N	O	S	0	0	0
			856	534	143	164	15			
1	W	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			
1	X	118	Total	C	N	O	S	0	0	0
			873	544	147	167	15			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	P	2	Total	Zn	0	0
			2	2		
2	K	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	W	2	Total	Zn	0	0
			2	2		
2	N	2	Total	Zn	0	0
			2	2		
2	X	2	Total	Zn	0	0
			2	2		
2	S	2	Total	Zn	0	0
			2	2		
2	J	2	Total	Zn	0	0
			2	2		
2	E	2	Total	Zn	0	0
			2	2		
2	V	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	R	2	Total 2	Zn 2	0	0
2	M	2	Total 2	Zn 2	0	0
2	D	2	Total 2	Zn 2	0	0
2	I	2	Total 2	Zn 2	0	0
2	U	2	Total 2	Zn 2	0	0
2	L	2	Total 2	Zn 2	0	0
2	G	2	Total 2	Zn 2	0	0
2	Q	2	Total 2	Zn 2	0	0
2	H	2	Total 2	Zn 2	0	0
2	C	2	Total 2	Zn 2	0	0
2	T	2	Total 2	Zn 2	0	0
2	O	2	Total 2	Zn 2	0	0
2	F	2	Total 2	Zn 2	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	69	Total 69	O 69	0	0
3	B	59	Total 59	O 59	0	0
3	C	66	Total 66	O 66	0	0
3	D	63	Total 63	O 63	0	0
3	E	63	Total 63	O 63	0	0
3	F	58	Total 58	O 58	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	65	Total 65	O 65	0	0
3	H	59	Total 59	O 59	0	0
3	I	55	Total 55	O 55	0	0
3	J	73	Total 73	O 73	0	0
3	K	71	Total 71	O 71	0	0
3	L	71	Total 71	O 71	0	0
3	M	74	Total 74	O 74	0	0
3	N	51	Total 51	O 51	0	0
3	O	45	Total 45	O 45	0	0
3	P	58	Total 58	O 58	0	0
3	Q	81	Total 81	O 81	0	0
3	R	61	Total 61	O 61	0	0
3	S	67	Total 67	O 67	0	0
3	T	62	Total 62	O 62	0	0
3	U	77	Total 77	O 77	0	0
3	V	49	Total 49	O 49	0	0
3	W	45	Total 45	O 45	0	0
3	X	65	Total 65	O 65	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide



- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide







- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide




- Molecule 1: orf1a polypeptide



- Molecule 1: orf1a polypeptide



- Molecule 1: orf1a polypeptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	159.11Å 321.83Å 161.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.10 49.80 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.10) 97.4 (49.80-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 2.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.217 , 0.247 0.221 , 0.251	Depositor DCC
$R_{free}$ test set	11393 reflections (4.83%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.5	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 56.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	22470	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.65	0/892	0.82	1/1211 (0.1%)
1	B	0.63	0/900	0.82	1/1222 (0.1%)
1	C	0.61	0/892	0.84	3/1211 (0.2%)
1	D	0.63	0/892	0.86	2/1211 (0.2%)
1	E	0.68	1/892 (0.1%)	0.82	1/1211 (0.1%)
1	F	0.60	0/880	0.79	1/1193 (0.1%)
1	G	0.66	1/892 (0.1%)	0.90	3/1211 (0.2%)
1	H	0.62	0/892	0.81	1/1211 (0.1%)
1	I	0.66	0/892	0.87	3/1211 (0.2%)
1	J	0.65	0/892	0.87	3/1211 (0.2%)
1	K	0.61	0/892	0.89	2/1211 (0.2%)
1	L	0.67	1/892 (0.1%)	0.86	1/1211 (0.1%)
1	M	0.66	0/892	0.89	3/1211 (0.2%)
1	N	0.58	0/892	0.79	1/1211 (0.1%)
1	O	0.57	0/892	0.77	1/1211 (0.1%)
1	P	0.61	0/892	0.78	0/1211
1	Q	0.67	0/892	0.83	2/1211 (0.2%)
1	R	0.64	0/892	0.82	1/1211 (0.1%)
1	S	0.64	1/892 (0.1%)	0.77	0/1211
1	T	0.70	1/892 (0.1%)	0.86	2/1211 (0.2%)
1	U	0.64	0/873	0.84	2/1183 (0.2%)
1	V	0.60	0/873	0.83	2/1183 (0.2%)
1	W	0.62	0/892	0.88	2/1211 (0.2%)
1	X	0.68	0/892	0.95	5/1211 (0.4%)
All	All	0.64	5/21366 (0.0%)	0.84	43/29001 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
1	G	0	1
1	H	0	1
1	M	0	1
1	O	0	1
1	Q	0	1
All	All	0	7

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	T	120	CYS	CB-SG	-9.17	1.66	1.82
1	E	73	CYS	CB-SG	-8.22	1.68	1.82
1	S	46	CYS	CB-SG	7.25	1.94	1.82
1	L	73	CYS	CB-SG	-6.80	1.70	1.82
1	G	46	CYS	CB-SG	5.68	1.92	1.82

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	K	78	ARG	NE-CZ-NH2	-11.94	114.33	120.30
1	I	78	ARG	NE-CZ-NH2	-11.20	114.70	120.30
1	G	78	ARG	NE-CZ-NH2	-11.14	114.73	120.30
1	W	78	ARG	NE-CZ-NH2	-10.43	115.08	120.30
1	M	78	ARG	NE-CZ-NH2	-10.17	115.21	120.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	96	TYR	Sidechain
1	E	76	TYR	Sidechain
1	G	76	TYR	Sidechain
1	H	76	TYR	Sidechain
1	M	76	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	873	0	829	7	0
1	B	880	0	837	19	0
1	C	873	0	829	27	0
1	D	873	0	829	37	0
1	E	873	0	829	30	0
1	F	863	0	821	30	0
1	G	873	0	829	27	0
1	H	873	0	829	13	0
1	I	873	0	829	21	0
1	J	873	0	829	20	0
1	K	873	0	829	30	0
1	L	873	0	829	19	0
1	M	873	0	829	32	0
1	N	873	0	829	10	0
1	O	873	0	829	23	0
1	P	873	0	829	25	0
1	Q	873	0	829	26	0
1	R	873	0	829	23	0
1	S	873	0	829	30	0
1	T	873	0	829	33	0
1	U	856	0	814	26	0
1	V	856	0	814	17	0
1	W	873	0	829	14	0
1	X	873	0	829	32	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
2	G	2	0	0	0	0
2	H	2	0	0	0	0
2	I	2	0	0	0	0
2	J	2	0	0	0	0
2	K	2	0	0	0	0
2	L	2	0	0	0	0
2	M	2	0	0	0	0
2	N	2	0	0	0	0
2	O	2	0	0	0	0
2	P	2	0	0	0	0
2	Q	2	0	0	0	0
2	R	2	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	S	2	0	0	0	0
2	T	2	0	0	0	0
2	U	2	0	0	0	0
2	V	2	0	0	0	0
2	W	2	0	0	0	0
2	X	2	0	0	0	0
3	A	69	0	0	4	0
3	B	59	0	0	4	0
3	C	66	0	0	5	0
3	D	63	0	0	4	0
3	E	63	0	0	8	0
3	F	58	0	0	6	0
3	G	65	0	0	12	0
3	H	59	0	0	2	0
3	I	55	0	0	8	0
3	J	73	0	0	5	0
3	K	71	0	0	15	0
3	L	71	0	0	0	0
3	M	74	0	0	11	0
3	N	51	0	0	1	0
3	O	45	0	0	7	0
3	P	58	0	0	11	0
3	Q	81	0	0	8	0
3	R	61	0	0	9	0
3	S	67	0	0	13	0
3	T	62	0	0	10	0
3	U	77	0	0	13	0
3	V	49	0	0	6	0
3	W	45	0	0	0	0
3	X	65	0	0	10	0
All	All	22470	0	19866	552	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 552 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:THR:HG23	1:B:49:THR:H	1.14	1.08
1:P:14:LEU:HD11	1:P:78:ARG:HH12	1.22	1.00
1:S:72:SER:HB3	3:S:1026:HOH:O	1.63	0.99

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Q:43:LYS:HD3	3:Q:1031:HOH:O	1.63	0.99
1:S:40:ASN:HB2	1:S:78:ARG:HH22	1.32	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	114/152 (75%)	110 (96%)	4 (4%)	0	100	100
1	B	115/152 (76%)	109 (95%)	5 (4%)	1 (1%)	20	14
1	C	114/152 (75%)	110 (96%)	3 (3%)	1 (1%)	20	14
1	D	114/152 (75%)	109 (96%)	4 (4%)	1 (1%)	20	14
1	E	114/152 (75%)	107 (94%)	5 (4%)	2 (2%)	10	4
1	F	111/152 (73%)	104 (94%)	6 (5%)	1 (1%)	20	14
1	G	114/152 (75%)	109 (96%)	4 (4%)	1 (1%)	20	14
1	H	114/152 (75%)	107 (94%)	6 (5%)	1 (1%)	20	14
1	I	114/152 (75%)	107 (94%)	7 (6%)	0	100	100
1	J	114/152 (75%)	111 (97%)	3 (3%)	0	100	100
1	K	114/152 (75%)	107 (94%)	6 (5%)	1 (1%)	20	14
1	L	114/152 (75%)	110 (96%)	3 (3%)	1 (1%)	20	14
1	M	114/152 (75%)	110 (96%)	2 (2%)	2 (2%)	10	4
1	N	114/152 (75%)	110 (96%)	4 (4%)	0	100	100
1	O	114/152 (75%)	105 (92%)	8 (7%)	1 (1%)	20	14
1	P	114/152 (75%)	107 (94%)	6 (5%)	1 (1%)	20	14
1	Q	114/152 (75%)	109 (96%)	4 (4%)	1 (1%)	20	14
1	R	114/152 (75%)	110 (96%)	3 (3%)	1 (1%)	20	14

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	S	114/152 (75%)	108 (95%)	5 (4%)	1 (1%)	20	14
1	T	114/152 (75%)	110 (96%)	3 (3%)	1 (1%)	20	14
1	U	110/152 (72%)	105 (96%)	4 (4%)	1 (1%)	20	14
1	V	110/152 (72%)	103 (94%)	6 (6%)	1 (1%)	20	14
1	W	114/152 (75%)	109 (96%)	4 (4%)	1 (1%)	20	14
1	X	114/152 (75%)	109 (96%)	4 (4%)	1 (1%)	20	14
All	All	2726/3648 (75%)	2595 (95%)	109 (4%)	22 (1%)	22	17

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	49	THR
1	H	84	PRO
1	P	84	PRO
1	S	84	PRO
1	T	84	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/123 (79%)	96 (99%)	1 (1%)	80	85
1	B	98/123 (80%)	94 (96%)	4 (4%)	35	35
1	C	97/123 (79%)	95 (98%)	2 (2%)	59	64
1	D	97/123 (79%)	94 (97%)	3 (3%)	45	48
1	E	97/123 (79%)	91 (94%)	6 (6%)	21	18
1	F	96/123 (78%)	94 (98%)	2 (2%)	59	64
1	G	97/123 (79%)	91 (94%)	6 (6%)	21	18
1	H	97/123 (79%)	94 (97%)	3 (3%)	45	48
1	I	97/123 (79%)	92 (95%)	5 (5%)	27	24
1	J	97/123 (79%)	96 (99%)	1 (1%)	80	85

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	97/123 (79%)	93 (96%)	4 (4%)	35	35
1	L	97/123 (79%)	94 (97%)	3 (3%)	45	48
1	M	97/123 (79%)	92 (95%)	5 (5%)	27	24
1	N	97/123 (79%)	95 (98%)	2 (2%)	59	64
1	O	97/123 (79%)	97 (100%)	0	100	100
1	P	97/123 (79%)	91 (94%)	6 (6%)	21	18
1	Q	97/123 (79%)	93 (96%)	4 (4%)	35	35
1	R	97/123 (79%)	93 (96%)	4 (4%)	35	35
1	S	97/123 (79%)	91 (94%)	6 (6%)	21	18
1	T	97/123 (79%)	92 (95%)	5 (5%)	27	24
1	U	95/123 (77%)	91 (96%)	4 (4%)	34	33
1	V	95/123 (77%)	92 (97%)	3 (3%)	44	46
1	W	97/123 (79%)	94 (97%)	3 (3%)	45	48
1	X	97/123 (79%)	94 (97%)	3 (3%)	45	48
All	All	2324/2952 (79%)	2239 (96%)	85 (4%)	39	39

5 of 85 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	85	ASN
1	P	45	LEU
1	V	124	LYS
1	L	105	ASN
1	M	89	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 59 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	105	ASN
1	O	40	ASN
1	W	105	ASN
1	M	48	HIS
1	M	85	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

Of 48 ligands modelled in this entry, 48 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers ⓘ

There are no such residues in this entry.

### 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	118/152 (77%)	0.07	1 (0%) 86 88	22, 28, 42, 54	0
1	B	119/152 (78%)	0.22	5 (4%) 37 44	21, 31, 44, 59	0
1	C	118/152 (77%)	0.23	5 (4%) 37 44	23, 31, 45, 59	0
1	D	118/152 (77%)	0.26	6 (5%) 29 35	21, 31, 47, 63	0
1	E	118/152 (77%)	0.25	4 (3%) 46 53	22, 31, 50, 61	0
1	F	117/152 (76%)	0.26	5 (4%) 36 43	24, 32, 46, 57	0
1	G	118/152 (77%)	0.28	7 (5%) 23 29	20, 29, 44, 59	0
1	H	118/152 (77%)	0.28	7 (5%) 23 29	23, 31, 47, 63	0
1	I	118/152 (77%)	0.15	4 (3%) 46 53	20, 27, 43, 52	0
1	J	118/152 (77%)	0.23	6 (5%) 29 35	22, 29, 44, 57	0
1	K	118/152 (77%)	0.23	4 (3%) 46 53	21, 31, 45, 62	0
1	L	118/152 (77%)	0.14	5 (4%) 37 44	20, 28, 41, 58	0
1	M	118/152 (77%)	0.23	4 (3%) 46 53	21, 29, 43, 60	0
1	N	118/152 (77%)	0.22	6 (5%) 29 35	25, 33, 46, 53	0
1	O	118/152 (77%)	0.37	6 (5%) 29 35	28, 37, 49, 62	0
1	P	118/152 (77%)	0.22	4 (3%) 46 53	23, 31, 44, 59	0
1	Q	118/152 (77%)	0.15	4 (3%) 46 53	18, 28, 42, 60	0
1	R	118/152 (77%)	0.10	3 (2%) 58 63	20, 28, 43, 62	0
1	S	118/152 (77%)	0.27	5 (4%) 37 44	24, 32, 45, 59	0
1	T	118/152 (77%)	0.30	7 (5%) 23 29	22, 31, 47, 58	0
1	U	116/152 (76%)	0.22	7 (6%) 23 28	19, 28, 47, 65	0
1	V	116/152 (76%)	0.45	8 (6%) 18 22	28, 34, 51, 61	0
1	W	118/152 (77%)	0.25	4 (3%) 46 53	22, 30, 45, 63	0
1	X	118/152 (77%)	0.26	8 (6%) 18 23	20, 28, 45, 59	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
All	All	2828/3648 (77%)	0.23	125 (4%) 35 41	18, 31, 46, 65	0

The worst 5 of 125 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	W	89	PHE	9.2
1	X	89	PHE	8.4
1	V	89	PHE	8.1
1	D	85	ASN	7.8
1	M	89	PHE	7.7

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	C	998	1/1	1.00	0.13	0.64	27,27,27,27	0
2	ZN	X	998	1/1	1.00	0.14	0.58	24,24,24,24	0
2	ZN	I	998	1/1	1.00	0.13	0.46	25,25,25,25	0
2	ZN	D	998	1/1	0.99	0.13	0.19	27,27,27,27	0
2	ZN	L	998	1/1	1.00	0.12	0.04	27,27,27,27	0
2	ZN	U	998	1/1	1.00	0.13	0.03	26,26,26,26	0
2	ZN	M	998	1/1	0.99	0.12	-0.01	27,27,27,27	0
2	ZN	B	998	1/1	0.99	0.12	-0.03	27,27,27,27	0
2	ZN	F	998	1/1	1.00	0.11	-0.43	27,27,27,27	0
2	ZN	G	999	1/1	0.98	0.10	-0.61	38,38,38,38	0
2	ZN	E	998	1/1	1.00	0.11	-0.63	27,27,27,27	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	S	998	1/1	1.00	0.12	-0.64	27,27,27,27	0
2	ZN	K	998	1/1	1.00	0.12	-0.69	29,29,29,29	0
2	ZN	N	998	1/1	1.00	0.12	-0.72	29,29,29,29	0
2	ZN	H	998	1/1	1.00	0.12	-0.72	29,29,29,29	0
2	ZN	P	998	1/1	0.99	0.10	-0.72	28,28,28,28	0
2	ZN	J	998	1/1	1.00	0.12	-0.74	27,27,27,27	0
2	ZN	H	999	1/1	0.99	0.10	-0.74	39,39,39,39	0
2	ZN	M	999	1/1	0.99	0.11	-0.77	43,43,43,43	0
2	ZN	U	999	1/1	0.99	0.10	-0.78	33,33,33,33	0
2	ZN	O	999	1/1	0.99	0.10	-0.84	42,42,42,42	0
2	ZN	L	999	1/1	0.99	0.10	-0.86	35,35,35,35	0
2	ZN	A	998	1/1	1.00	0.11	-0.86	27,27,27,27	0
2	ZN	S	999	1/1	1.00	0.08	-0.86	32,32,32,32	0
2	ZN	Q	998	1/1	1.00	0.11	-0.88	25,25,25,25	0
2	ZN	P	999	1/1	0.99	0.10	-0.88	38,38,38,38	0
2	ZN	I	999	1/1	0.98	0.10	-0.89	35,35,35,35	0
2	ZN	R	999	1/1	0.99	0.10	-0.90	34,34,34,34	0
2	ZN	C	999	1/1	0.99	0.09	-0.91	35,35,35,35	0
2	ZN	O	998	1/1	0.99	0.11	-0.95	34,34,34,34	0
2	ZN	G	998	1/1	0.99	0.12	-0.99	27,27,27,27	0
2	ZN	B	999	1/1	0.98	0.09	-0.99	41,41,41,41	0
2	ZN	W	998	1/1	0.99	0.12	-1.04	28,28,28,28	0
2	ZN	E	999	1/1	0.98	0.08	-1.16	35,35,35,35	0
2	ZN	T	998	1/1	1.00	0.11	-1.17	27,27,27,27	0
2	ZN	J	999	1/1	0.99	0.08	-1.20	35,35,35,35	0
2	ZN	A	999	1/1	0.99	0.08	-1.22	36,36,36,36	0
2	ZN	K	999	1/1	0.98	0.08	-1.31	37,37,37,37	0
2	ZN	V	998	1/1	0.99	0.09	-1.34	29,29,29,29	0
2	ZN	F	999	1/1	0.99	0.09	-1.42	38,38,38,38	0
2	ZN	D	999	1/1	0.98	0.07	-1.50	36,36,36,36	0
2	ZN	W	999	1/1	1.00	0.07	-1.62	34,34,34,34	0
2	ZN	T	999	1/1	0.99	0.10	-1.76	43,43,43,43	0
2	ZN	R	998	1/1	1.00	0.09	-2.08	28,28,28,28	0
2	ZN	X	999	1/1	0.98	0.06	-2.56	36,36,36,36	0
2	ZN	V	999	1/1	0.95	0.09	-	73,73,73,73	0
2	ZN	N	999	1/1	0.99	0.10	-	41,41,41,41	0
2	ZN	Q	999	1/1	0.99	0.07	-	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.