



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 09:16 pm GMT

PDB ID : 3GAA
Title : The crystal structure of the protein with unknown function from *Thermoplasma acidophilum*
Authors : Zhang, R.; Borovilos, M.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2009-02-16
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	252	<div> <div>2%</div> <div> <div></div> <div>57%</div> <div>28%</div> <div>11%</div> <div>.</div> </div> </div>
1	B	252	<div> <div>3%</div> <div> <div></div> <div>60%</div> <div>25%</div> <div>10%</div> <div>.</div> </div> </div>
1	C	252	<div> <div>4%</div> <div> <div></div> <div>55%</div> <div>32%</div> <div>8%</div> <div>.</div> </div> </div>
1	D	252	<div> <div>3%</div> <div> <div></div> <div>60%</div> <div>30%</div> <div>6%</div> <div>.</div> </div> </div>
1	E	252	<div> <div>5%</div> <div> <div></div> <div>58%</div> <div>27%</div> <div>12%</div> <div>.</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9299 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein Ta1441.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	S	0	0	0
			1856	1183	310	352	11			
1	B	242	Total	C	N	O	S	0	0	0
			1848	1178	309	351	10			
1	C	243	Total	C	N	O	S	0	0	0
			1856	1183	310	352	11			
1	D	242	Total	C	N	O	S	0	0	0
			1848	1177	309	351	11			
1	E	243	Total	C	N	O	S	0	0	0
			1856	1183	310	352	11			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	10	Total	O	0	0
			10	10		
2	B	10	Total	O	0	0
			10	10		
2	C	6	Total	O	0	0
			6	6		
2	D	6	Total	O	0	0
			6	6		
2	E	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: uncharacterized protein Ta1441



• Molecule 1: uncharacterized protein Ta1441




• Molecule 1: uncharacterized protein Ta1441






• Molecule 1: uncharacterized protein Ta1441



• Molecule 1: uncharacterized protein Ta1441



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	184.32Å 184.32Å 260.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	101.02 – 2.70 45.37 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.6 (101.02-2.70) 99.6 (45.37-2.70)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.5.0054	Depositor
R, R_{free}	0.196 , 0.256 0.206 , 0.264	Depositor DCC
R_{free} test set	2360 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	65.7	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9299	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.13	8/1888 (0.4%)	0.98	4/2558 (0.2%)
1	B	1.07	3/1880 (0.2%)	0.96	1/2548 (0.0%)
1	C	1.00	3/1888 (0.2%)	0.97	3/2558 (0.1%)
1	D	1.05	6/1879 (0.3%)	0.93	2/2543 (0.1%)
1	E	0.97	2/1888 (0.1%)	0.95	5/2558 (0.2%)
All	All	1.05	22/9423 (0.2%)	0.96	15/12765 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	234	GLU	CG-CD	9.39	1.66	1.51
1	D	175	CYS	CB-SG	-9.08	1.66	1.82
1	D	13	GLU	CG-CD	8.89	1.65	1.51
1	D	234	GLU	CB-CG	8.80	1.68	1.52
1	A	234	GLU	CG-CD	8.45	1.64	1.51

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	19	LEU	CA-CB-CG	7.95	133.59	115.30
1	A	19	LEU	CA-CB-CG	7.90	133.48	115.30
1	B	19	LEU	CA-CB-CG	7.62	132.84	115.30
1	E	19	LEU	CA-CB-CG	7.60	132.77	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	19	LEU	CA-CB-CG	6.73	130.79	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	7	VAL	Peptide
1	C	135	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1856	0	1903	113	1
1	B	1848	0	1894	84	2
1	C	1856	0	1903	115	3
1	D	1848	0	1892	90	1
1	E	1856	0	1903	116	2
2	A	10	0	0	2	0
2	B	10	0	0	2	0
2	C	6	0	0	2	0
2	D	6	0	0	0	0
2	E	3	0	0	0	0
All	All	9299	0	9495	482	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 482 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:LYS:HZ3	1:A:10:LYS:HB3	1.02	1.17
1:D:10:LYS:HB3	1:D:10:LYS:NZ	1.60	1.14
1:C:177:VAL:HG11	1:D:60:ILE:HD11	1.16	1.11
1:B:177:VAL:HG11	1:C:60:ILE:HD11	1.17	1.10
1:D:10:LYS:HZ3	1:D:10:LYS:HB3	0.98	1.10

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:24:TYR:OH	1:C:9:LYS:NZ[4_556]	1.67	0.53
1:D:131:ASN:ND2	1:E:245:LYS:O[18_655]	1.82	0.38
1:E:9:LYS:NZ	1:E:20:GLU:OE2[4_556]	1.87	0.33
1:B:5:ILE:CD1	1:C:105:GLU:OE1[4_556]	2.07	0.13
1:A:13:GLU:OE1	1:C:177:VAL:O[4_556]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/252 (96%)	224 (93%)	16 (7%)	1 (0%)	38	66
1	B	240/252 (95%)	216 (90%)	23 (10%)	1 (0%)	38	66
1	C	241/252 (96%)	211 (88%)	23 (10%)	7 (3%)	5	13
1	D	238/252 (94%)	216 (91%)	20 (8%)	2 (1%)	22	49
1	E	241/252 (96%)	216 (90%)	21 (9%)	4 (2%)	11	27
All	All	1201/1260 (95%)	1083 (90%)	103 (9%)	15 (1%)	15	37

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	130	ALA
1	C	136	GLU
1	A	135	GLU
1	C	10	LYS
1	C	191	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/214 (96%)	169 (82%)	36 (18%)	2	5
1	B	204/214 (95%)	156 (76%)	48 (24%)	1	2
1	C	205/214 (96%)	165 (80%)	40 (20%)	1	4
1	D	204/214 (95%)	170 (83%)	34 (17%)	2	6
1	E	205/214 (96%)	165 (80%)	40 (20%)	1	4
All	All	1023/1070 (96%)	825 (81%)	198 (19%)	1	4

5 of 198 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	14	SER
1	C	145	LYS
1	E	131	ASN
1	C	19	LEU
1	C	69	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	58	HIS
1	C	222	GLN
1	E	112	ASN
1	C	116	GLN
1	D	15	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	243/252 (96%)	0.07	5 (2%) 64 65	22, 40, 73, 93	0
1	B	242/252 (96%)	0.14	8 (3%) 47 46	32, 43, 63, 92	0
1	C	243/252 (96%)	0.18	9 (3%) 42 41	35, 48, 73, 88	0
1	D	242/252 (96%)	0.26	8 (3%) 47 46	20, 48, 78, 95	0
1	E	243/252 (96%)	0.26	13 (5%) 27 25	33, 47, 73, 85	0
All	All	1213/1260 (96%)	0.18	43 (3%) 44 44	20, 45, 73, 95	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	218	LEU	5.9
1	B	7	VAL	5.5
1	C	136	GLU	5.5
1	E	136	GLU	4.6
1	E	132	GLY	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.