



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 05:26 am GMT

PDB ID : 4GBE  
Title : Crystal Structure of E. coli DNA Adenine Methyltransferase in Complex with SAH  
Authors : Harmer, J.E.; Roach, P.L.  
Deposited on : 2012-07-27  
Resolution : 2.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3491 (2.70-2.62)
Clashscore	112137	1026 (2.68-2.64)
Ramachandran outliers	110173	1010 (2.68-2.64)
Sidechain outliers	110143	1010 (2.68-2.64)
RSRZ outliers	101464	3511 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	284	<div> <div> <div>2%</div> <div>73%</div> <div>12%</div> <div>•</div> <div>14%</div> </div> </div>
1	E	284	<div> <div> <div>2%</div> <div>75%</div> <div>9%</div> <div>••</div> <div>14%</div> </div> </div>
1	F	284	<div> <div> <div>2%</div> <div>71%</div> <div>13%</div> <div>•</div> <div>14%</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6192 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA adenine methylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	243	Total	C	N	O	S	0	0	0
			2008	1300	338	362	8			
1	E	244	Total	C	N	O	S	0	0	0
			2016	1307	339	362	8			
1	F	244	Total	C	N	O	S	3	0	0
			2015	1305	339	363	8			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	HIS	-	EXPRESSION TAG	UNP P0AEE8
D	-4	HIS	-	EXPRESSION TAG	UNP P0AEE8
D	-3	HIS	-	EXPRESSION TAG	UNP P0AEE8
D	-2	HIS	-	EXPRESSION TAG	UNP P0AEE8
D	-1	HIS	-	EXPRESSION TAG	UNP P0AEE8
D	0	HIS	-	EXPRESSION TAG	UNP P0AEE8
E	-5	HIS	-	EXPRESSION TAG	UNP P0AEE8
E	-4	HIS	-	EXPRESSION TAG	UNP P0AEE8
E	-3	HIS	-	EXPRESSION TAG	UNP P0AEE8
E	-2	HIS	-	EXPRESSION TAG	UNP P0AEE8
E	-1	HIS	-	EXPRESSION TAG	UNP P0AEE8
E	0	HIS	-	EXPRESSION TAG	UNP P0AEE8
F	-5	HIS	-	EXPRESSION TAG	UNP P0AEE8
F	-4	HIS	-	EXPRESSION TAG	UNP P0AEE8
F	-3	HIS	-	EXPRESSION TAG	UNP P0AEE8
F	-2	HIS	-	EXPRESSION TAG	UNP P0AEE8
F	-1	HIS	-	EXPRESSION TAG	UNP P0AEE8
F	0	HIS	-	EXPRESSION TAG	UNP P0AEE8

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	E	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	F	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	31	Total	O	0	0
			31	31		
3	E	17	Total	O	0	0
			17	17		
3	F	27	Total	O	0	0
			27	27		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.18Å 161.18Å 95.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.02 – 2.66 29.02 – 2.66	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.02-2.66) 99.3 (29.02-2.66)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.27 (at 2.64Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.196 , 0.239 0.188 , 0.229	Depositor DCC
$R_{free}$ test set	2092 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	53.6	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 37.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6192	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	D	1.15	0/2063	0.94	0/2794
1	E	1.07	3/2071 (0.1%)	0.91	2/2804 (0.1%)
1	F	1.09	5/2070 (0.2%)	0.93	4/2804 (0.1%)
All	All	1.10	8/6204 (0.1%)	0.92	6/8402 (0.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	111	PHE	CD1-CE1	6.84	1.52	1.39
1	F	162	CYS	CB-SG	5.47	1.91	1.82
1	E	204	GLU	CG-CD	5.46	1.60	1.51
1	E	66	VAL	CB-CG2	-5.41	1.41	1.52
1	F	211	GLU	CB-CG	-5.40	1.41	1.52
1	F	84	GLU	CG-CD	5.32	1.59	1.51
1	E	218	GLU	CG-CD	5.21	1.59	1.51
1	F	204	GLU	CG-CD	5.15	1.59	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	229	ASP	CB-CG-OD1	6.52	124.17	118.30
1	F	69	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	E	31	LEU	CA-CB-CG	5.77	128.56	115.30
1	E	58	ASP	CB-CG-OD1	5.49	123.24	118.30
1	F	234	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	F	128	ARG	NE-CZ-NH2	-5.42	117.59	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2008	0	1944	25	0
1	E	2016	0	1960	22	0
1	F	2015	0	1953	16	0
2	D	26	0	17	2	0
2	E	26	0	18	0	0
2	F	26	0	17	0	0
3	D	31	0	0	0	0
3	E	17	0	0	0	0
3	F	27	0	0	0	0
All	All	6192	0	5909	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:139:LYS:HD2	1:D:139:LYS:H	1.06	1.08
1:D:139:LYS:CD	1:D:139:LYS:H	1.65	1.05
1:D:15:TYR:CD1	1:D:16:PRO:HD3	1.92	1.04
1:D:139:LYS:HD2	1:D:139:LYS:N	1.71	1.03
1:E:261:VAL:O	1:E:262:ASP:HB2	1.77	0.83
1:D:231:MET:HE1	1:E:122:LEU:HD13	1.74	0.69
1:D:139:LYS:N	1:D:139:LYS:CD	2.40	0.66
1:D:15:TYR:CD1	1:D:16:PRO:CD	2.77	0.66
1:E:67:LYS:HE3	1:E:155:ALA:O	1.96	0.64
1:F:124:ARG:CG	1:F:124:ARG:HH11	2.11	0.64
1:F:15:TYR:HB3	1:F:16:PRO:HD3	1.83	0.61
1:F:124:ARG:HH11	1:F:124:ARG:HG2	1.66	0.61
1:D:78:ARG:HH21	1:D:145:GLU:CD	2.04	0.60
1:D:231:MET:CE	1:E:122:LEU:HD13	2.32	0.60
1:D:54:ASP:OD2	2:D:301:SAH:O3'	2.20	0.60
1:D:231:MET:CE	1:E:122:LEU:CD1	2.82	0.58
1:E:247:VAL:HG22	1:E:260:LYS:NZ	2.19	0.58
1:D:80:LEU:HD22	1:D:109:VAL:HG12	1.84	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:246:LYS:O	1:F:247:VAL:HG13	2.06	0.56
1:E:78:ARG:NH1	1:E:145:GLU:OE2	2.38	0.56
1:E:246:LYS:O	1:E:247:VAL:HB	2.08	0.54
1:D:3:LYS:HG3	1:D:3:LYS:O	2.07	0.54
1:E:139:LYS:HG3	1:E:140:LYS:H	1.76	0.51
1:D:96:GLU:HG3	1:D:100:LYS:NZ	2.26	0.51
1:E:124:ARG:HG3	1:E:124:ARG:HH11	1.75	0.50
1:E:247:VAL:HG22	1:E:260:LYS:HZ3	1.77	0.50
1:D:177:VAL:HG22	1:D:222:PRO:HG2	1.95	0.49
1:D:20:ASP:OD1	1:D:23:ARG:NH1	2.44	0.48
1:D:231:MET:HE1	1:E:122:LEU:CD1	2.39	0.48
1:E:139:LYS:CG	1:E:140:LYS:H	2.28	0.47
1:D:12:GLY:HA3	2:D:301:SAH:HG1	1.96	0.47
1:D:242:LEU:HB3	1:D:264:LEU:HD11	1.98	0.46
1:D:17:LEU:HD21	1:D:263:GLU:HG2	1.97	0.45
1:E:174:ASP:N	1:E:174:ASP:OD1	2.48	0.45
1:F:261:VAL:HB	1:F:262:ASP:H	1.68	0.44
1:E:124:ARG:NH1	1:E:124:ARG:HG3	2.32	0.44
1:E:18:LEU:O	1:E:22:LYS:HG2	2.17	0.44
1:F:124:ARG:NH1	1:F:124:ARG:CG	2.77	0.43
1:D:143:PHE:CE2	1:D:145:GLU:HB2	2.54	0.43
1:F:117:TYR:CE2	1:F:141:PRO:HG2	2.53	0.43
1:D:14:LYS:HE2	1:D:181:ASP:OD2	2.19	0.42
1:E:187:LEU:HD21	1:E:228:HIS:CD2	2.54	0.42
1:F:218:GLU:O	1:F:218:GLU:HG2	2.18	0.42
1:E:31:LEU:HG	1:E:47:PHE:CE1	2.55	0.42
1:F:230:THR:O	1:F:234:ARG:HG3	2.20	0.42
1:D:15:TYR:CG	1:D:16:PRO:HD3	2.51	0.41
1:F:117:TYR:CZ	1:F:141:PRO:HG2	2.56	0.41
1:F:180:CYS:HB2	1:F:225:ILE:HG12	2.01	0.41
1:F:18:LEU:HD23	1:F:18:LEU:HA	1.94	0.41
1:D:14:LYS:HD3	1:D:14:LYS:HA	1.88	0.41
1:F:241:LYS:HA	1:F:241:LYS:HD3	1.72	0.41
1:D:10:TRP:CB	1:D:38:ALA:HB2	2.51	0.41
1:F:103:ASP:C	1:F:103:ASP:OD1	2.58	0.41
1:E:239:ARG:HH11	1:E:239:ARG:HG3	1.84	0.41
1:F:39:GLY:O	1:F:42:PHE:HB3	2.21	0.41
1:F:52:LEU:O	1:F:160:PHE:HA	2.21	0.41
1:E:52:LEU:O	1:E:160:PHE:HA	2.21	0.40
1:E:55:ILE:HA	1:E:55:ILE:HD12	1.86	0.40
1:E:14:LYS:HD3	1:E:14:LYS:HA	1.88	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	237/284 (84%)	221 (93%)	14 (6%)	2 (1%)	22	34
1	E	238/284 (84%)	224 (94%)	12 (5%)	2 (1%)	22	34
1	F	238/284 (84%)	221 (93%)	14 (6%)	3 (1%)	14	22
All	All	713/852 (84%)	666 (93%)	40 (6%)	7 (1%)	18	28

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	139	LYS
1	D	13	GLY
1	F	137	ARG
1	E	262	ASP
1	F	139	LYS
1	F	238	GLN
1	D	15	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	211/245 (86%)	198 (94%)	13 (6%)	21	34
1	E	212/245 (86%)	203 (96%)	9 (4%)	34	53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	212/245 (86%)	200 (94%)	12 (6%)	24	38
All	All	635/735 (86%)	601 (95%)	34 (5%)	26	41

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	3	LYS
1	D	18	LEU
1	D	40	SER
1	D	48	SER
1	D	55	ILE
1	D	80	LEU
1	D	137	ARG
1	D	139	LYS
1	D	187	LEU
1	D	226	SER
1	D	234	ARG
1	D	246	LYS
1	D	269	LYS
1	E	31	LEU
1	E	70	THR
1	E	128	ARG
1	E	139	LYS
1	E	140	LYS
1	E	174	ASP
1	E	226	SER
1	E	260	LYS
1	E	262	ASP
1	F	14	LYS
1	F	17	LEU
1	F	48	SER
1	F	49	ARG
1	F	132	ASN
1	F	174	ASP
1	F	187	LEU
1	F	199	ASN
1	F	218	GLU
1	F	247	VAL
1	F	261	VAL
1	F	269	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	D	4	ASN
1	D	120	ASN
1	D	157	ASN
1	E	150	HIS
1	E	228	HIS
1	F	120	ASN
1	F	132	ASN
1	F	199	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SAH	D	301	-	20,28,28	2.07	8 (40%)	20,40,40	3.17	5 (25%)
2	SAH	E	301	-	20,28,28	2.05	8 (40%)	20,40,40	3.37	7 (35%)
2	SAH	F	301	-	20,28,28	2.03	8 (40%)	20,40,40	3.46	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	D	301	-	-	0/7/31/31	0/3/3/3
2	SAH	E	301	-	-	0/7/31/31	0/3/3/3
2	SAH	F	301	-	-	0/7/31/31	0/3/3/3

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	301	SAH	C8-N7	-5.09	1.25	1.34
2	E	301	SAH	C8-N7	-4.99	1.25	1.34
2	F	301	SAH	C8-N7	-4.34	1.26	1.34
2	F	301	SAH	C2'-C1'	-3.88	1.47	1.53
2	E	301	SAH	O3'-C3'	-3.17	1.35	1.43
2	E	301	SAH	C2'-C1'	-3.01	1.48	1.53
2	F	301	SAH	O3'-C3'	-2.97	1.36	1.43
2	F	301	SAH	O2'-C2'	-2.79	1.36	1.43
2	D	301	SAH	O3'-C3'	-2.77	1.36	1.43
2	D	301	SAH	O2'-C2'	-2.67	1.36	1.43
2	F	301	SAH	C2'-C3'	-2.27	1.47	1.53
2	D	301	SAH	C2'-C3'	-2.26	1.47	1.53
2	D	301	SAH	C3'-C4'	-2.23	1.47	1.53
2	D	301	SAH	C5-N7	-2.16	1.32	1.39
2	F	301	SAH	C5-N7	-2.12	1.32	1.39
2	E	301	SAH	C3'-C4'	-2.10	1.47	1.53
2	E	301	SAH	C5-N7	-2.06	1.32	1.39
2	F	301	SAH	C2-N3	2.17	1.35	1.32
2	E	301	SAH	C6-N6	2.41	1.43	1.34
2	F	301	SAH	C6-N6	2.48	1.44	1.34
2	E	301	SAH	C5-C4	2.48	1.46	1.40
2	D	301	SAH	C2-N3	2.57	1.36	1.32
2	E	301	SAH	C2-N3	2.58	1.36	1.32
2	D	301	SAH	C5-C4	3.23	1.47	1.40

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	301	SAH	N3-C2-N1	-12.12	118.30	128.86
2	E	301	SAH	N3-C2-N1	-10.93	119.34	128.86
2	D	301	SAH	N3-C2-N1	-8.94	121.07	128.86

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	301	SAH	C4-C5-N7	-6.95	102.70	109.41
2	D	301	SAH	C4'-O4'-C1'	-6.36	103.00	109.77
2	D	301	SAH	C4-C5-N7	-6.34	103.28	109.41
2	E	301	SAH	C4'-O4'-C1'	-5.90	103.49	109.77
2	E	301	SAH	C4-C5-N7	-5.88	103.73	109.41
2	F	301	SAH	C4'-O4'-C1'	-3.15	106.42	109.77
2	E	301	SAH	C1'-N9-C4	-2.52	122.27	126.64
2	F	301	SAH	O2'-C2'-C1'	-2.20	104.73	111.61
2	D	301	SAH	O4'-C4'-C3'	2.03	109.19	105.17
2	F	301	SAH	CB-CG-SD	2.03	117.49	113.57
2	E	301	SAH	C2-N1-C6	2.04	122.34	118.77
2	E	301	SAH	CG-CB-CA	2.07	118.90	112.97
2	F	301	SAH	C2-N1-C6	2.55	123.23	118.77
2	F	301	SAH	C2'-C3'-C4'	2.83	108.13	102.62
2	E	301	SAH	C2'-C3'-C4'	3.00	108.46	102.62
2	D	301	SAH	CB-CG-SD	4.97	123.17	113.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	SAH	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	D	243/284 (85%)	-0.42	4 (1%) 72 72	26, 42, 69, 94	0
1	E	244/284 (85%)	-0.37	4 (1%) 72 72	30, 46, 69, 90	0
1	F	244/284 (85%)	-0.28	6 (2%) 58 56	28, 45, 68, 82	1 (0%)
All	All	731/852 (85%)	-0.36	14 (1%) 67 66	26, 44, 69, 94	1 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	261	VAL	5.2
1	D	199	ASN	4.9
1	F	261	VAL	3.9
1	F	199	ASN	3.9
1	E	260	LYS	3.2
1	F	247	VAL	3.0
1	F	179	TYR	2.7
1	D	261	VAL	2.6
1	D	16	PRO	2.5
1	D	15	TYR	2.5
1	E	89	GLU	2.0
1	F	218	GLU	2.0
1	F	139	LYS	2.0
1	E	246	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SAH	F	301	26/26	0.97	0.16	-0.32	31,36,54,55	0
2	SAH	E	301	26/26	0.96	0.12	-0.47	32,38,57,61	0
2	SAH	D	301	26/26	0.96	0.14	-0.78	40,47,70,71	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.