



# Full wwPDB X-ray Structure Validation Report i

Feb 13, 2017 – 05:07 am GMT

PDB ID : 4GCW  
Title : Crystal structure of RNase Z in complex with precursor tRNA(Thr)  
Authors : Pellegrini, O.; Li de la Sierra-Gallay, I.; Piton, J.; Gilet, L.; Condon, C.  
Deposited on : 2012-07-31  
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1692 (3.00-3.00)
Clashscore	112137	2037 (3.00-3.00)
Ramachandran outliers	110173	1973 (3.00-3.00)
Sidechain outliers	110143	1976 (3.00-3.00)
RSRZ outliers	101464	1716 (3.00-3.00)
RNA backbone	2435	1007 (3.34-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 3618 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease Z.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	307	Total	C 2400	N 1527	O 410	S 456	7	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	MET	ILE	ENGINEERED MUTATION	UNP P54548
A	228	MET	LEU	ENGINEERED MUTATION	UNP P54548
A	308	LYS	-	EXPRESSION TAG	UNP P54548
A	309	LEU	-	EXPRESSION TAG	UNP P54548
A	310	ALA	-	EXPRESSION TAG	UNP P54548
A	311	ALA	-	EXPRESSION TAG	UNP P54548
A	312	ALA	-	EXPRESSION TAG	UNP P54548
A	313	LEU	-	EXPRESSION TAG	UNP P54548
A	314	GLU	-	EXPRESSION TAG	UNP P54548
A	315	HIS	-	EXPRESSION TAG	UNP P54548
A	316	HIS	-	EXPRESSION TAG	UNP P54548
A	317	HIS	-	EXPRESSION TAG	UNP P54548
A	318	HIS	-	EXPRESSION TAG	UNP P54548
A	319	HIS	-	EXPRESSION TAG	UNP P54548
A	320	HIS	-	EXPRESSION TAG	UNP P54548


- Molecule 2 is a RNA chain called TRNA(THR).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	55	Total	C 1172	N 525	O 209	P 384	54	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total    Zn 2      2	0	0

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total    C    N    O    S 12    6    1    4    1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	26	Total    O 26    26	0	0
5	B	6	Total    O 6      6	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribonuclease Z



- Molecule 2: tRNA(THR)



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	174.08Å    42.55Å    110.89Å 90.00°    127.18°    90.00°	Depositor
Resolution (Å)	43.30 – 3.00 42.44 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (43.30-3.00) 99.1 (42.44-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle^1$	2.19 (at 3.01Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
$R$ , $R_{free}$	0.227 , 0.256 0.244 , 0.281	Depositor DCC
$R_{free}$ test set	662 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.5	Xtriage
Anisotropy	0.668	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 56.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.024 for -h-2*l,-k,l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	3618	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.39% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MES, OMU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.36	0/2447	0.60	0/3309
2	B	0.90	0/1262	1.52	20/1963 (1.0%)
All	All	0.60	0/3709	1.04	20/5272 (0.4%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	9	A	P-O3'-C3'	11.04	132.94	119.70
2	B	12	U	P-O3'-C3'	10.54	132.35	119.70
2	B	21	A	P-O3'-C3'	9.71	131.35	119.70
2	B	11	C	P-O3'-C3'	9.50	131.09	119.70
2	B	46	G	P-O3'-C3'	9.41	131.00	119.70
2	B	13	C	P-O3'-C3'	7.33	128.50	119.70
2	B	3	U	O4'-C1'-N1	6.08	113.06	108.20
2	B	15	G	P-O3'-C3'	5.85	126.72	119.70
2	B	47	U	C3'-C2'-C1'	5.63	106.00	101.50
2	B	25	C	O4'-C1'-N1	5.55	112.64	108.20
2	B	62	C	O4'-C1'-N1	5.50	112.60	108.20
2	B	51	C	O4'-C1'-N1	5.30	112.44	108.20
2	B	56	C	O4'-C1'-N1	5.30	112.44	108.20
2	B	65	U	O4'-C1'-N1	5.24	112.39	108.20
2	B	13	C	O4'-C1'-N1	5.17	112.34	108.20
2	B	10	G	N9-C1'-C2'	5.16	120.70	114.00
2	B	72	C	O4'-C1'-N1	5.09	112.27	108.20
2	B	19	G	P-O3'-C3'	5.07	125.79	119.70
2	B	64	C	O4'-C1'-N1	5.02	112.22	108.20
2	B	12	U	O4'-C1'-N1	5.02	112.22	108.20

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2400	0	2413	18	0
2	B	1172	0	597	1	0
3	A	2	0	0	0	0
4	A	12	0	13	1	0
5	A	26	0	0	0	0
5	B	6	0	0	0	0
All	All	3618	0	3023	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:CYS:HB3	1:A:72:LEU:HD22	1.81	0.62
1:A:122:GLU:HG3	1:A:136:VAL:HA	1.82	0.60
1:A:87:GLU:HG3	1:A:114:PRO:HB2	1.94	0.49
1:A:53:PRO:HA	1:A:56:ILE:HD12	1.94	0.49
1:A:207:VAL:HG21	1:A:222:ALA:HA	1.96	0.48
1:A:168:ILE:HG23	1:A:185:LEU:HD21	1.95	0.48
1:A:61:ILE:HD11	1:A:72:LEU:HD21	1.95	0.47
1:A:140:HIS:HA	1:A:246:TYR:HB3	1.97	0.47
2:B:72:C:H5"	2:B:73:OMU:H5	1.95	0.47
1:A:257:ALA:HB2	1:A:265:LEU:HD13	1.96	0.46
1:A:178:LYS:HA	1:A:197:LEU:HD21	1.98	0.45
1:A:33:VAL:HG21	1:A:51:ILE:HG23	1.99	0.45
1:A:209:SER:HB2	1:A:218:LEU:HD21	1.97	0.45
1:A:32:SER:HB3	4:A:403:MES:H62	1.99	0.44
1:A:104:LEU:HD23	1:A:111:LEU:HD21	2.00	0.44
1:A:177:ILE:HG23	1:A:193:GLY:HA2	2.01	0.42
1:A:194:ASN:HA	1:A:197:LEU:HD12	2.01	0.41
1:A:52:LYS:HD2	1:A:55:LYS:HE3	2.02	0.41
1:A:44:HIS:HA	1:A:47:LEU:HD12	2.03	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	305/320 (95%)	288 (94%)	16 (5%)	1 (0%)	44 81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	GLY

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	258/268 (96%)	247 (96%)	11 (4%)	33 72

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	20	THR
1	A	58	LYS
1	A	72	LEU
1	A	88	LEU
1	A	109	THR
1	A	115	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	158	LEU
1	A	185	LEU
1	A	214	VAL
1	A	231	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	53/76 (69%)	24 (45%)	0

All (24) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	2	C
2	B	6	C
2	B	10	G
2	B	11	C
2	B	12	U
2	B	13	C
2	B	14	A
2	B	16	C
2	B	17	A
2	B	20	U
2	B	21	A
2	B	22	G
2	B	23	A
2	B	25	C
2	B	26	A
2	B	47	U
2	B	48	C
2	B	49	A
2	B	60	C
2	B	61	C
2	B	63	G
2	B	67	G
2	B	73	OMU
2	B	74	OMU

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	OMU	B	73	2	14,22,23	1.35	2 (14%)	18,31,34	4.12	6 (33%)
2	OMU	B	74	2	14,22,23	1.39	2 (14%)	18,31,34	3.71	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OMU	B	73	2	-	0/5/27/28	0/2/2/2
2	OMU	B	74	2	-	0/5/27/28	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	74	OMU	C6-N1	2.92	1.39	1.35
2	B	73	OMU	C6-N1	3.00	1.39	1.35
2	B	73	OMU	C4-N3	3.46	1.39	1.33
2	B	74	OMU	C4-N3	3.50	1.39	1.33

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	73	OMU	C5-C4-N3	-3.47	114.82	123.12
2	B	74	OMU	C5-C4-N3	-3.35	115.12	123.12
2	B	74	OMU	CM2-O2'-C2'	2.03	120.09	114.54
2	B	73	OMU	O2'-C2'-C3'	2.38	117.51	111.21
2	B	73	OMU	C4'-O4'-C1'	2.42	112.35	109.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	73	OMU	O2'-C2'-C1'	4.15	117.23	108.75
2	B	73	OMU	CM2-O2'-C2'	4.26	126.19	114.54
2	B	74	OMU	C4-N3-C2	14.98	126.99	114.13
2	B	73	OMU	C4-N3-C2	15.31	127.28	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	73	OMU	1	0

## 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MES	A	403	-	12,12,12	0.68	0	14,16,16	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	A	403	-	-	0/6/14/14	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	MES	1	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/320 (95%)	0.23	8 (2%) 56 27	44, 63, 90, 105	0
2	B	53/76 (69%)	1.12	16 (30%) 1 0	62, 117, 154, 155	0
All	All	360/396 (90%)	0.36	24 (6%) 19 7	44, 66, 129, 155	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	46	G	4.9
2	B	16	C	4.8
2	B	15	G	4.5
2	B	47	U	3.9
2	B	9	A	3.7
1	A	240	LYS	3.2
2	B	13	C	3.2
1	A	237	GLU	3.2
1	A	242	ALA	3.1
2	B	22	G	3.1
2	B	25	C	3.1
2	B	8	U	3.0
1	A	236	LYS	3.0
2	B	26	A	2.9
2	B	23	A	2.9
1	A	239	ARG	2.7
2	B	10	G	2.6
2	B	24	G	2.6
2	B	17	A	2.4
1	A	241	LEU	2.4
2	B	21	A	2.3
1	A	175	GLN	2.3
2	B	14	A	2.1
1	A	235	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	OMU	B	73	21/22	0.79	0.33	-	106,109,113,113	0
2	OMU	B	74	21/22	0.91	0.26	-	105,107,110,110	0

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MES	A	403	12/12	0.87	0.35	1.13	110,111,116,116	0
3	ZN	A	402	1/1	0.96	0.21	0.90	68,68,68,68	0
3	ZN	A	401	1/1	0.99	0.18	-0.90	70,70,70,70	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.