



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2017 – 07:22 AM EDT

PDB ID : 3GG0  
Title : Klebsiella pneumoniae BlrP1 pH 9.0 manganese/cy-diGMP complex  
Authors : Barends, T.; Hartmann, E.; Griesse, J.; Beitlich, T.; Kirienko, N.; Ryjenkov, D.; Reinstein, J.; Shoeman, R.; Gomelsky, M.; Schlichting, I.  
Deposited on : unknown  
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4993 (2.58-2.50)
Clashscore	112137	5755 (2.58-2.50)
Ramachandran outliers	110173	5652 (2.58-2.50)
Sidechain outliers	110143	5654 (2.58-2.50)
RSRZ outliers	101464	5026 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	<div> <div>2%</div> <div>86%</div> <div>8%</div> <div>5%</div> </div>
1	B	413	<div> <div>6%</div> <div>84%</div> <div>11%</div> <div>5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	C2E	A	501	X	-	-	-
2	C2E	B	501	X	-	-	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6457 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called *Klebsiella pneumoniae* BlrP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	0	0
			3075	1970	532	555	18			
1	B	394	Total	C	N	O	S	0	0	0
			3094	1983	534	559	18			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	ILE	-	EXPRESSION TAG	UNP A6T8V8
A	-6	SER	-	EXPRESSION TAG	UNP A6T8V8
A	-5	GLU	-	EXPRESSION TAG	UNP A6T8V8
A	-4	PHE	-	EXPRESSION TAG	UNP A6T8V8
A	-3	GLY	-	EXPRESSION TAG	UNP A6T8V8
A	-2	SER	-	EXPRESSION TAG	UNP A6T8V8
A	-1	SER	-	EXPRESSION TAG	UNP A6T8V8
A	0	ARG	-	EXPRESSION TAG	UNP A6T8V8
B	-7	ILE	-	EXPRESSION TAG	UNP A6T8V8
B	-6	SER	-	EXPRESSION TAG	UNP A6T8V8
B	-5	GLU	-	EXPRESSION TAG	UNP A6T8V8
B	-4	PHE	-	EXPRESSION TAG	UNP A6T8V8
B	-3	GLY	-	EXPRESSION TAG	UNP A6T8V8
B	-2	SER	-	EXPRESSION TAG	UNP A6T8V8
B	-1	SER	-	EXPRESSION TAG	UNP A6T8V8
B	0	ARG	-	EXPRESSION TAG	UNP A6T8V8

- Molecule 2 is 9,9'-[(2R,3R,3aS,5S,7aR,9R,10R,10aS,12S,14aR)-3,5,10,12-tetrahydroxy-5,12-dioxidoctahydro-2H,7H-difuro[3,2-d:3',2'-j][1,3,7,9,2,8]tetraoxadiphosphacyclododecine-2,9-diyl]bis(2-amino-1,9-dihydro-6H-purin-6-one) (three-letter code: C2E) (formula: C<sub>20</sub>H<sub>24</sub>N<sub>10</sub>O<sub>14</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 46	C 20	N 10	O 14	P 2	0	0
2	B	1	Total 46	C 20	N 10	O 14	P 2	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Mn 2 2	0	0
3	A	2	Total Mn 2 2	0	0

- Molecule 4 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula:  $\text{C}_{17}\text{H}_{21}\text{N}_4\text{O}_9\text{P}$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 17	N 4	O 9	P 1	0	0
4	B	1	Total 31	C 17	N 4	O 9	P 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	73	Total O 73 73	0	0
5	B	57	Total O 57 57	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: *Klebsiella pneumoniae* BlrP1



- Molecule 1: *Klebsiella pneumoniae* BlrP1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.02Å 96.89Å 126.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.45 – 2.55 48.45 – 2.55	Depositor EDS
% Data completeness (in resolution range)	97.4 (48.45-2.55) 97.4 (48.45-2.55)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.53 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.240 , 0.295 0.237 , 0.293	Depositor DCC
$R_{free}$ test set	1392 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.3	Xtriage
Anisotropy	0.787	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 25.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	6457	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, MN, C2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/3143	0.38	0/4259
1	B	0.34	0/3163	0.38	0/4287
All	All	0.35	0/6306	0.38	0/8546

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3075	0	3095	17	0
1	B	3094	0	3110	18	0
2	A	46	0	19	3	0
2	B	46	0	19	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	31	0	19	0	0
4	B	31	0	19	0	0
5	A	73	0	0	1	0
5	B	57	0	0	0	0
All	All	6457	0	6281	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:VAL:HG21	1:A:94:ILE:HB	1.75	0.68
1:B:32:LEU:HD13	1:B:90:THR:HG23	1.79	0.65
2:A:501:C2E:O2A	5:A:408:HOH:O	2.14	0.64
1:A:229:LEU:N	1:A:230:PRO:HA	2.17	0.58
1:A:352:LEU:HD21	1:B:337:THR:HA	1.88	0.56
1:B:245:LEU:HB3	1:B:288:VAL:HG11	1.88	0.56
1:B:30:LYS:NZ	1:B:64:GLU:OE1	2.39	0.55
1:A:143:GLU:N	1:A:144:PRO:HD2	2.22	0.55
1:A:105:ALA:O	1:A:109:ILE:HG12	2.11	0.51
1:A:276:THR:HG22	1:A:307:GLY:HA3	1.93	0.50
1:A:309:SER:HA	1:B:309:SER:HA	1.92	0.50
1:A:321:LYS:HE2	1:A:355:THR:HG21	1.93	0.49
1:A:307:GLY:O	1:A:309:SER:N	2.39	0.48
1:B:275:GLU:CD	1:B:309:SER:HB2	2.34	0.47
1:B:270:LEU:HD12	1:B:298:LYS:HB3	1.96	0.47
1:A:229:LEU:H	1:A:230:PRO:HA	1.78	0.47
1:A:6:ILE:HD11	1:A:125:MET:HG3	1.97	0.47
1:B:286:ARG:HA	1:B:289:LEU:HD12	1.97	0.47
1:B:329:VAL:HG13	1:B:370:LEU:HD11	1.97	0.47
1:A:324:VAL:O	1:A:359:GLU:HB2	2.16	0.46
1:A:46:GLN:HE22	1:A:137:GLY:HA2	1.81	0.46
1:A:36:ILE:HD13	1:A:61:LEU:HD22	1.97	0.45
1:A:226:ALA:HB1	1:A:231:LEU:HD22	1.99	0.45
1:B:141:LEU:HA	1:B:142:PRO:HD3	1.89	0.43
1:B:231:LEU:HD12	1:B:268:GLN:HE22	1.84	0.43
2:A:501:C2E:O2P	2:A:501:C2E:H2A	2.19	0.42
1:B:306:ALA:HB2	1:B:325:ASP:CG	2.39	0.42
1:B:333:HIS:CD2	1:B:334:ILE:HG23	2.54	0.42
1:A:270:LEU:HD21	1:A:300:ALA:HB2	2.01	0.42
1:B:189:ALA:HB3	1:B:238:ILE:HG22	2.02	0.41
1:B:40:LEU:HD13	1:B:49:GLN:HB2	2.01	0.41
1:B:330:ARG:HA	1:B:330:ARG:HD3	1.80	0.41
1:A:174:GLN:OE1	2:A:501:C2E:H8	2.20	0.41
1:B:208:ALA:HA	1:B:211:ARG:HD2	2.03	0.41
1:B:176:ILE:HG12	1:B:379:GLN:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/413 (94%)	373 (97%)	13 (3%)	0	100	100
1	B	390/413 (94%)	376 (96%)	11 (3%)	3 (1%)	22	38
All	All	776/826 (94%)	749 (96%)	24 (3%)	3 (0%)	38	57

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	281	CYS
1	B	230	PRO
1	B	232	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/346 (95%)	321 (97%)	9 (3%)	50	75
1	B	332/346 (96%)	317 (96%)	15 (4%)	32	54
All	All	662/692 (96%)	638 (96%)	24 (4%)	40	64

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	71	HIS
1	A	238	ILE
1	A	267	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	274	THR
1	A	276	THR
1	A	283	ASP
1	A	311	LEU
1	A	347	ARG
1	A	393	GLU
1	B	67	SER
1	B	71	HIS
1	B	88	HIS
1	B	95	LEU
1	B	231	LEU
1	B	235	GLN
1	B	249	PRO
1	B	264	LEU
1	B	283	ASP
1	B	299	LEU
1	B	309	SER
1	B	327	GLU
1	B	365	GLU
1	B	371	GLN
1	B	400	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	167	GLN
1	B	167	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	C2E	A	501	3	44,52,52	3.09	8 (18%)	50,82,82	2.13	14 (28%)
4	FMN	A	504	-	31,33,33	1.27	5 (16%)	38,50,50	1.37	5 (13%)
2	C2E	B	501	3	44,52,52	3.02	8 (18%)	50,82,82	2.08	14 (28%)
4	FMN	B	504	-	31,33,33	1.39	5 (16%)	38,50,50	1.38	4 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C2E	A	501	3	1/1/10/10	0/22/62/62	0/6/7/7
4	FMN	A	504	-	-	0/16/18/18	0/3/3/3
2	C2E	B	501	3	1/1/10/10	0/22/62/62	0/6/7/7
4	FMN	B	504	-	-	0/16/18/18	0/3/3/3

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	C2E	C2A-C1A	-13.50	1.32	1.53
2	B	501	C2E	C2A-C1A	-13.01	1.33	1.53
2	A	501	C2E	C2'-C1'	-12.17	1.34	1.53
2	B	501	C2E	C2'-C1'	-11.83	1.34	1.53
2	A	501	C2E	O2A-C2A	-4.30	1.33	1.43
2	B	501	C2E	O2'-C2'	-4.16	1.33	1.43
2	B	501	C2E	O2A-C2A	-4.13	1.33	1.43
2	A	501	C2E	O2'-C2'	-4.06	1.33	1.43
2	A	501	C2E	C8-N7	-3.11	1.28	1.34
2	B	501	C2E	C8-N7	-3.00	1.29	1.34
4	A	504	FMN	C5A-N5	2.01	1.38	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	504	FMN	C1'-N10	2.15	1.50	1.48
4	B	504	FMN	C5A-N5	2.39	1.39	1.35
2	A	501	C2E	O4A-C1A	2.42	1.44	1.41
4	A	504	FMN	C1'-N10	2.46	1.50	1.48
2	A	501	C2E	P11-O11	2.47	1.60	1.50
4	A	504	FMN	C4-N3	2.56	1.37	1.33
2	B	501	C2E	O4A-C1A	2.70	1.45	1.41
2	B	501	C2E	P11-O11	2.76	1.61	1.50
4	B	504	FMN	C4-N3	2.92	1.38	1.33
4	A	504	FMN	C10-N1	3.13	1.37	1.33
4	A	504	FMN	C4A-N5	3.20	1.37	1.33
2	B	501	C2E	O4'-C1'	3.39	1.45	1.41
4	B	504	FMN	C4A-N5	3.68	1.38	1.33
4	B	504	FMN	C10-N1	3.69	1.38	1.33
2	A	501	C2E	O4'-C1'	3.83	1.46	1.41

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	C2E	C4A-O4A-C1A	-5.69	103.71	109.77
2	B	501	C2E	C4A-O4A-C1A	-5.46	103.96	109.77
2	A	501	C2E	C4-C5-N7	-5.00	104.58	109.41
2	A	501	C2E	N31-C21-N11	-4.57	120.78	127.46
2	B	501	C2E	C4-C5-N7	-4.51	105.06	109.41
2	B	501	C2E	N3-C2-N1	-4.50	120.89	127.46
2	B	501	C2E	N31-C21-N11	-4.46	120.95	127.46
2	A	501	C2E	N3-C2-N1	-4.33	121.14	127.46
2	A	501	C2E	C5-C6-N1	-3.49	118.51	123.48
2	B	501	C2E	O5A-C5A-C4A	-3.33	97.18	109.00
2	A	501	C2E	O5A-C5A-C4A	-3.28	97.38	109.00
2	B	501	C2E	C5-C6-N1	-3.03	119.17	123.48
2	A	501	C2E	C51-C61-N11	-3.01	119.20	123.48
2	B	501	C2E	C51-C61-N11	-2.91	119.34	123.48
2	B	501	C2E	O5A-P11-O11	-2.82	97.85	109.25
2	A	501	C2E	O5A-P11-O11	-2.59	98.80	109.25
4	A	504	FMN	C4A-C4-N3	-2.57	119.82	123.48
4	B	504	FMN	C4A-C4-N3	-2.55	119.86	123.48
2	B	501	C2E	C1'-N9-C4	-2.42	122.45	126.64
4	A	504	FMN	C1'-N10-C9A	2.02	120.19	118.35
2	A	501	C2E	C3'-C2'-C1'	2.10	104.66	99.95
2	B	501	C2E	C3'-C2'-C1'	2.25	105.01	99.95
2	A	501	C2E	O21-P11-O5A	2.41	119.51	108.14

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	504	FMN	C5A-C9A-N10	2.65	119.62	117.66
4	B	504	FMN	C4A-N5-C5A	2.93	119.85	116.76
4	B	504	FMN	C5A-C9A-N10	3.05	119.92	117.66
2	B	501	C2E	C6-N1-C2	3.30	120.80	116.06
2	B	501	C2E	C61-N11-C21	3.37	120.91	116.06
2	B	501	C2E	C21-N31-C41	3.47	119.21	115.16
2	A	501	C2E	C61-N11-C21	3.48	121.06	116.06
2	A	501	C2E	C6-N1-C2	3.54	121.15	116.06
2	A	501	C2E	C21-N31-C41	3.57	119.33	115.16
4	A	504	FMN	C4A-N5-C5A	3.59	120.55	116.76
2	A	501	C2E	C2-N3-C4	3.62	119.39	115.16
2	B	501	C2E	C2-N3-C4	3.63	119.39	115.16
4	A	504	FMN	C4-N3-C2	5.32	119.81	115.16
4	B	504	FMN	C4-N3-C2	5.64	120.09	115.16

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	501	C2E	C1A
2	A	501	C2E	C1A

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	C2E	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	392/413 (94%)	0.38	10 (2%) 56 60	18, 29, 49, 58	0
1	B	394/413 (95%)	0.47	25 (6%) 21 22	19, 31, 47, 57	0
All	All	786/826 (95%)	0.43	35 (4%) 34 37	18, 30, 49, 58	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	308	TYR	8.5
1	A	308	TYR	7.4
1	B	310	GLY	5.2
1	B	312	SER	4.7
1	B	232	GLY	4.3
1	B	161	PRO	3.8
1	B	230	PRO	3.7
1	B	311	LEU	3.6
1	B	233	LYS	3.5
1	B	307	GLY	3.5
1	A	166	GLY	3.4
1	A	159	ALA	3.3
1	B	309	SER	3.3
1	B	282	PHE	3.2
1	B	231	LEU	2.8
1	B	14	ASP	2.8
1	B	286	ARG	2.7
1	B	160	ALA	2.6
1	A	282	PHE	2.6
1	B	13	PRO	2.5
1	B	102	THR	2.5
1	B	314	LEU	2.5
1	A	309	SER	2.5
1	B	90	THR	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	161	PRO	2.5
1	B	280	THR	2.5
1	A	306	ALA	2.4
1	B	285	PHE	2.3
1	B	316	ARG	2.2
1	A	294	VAL	2.2
1	B	265	ARG	2.1
1	A	247	HIS	2.1
1	B	152	MET	2.1
1	A	14	ASP	2.0
1	B	313	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	FMN	A	504	31/31	0.92	0.16	-0.08	13,14,35,35	0
4	FMN	B	504	31/31	0.88	0.16	-0.30	21,23,41,41	0
2	C2E	B	501	46/46	0.93	0.13	-0.65	19,25,26,27	0
2	C2E	A	501	46/46	0.95	0.14	-0.80	18,19,21,22	0
3	MN	A	502	1/1	0.99	0.07	-2.65	16,16,16,16	0
3	MN	B	503	1/1	0.95	0.06	-7.87	29,29,29,29	0
3	MN	A	503	1/1	0.97	0.09	-	17,17,17,17	0
3	MN	B	502	1/1	0.99	0.06	-	23,23,23,23	0

## 6.5 Other polymers

There are no such residues in this entry.