Full wwPDB X-ray Structure Validation Report (i)

Sep 26, 2017 – 06:41 AM EDT

PDB ID	:	1GHE
Title	:	CRYSTAL STRUCTURE OF TABTOXIN RESISTANCE PROTEIN COMPLEXED WITH AN ACYL COENZYME A
Authors	:	He, H.; Ding, Y.; Bartlam, M.; Sun, F.; Le, Y.; Qin, X.; Tang, H.; Zhang, R.; Joachimiak, A.; Liu, Y.; Zhao, N.; Rao, Z.
Deposited on	:	unknown
Resolution	:	1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

|		
MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percentile Ranks	Value
Clashscore		5
Ramachandran outliers		0
Sidechain outliers		1.5%

Worse ◄ ► *Better*

■ Percentile relative to all X-ray structures
▯ Percentile relative to X-ray structures of similar resolution

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	177	82% 13% · ·
1	B	177	89% 7% · ·

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLTRANSFERASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	170	1298	820	232	239	3	4	0	0	0
1	B	171	Total	C	N	O	S	Se	0	0	0
			1308	826	235	240	3	4			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP P16966
A	38	MSE	MET	MODIFIED RESIDUE	UNP P16966
A	43	MSE	MET	MODIFIED RESIDUE	UNP P16966
A	97	MSE	MET	MODIFIED RESIDUE	UNP P16966
A	112	MSE	MET	MODIFIED RESIDUE	UNP P16966
B	1	MSE	MET	MODIFIED RESIDUE	UNP P16966
B	38	MSE	MET	MODIFIED RESIDUE	UNP P16966
B	43	MSE	MET	MODIFIED RESIDUE	UNP P16966
B	97	MSE	MET	MODIFIED RESIDUE	UNP P16966
B	112	MSE	MET	MODIFIED RESIDUE	UNP P16966

- Molecule 2 is ACETYL COENZYME *A (three-letter code: ACO) (formula: $C_{23}H_{38}N_7O_{17}P_3S$).

ACO

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			51	23	7	17	3	1		

• Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	83	Total	O	0	0
			83	83		
3	B	79	Total	O	0	0
			79	79		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: ACETYLTRANSFERASE

Chain A:

- Molecule 1: ACETYLTRANSFERASE

Chain B:

4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	101.76Å 45.70Å 84.24Å 90.00° 105.79° 90.00°	Depositor
Resolution (Å)	30.00 – 1.55	Depositor
% Data completeness (in resolution range)	87.8 (30.00-1.55)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.209 , 0.230	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2870	wwPDB-VP
Average B, all atoms (Å^2)	17.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ACO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$|Z|$>5	RMSZ	#$|Z|$>5
1	A	0.48	0/1319	0.69	1/1780 (0.1%)
1	B	0.51	1/1330 (0.1%)	0.72	0/1795
All	All	0.50	1/2649 (0.0%)	0.70	1/3575 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	50	CYS	CB-SG	-5.59	1.72	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	67	VAL	N-CA-C	-5.71	95.57	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1298	0	1291	19	0
1	B	1308	0	1298	10	0
2	A	51	0	32	6	0
2	B	51	0	33	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	83	0	0	1	0
3	B	79	0	0	1	0
All	All	2870	0	2654	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:ARG:N	1:B:162:ARG:HD2	1.98	0.78
1:A:129:LEU:HB2	2:A:301:ACO:CH3	2.16	0.75
1:A:19:HIS:HD2	1:A:47:TYR:OH	1.71	0.72
1:A:132:GLU:HG3	1:A:164:HIS:CD2	2.26	0.70
1:A:129:LEU:HD13	2:A:301:ACO:HH32	1.76	0.66
1:B:19:HIS:HD2	1:B:47:TYR:OH	1.78	0.66
1:A:105:ARG:NH2	1:B:44:GLN:OE1	2.31	0.64
1:A:129:LEU:HB2	2:A:301:ACO:HH31	1.81	0.62
1:A:6:LEU:HD13	1:A:6:LEU:C	2.27	0.55
1:A:19:HIS:CD2	1:A:47:TYR:OH	2.55	0.55
1:A:129:LEU:HD13	2:A:301:ACO:CH3	2.37	0.55
1:A:38:MSE:HE1	1:A:161:GLY:HA2	1.89	0.53
1:B:158:THR:OG1	1:B:160:ASP:OD2	2.28	0.52
1:B:161:GLY:C	1:B:162:ARG:HD2	2.30	0.51
1:B:162:ARG:CD	1:B:162:ARG:N	2.72	0.51
1:B:128:HIS:HE1	3:B:310:HOH:O	1.94	0.51
1:B:43:MSE:HE3	1:B:47:TYR:CE1	2.46	0.50
1:A:67:VAL:HG11	1:A:107:LEU:HD22	1.92	0.50
1:A:98:VAL:HG23	2:A:301:ACO:H132	1.93	0.50
1:A:164:HIS:HD2	1:A:165:PRO:O	1.94	0.50
1:A:128:HIS:HD2	3:A:319:HOH:O	1.96	0.49
1:A:63:LEU:N	1:A:63:LEU:HD12	2.31	0.46
1:A:129:LEU:HB2	2:A:301:ACO:HH33	1.95	0.44
1:B:113:ASP:O	1:B:117:GLN:HG3	2.19	0.42
1:A:127:LEU:HG	1:A:173:LEU:HD11	2.01	0.41
1:A:160:ASP:OD2	1:A:162:ARG:NH1	2.54	0.41
1:A:89:ASN:ND2	1:A:125:GLY:H	2.18	0.41
1:B:38:MSE:HB2	1:B:40:ASP:OD2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/177 (95%)	166 (99%)	2 (1%)	0	100	100
1	B	169/177 (96%)	167 (99%)	2 (1%)	0	100	100
All	All	337/354 (95%)	333 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/132 (99%)	128 (98%)	3 (2%)	56	23
1	B	132/132 (100%)	131 (99%)	1 (1%)	85	70
All	All	263/264 (100%)	259 (98%)	4 (2%)	70	41

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	LYS
1	A	86	ASN
1	A	158	THR
1	B	162	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	HIS
1	A	72	ASN
1	A	83	GLN
1	A	86	ASN
1	A	89	ASN
1	A	94	GLN
1	A	117	GLN
1	A	128	HIS
1	A	164	HIS
1	B	19	HIS
1	B	72	ASN
1	B	94	GLN
1	B	122	HIS
1	B	128	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#$|Z| > 2$	Counts	RMSZ	#$|Z| > 2$
2	ACO	A	301	-	46,53,53	3.63	16 (34%)	53,79,79	3.41	27 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#|Z| > 2	Counts	RMSZ	#|Z| > 2
2	ACO	B	302	-	46,53,53	2.25	15 (32%)	53,79,79	2.85	28 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACO	A	301	-	-	0/47/67/67	0/3/3/3
2	ACO	B	302	-	-	0/47/67/67	0/3/3/3

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	ACO	O2B-C2B	-13.64	1.11	1.43
2	B	302	ACO	P3B-O3B	-4.13	1.52	1.59
2	B	302	ACO	C-S1P	-3.98	1.48	1.75
2	A	301	ACO	C5B-C4B	-3.97	1.39	1.51
2	B	302	ACO	C3B-C4B	-2.88	1.44	1.52
2	B	302	ACO	P2A-O5A	-2.73	1.41	1.55
2	A	301	ACO	P2A-O5A	-2.64	1.41	1.55
2	A	301	ACO	C9P-N8P	2.08	1.37	1.33
2	B	302	ACO	P2A-O6A	2.19	1.68	1.59
2	A	301	ACO	CH3-C	2.35	1.60	1.50
2	A	301	ACO	C2A-N1A	2.35	1.38	1.33
2	B	302	ACO	CCP-CBP	2.55	1.56	1.52
2	B	302	ACO	OAP-CAP	2.58	1.47	1.42
2	A	301	ACO	OAP-CAP	2.63	1.47	1.42
2	B	302	ACO	O4B-C1B	2.68	1.45	1.41
2	A	301	ACO	O5P-C5P	2.88	1.29	1.23
2	B	302	ACO	O5B-C5B	3.01	1.56	1.44
2	A	301	ACO	P3B-O3B	3.07	1.64	1.59
2	B	302	ACO	O6A-CCP	3.09	1.54	1.43
2	B	302	ACO	O9P-C9P	3.17	1.29	1.23
2	A	301	ACO	O4B-C1B	3.22	1.45	1.41
2	B	302	ACO	C7P-N8P	3.85	1.55	1.46
2	A	301	ACO	C4A-N3A	3.97	1.41	1.35
2	B	302	ACO	O4B-C4B	4.20	1.54	1.45
2	B	302	ACO	O2B-C2B	4.91	1.54	1.43
2	A	301	ACO	C7P-N8P	4.92	1.57	1.46
2	A	301	ACO	O-C	5.13	1.47	1.21
2	B	302	ACO	O-C	6.93	1.56	1.21

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	ACO	O9P-C9P	7.43	1.38	1.23
2	A	301	ACO	CEP-CBP	8.29	1.72	1.53
2	A	301	ACO	CCP-CBP	11.24	1.67	1.52

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	ACO	CEP-CBP-CCP	-10.97	92.26	108.37
2	B	302	ACO	O-C-CH3	-7.36	90.36	122.88
2	A	301	ACO	O9A-P3B-O7A	-7.04	82.94	110.50
2	A	301	ACO	O9A-P3B-O3B	-5.74	79.88	106.00
2	B	302	ACO	O5B-C5B-C4B	-5.05	91.08	109.00
2	B	302	ACO	C7P-N8P-C9P	-4.82	113.60	122.59
2	B	302	ACO	O2B-C2B-C3B	-4.25	99.09	111.18
2	A	301	ACO	O9A-P3B-O8A	-3.98	91.56	107.61
2	A	301	ACO	O5P-C5P-C6P	-3.85	114.77	122.01
2	B	302	ACO	CDP-CBP-CCP	-3.83	102.74	108.37
2	A	301	ACO	C7P-N8P-C9P	-3.83	115.44	122.59
2	B	302	ACO	O5P-C5P-C6P	-3.73	115.01	122.01
2	A	301	ACO	O-C-CH3	-3.71	106.51	122.88
2	A	301	ACO	O5B-C5B-C4B	-3.67	95.98	109.00
2	B	302	ACO	C2P-C3P-N4P	-3.62	104.50	112.49
2	A	301	ACO	O3B-C3B-C4B	-3.35	97.46	110.04
2	A	301	ACO	O2B-C2B-C3B	-3.14	102.25	111.18
2	A	301	ACO	O6A-CCP-CBP	-3.12	105.54	110.55
2	A	301	ACO	O4B-C4B-C5B	-3.05	99.12	109.40
2	A	301	ACO	C2P-C3P-N4P	-2.60	106.76	112.49
2	B	302	ACO	O6A-P2A-O4A	-2.52	99.07	109.25
2	B	302	ACO	C3P-N4P-C5P	-2.43	118.17	122.84
2	A	301	ACO	O6A-P2A-O4A	-2.39	99.62	109.25
2	B	302	ACO	CEP-CBP-CDP	-2.37	104.14	109.19
2	B	302	ACO	O4B-C4B-C5B	-2.28	101.72	109.40
2	B	302	ACO	O5A-P2A-O6A	-2.20	97.76	108.14
2	A	301	ACO	N3A-C2A-N1A	-2.15	126.99	128.86
2	A	301	ACO	C3P-N4P-C5P	-2.09	118.83	122.84
2	B	302	ACO	C1B-N9A-C4A	2.04	130.15	126.64
2	B	302	ACO	O8A-P3B-O7A	2.07	118.59	110.50
2	B	302	ACO	O2A-P1A-O5B	2.08	117.97	108.14
2	A	301	ACO	O-C-S1P	2.18	133.46	122.65
2	B	302	ACO	O9A-P3B-O8A	2.29	116.87	107.61
2	A	301	ACO	O2A-P1A-O1A	2.41	124.77	112.28
2	B	302	ACO	O9P-C9P-N8P	2.45	127.79	123.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	A	301	ACO	C3B-C2B-C1B	2.49	105.53	99.95
2	B	302	ACO	CH3-C-S1P	2.50	133.80	114.53
2	B	302	ACO	O2A-P1A-O1A	2.52	125.32	112.28
2	B	302	ACO	C3B-C2B-C1B	2.67	105.94	99.95
2	B	302	ACO	O3B-C3B-C2B	2.93	122.47	111.63
2	A	301	ACO	O3B-C3B-C2B	3.16	123.34	111.63
2	B	302	ACO	C7P-C6P-C5P	3.23	117.42	112.22
2	A	301	ACO	C4A-C5A-N7A	3.27	112.57	109.41
2	B	302	ACO	C6P-C5P-N4P	3.37	122.30	116.49
2	B	302	ACO	CEP-CBP-CAP	3.46	114.81	108.82
2	B	302	ACO	C4A-C5A-N7A	3.51	112.81	109.41
2	B	302	ACO	C2P-S1P-C	3.63	122.13	101.67
2	A	301	ACO	O2B-C2B-C1B	3.87	123.72	111.61
2	A	301	ACO	CEP-CBP-CDP	3.93	117.54	109.19
2	A	301	ACO	CDP-CBP-CCP	4.32	114.72	108.37
2	A	301	ACO	O3B-P3B-O7A	4.42	126.58	109.26
2	B	302	ACO	O2B-C2B-C1B	5.11	127.59	111.61
2	A	301	ACO	CEP-CBP-CAP	6.70	120.44	108.82
2	B	302	ACO	C5B-C4B-C3B	8.37	142.67	114.30
2	A	301	ACO	C5B-C4B-C3B	9.24	145.62	114.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	ACO	6	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.