



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:57 pm GMT

PDB ID : 1GJR
Title : FERREDOXIN-NADP⁺ REDUCTASE COMPLEXED WITH NADP⁺ BY COCRYSTALLIZATION
Authors : Hermoso, J.A.; Mayoral, T.; Medina, M.; Sanz-Aparicio, J.; Gomez-Moreno, C.
Deposited on : 2001-08-01
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	<div> <div>6%</div> <div> <div></div> <div>75%</div> <div>20%</div> <div>• •</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2639 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FERREDONIN-NADP REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	S	0	0	0
			2338	1488	399	442	9			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	200	Total	O	0	0
			200	200		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: FERREDOXIN-NADP REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, α , β , γ	151.69Å 151.69Å 35.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.70 – 2.10 29.75 – 2.02	Depositor EDS
% Data completeness (in resolution range)	98.8 (29.70-2.10) 98.5 (29.75-2.02)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 2.01Å)	Xtriage
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.200 , 0.270 0.202 , 0.266	Depositor DCC
R_{free} test set	1638 reflections (6.89%)	DCC
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 67.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.035 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2639	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/2394	0.61	0/3243

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	108	GLU	Peptide
1	A	79	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2338	0	2314	37	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	53	0	31	1	0
3	A	48	0	25	3	0
4	A	200	0	0	4	0
All	All	2639	0	2370	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:257:HIS:HD2	1:A:299:HIS:HE1	1.09	0.94
1:A:257:HIS:CD2	1:A:299:HIS:HE1	2.00	0.77
1:A:257:HIS:HD2	1:A:299:HIS:CE1	2.01	0.72
3:A:305:NAP:O1A	3:A:305:NAP:H51N	1.94	0.67
1:A:243:HIS:CD2	1:A:246:GLN:HE22	2.16	0.64
1:A:25:VAL:HG21	1:A:124:ILE:HG12	1.82	0.61
1:A:15:TYR:HD2	1:A:52:LEU:HD13	1.68	0.58
1:A:190:VAL:O	1:A:236:ILE:HD11	2.03	0.57
1:A:263:LEU:O	1:A:266:MET:HG2	2.04	0.57
1:A:243:HIS:HD2	1:A:246:GLN:HE22	1.53	0.55
1:A:105:LYS:HD2	1:A:110:GLY:O	2.07	0.54
1:A:28:ASN:C	1:A:28:ASN:HD22	2.11	0.53
1:A:236:ILE:HG21	1:A:266:MET:CE	2.39	0.53
1:A:228:ASN:ND2	1:A:232:GLY:H	2.09	0.51
1:A:28:ASN:ND2	1:A:41:GLN:HE21	2.08	0.51
1:A:104:TYR:CD1	1:A:104:TYR:N	2.80	0.50
1:A:271:ASP:O	1:A:275:SER:HB2	2.11	0.50
1:A:169:MET:HE2	1:A:170:PHE:CE2	2.48	0.49
1:A:149:ASN:ND2	4:A:2106:HOH:O	2.46	0.49
1:A:223:SER:HG	3:A:305:NAP:P2B	2.37	0.48
1:A:241:ALA:HA	1:A:273:ALA:HB1	1.97	0.47
1:A:121:LEU:O	1:A:124:ILE:HG22	2.14	0.47
1:A:217:ARG:HD2	4:A:2131:HOH:O	2.14	0.46
1:A:15:TYR:CD2	1:A:52:LEU:HD13	2.49	0.46
1:A:139:GLU:O	1:A:299:HIS:HD2	1.98	0.46
1:A:196:THR:HB	1:A:197:PRO:HD3	1.96	0.46
1:A:199:ILE:CG2	1:A:202:LYS:HB2	2.46	0.46
1:A:149:ASN:HA	1:A:186:PHE:O	2.17	0.45
1:A:45:PHE:HE2	1:A:130:VAL:HG21	1.83	0.43
1:A:236:ILE:HG21	1:A:266:MET:HE2	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:GLN:OE1	3:A:305:NAP:H2A	2.18	0.43
1:A:236:ILE:HG21	1:A:266:MET:HE3	2.00	0.43
1:A:223:SER:O	1:A:233:ARG:HG2	2.18	0.43
1:A:33:LYS:HE2	4:A:2120:HOH:O	2.18	0.43
1:A:168:ARG:HD2	4:A:2039:HOH:O	2.19	0.42
1:A:22:ILE:CD1	1:A:131:LYS:HG2	2.50	0.41
2:A:304:FAD:H1'1	2:A:304:FAD:H9	1.83	0.41
1:A:228:ASN:HD22	1:A:230:GLN:H	1.69	0.40
1:A:56:GLU:HB3	1:A:164:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	293/304 (96%)	280 (96%)	12 (4%)	1 (0%)	44 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	107	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	250/256 (98%)	226 (90%)	24 (10%)	10 6

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	VAL
1	A	24	LYS
1	A	28	ASN
1	A	34	GLU
1	A	70	ASN
1	A	90	VAL
1	A	101	GLN
1	A	104	TYR
1	A	109	SER
1	A	124	ILE
1	A	167	TRP
1	A	172	ASP
1	A	187	SER
1	A	193	VAL
1	A	204	GLU
1	A	214	ASP
1	A	223	SER
1	A	227	LYS
1	A	228	ASN
1	A	230	GLN
1	A	239	ARG
1	A	264	ARG
1	A	284	THR
1	A	290	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	70	ASN
1	A	101	GLN
1	A	106	HIS
1	A	149	ASN
1	A	182	GLN
1	A	226	GLN
1	A	228	ASN
1	A	243	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	246	GLN
1	A	257	HIS
1	A	299	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	304	-	51,58,58	1.73	11 (21%)	54,89,89	1.76	12 (22%)
3	NAP	A	305	-	44,52,52	1.47	10 (22%)	51,80,80	1.68	9 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	304	-	-	0/28/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	305	-	-	0/27/67/67	0/5/5/5

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	304	FAD	P-O2P	-4.05	1.34	1.55
2	A	304	FAD	C2-N1	-3.63	1.31	1.38
2	A	304	FAD	PA-O2A	-3.17	1.39	1.55
3	A	305	NAP	P2B-O2X	-2.75	1.43	1.54
3	A	305	NAP	P2B-O3X	-2.43	1.44	1.54
3	A	305	NAP	C3N-C7N	-2.40	1.46	1.50
3	A	305	NAP	PA-O2A	-2.37	1.43	1.55
2	A	304	FAD	C2B-C1B	-2.30	1.50	1.53
3	A	305	NAP	C5N-C4N	-2.23	1.34	1.38
3	A	305	NAP	C5A-C4A	-2.20	1.35	1.40
2	A	304	FAD	PA-O5B	-2.05	1.50	1.59
3	A	305	NAP	PA-O5B	2.12	1.68	1.59
2	A	304	FAD	C7M-C7	2.51	1.56	1.51
2	A	304	FAD	C4X-C10	2.64	1.45	1.41
3	A	305	NAP	C2D-C1D	2.67	1.57	1.53
2	A	304	FAD	C9A-N10	3.05	1.42	1.38
3	A	305	NAP	P2B-O2B	3.37	1.65	1.59
2	A	304	FAD	O5'-C5'	3.49	1.58	1.44
2	A	304	FAD	O4B-C1B	3.55	1.46	1.41
3	A	305	NAP	C2N-C3N	3.73	1.44	1.39
2	A	304	FAD	C4-N3	3.85	1.40	1.33

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	304	FAD	C4X-C4-N3	-4.25	117.44	123.48
3	A	305	NAP	O7N-C7N-N7N	-3.90	117.04	122.58
3	A	305	NAP	C3N-C2N-N1N	-3.18	117.23	120.43
3	A	305	NAP	C4B-O4B-C1B	-2.94	106.64	109.77
3	A	305	NAP	N3A-C2A-N1A	-2.63	126.57	128.86
2	A	304	FAD	O5'-P-O1P	-2.61	98.70	109.25
2	A	304	FAD	C4-C4X-C10	-2.55	117.90	119.96
2	A	304	FAD	O5B-PA-O1A	-2.20	100.39	109.25
3	A	305	NAP	O3B-C3B-C2B	-2.09	105.24	111.18
2	A	304	FAD	N3A-C2A-N1A	-2.02	127.10	128.86
2	A	304	FAD	O4B-C4B-C3B	-2.01	101.17	105.17
3	A	305	NAP	O5D-PN-O1N	2.15	117.92	109.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	304	FAD	C2A-N1A-C6A	2.24	122.68	118.77
2	A	304	FAD	C4X-N5-C5X	2.40	119.30	116.76
2	A	304	FAD	O3B-C3B-C4B	2.77	119.18	111.09
2	A	304	FAD	C4-C4X-N5	3.03	122.01	118.68
3	A	305	NAP	O2A-PA-O1A	3.37	129.71	112.28
3	A	305	NAP	C3N-C7N-N7N	3.38	121.63	117.77
2	A	304	FAD	C1'-N10-C10	3.47	122.06	118.50
3	A	305	NAP	O3X-P2B-O2X	5.49	129.75	107.61
2	A	304	FAD	C4-N3-C2	5.99	120.39	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	304	FAD	1	0
3	A	305	NAP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	295/304 (97%)	0.18	18 (6%)	22 27	17, 32, 52, 73	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	229	PRO	4.4
1	A	107	PRO	3.8
1	A	69	LYS	3.7
1	A	9	ASP	3.3
1	A	153	LEU	3.0
1	A	70	ASN	3.0
1	A	279	ALA	2.9
1	A	108	GLU	2.6
1	A	228	ASN	2.5
1	A	176	ALA	2.4
1	A	179	PRO	2.4
1	A	276	ALA	2.4
1	A	67	VAL	2.4
1	A	159	ILE	2.4
1	A	278	ALA	2.2
1	A	152	MET	2.1
1	A	189	LEU	2.1
1	A	151	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FAD	A	304	53/53	0.95	0.11	-0.40	12,20,61,64	0
3	NAP	A	305	48/48	0.95	0.11	-0.66	24,37,46,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.