



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 05:28 pm GMT

PDB ID : 2GJM
Title : Crystal structure of Buffalo lactoperoxidase at 2.75Å resolution
Authors : Sheikh, I.A.; Ethayathulla, A.S.; Singh, A.K.; Singh, N.; Sharma, S.; Singh, T.P.
Deposited on : 2006-03-31
Resolution : 2.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3666 (2.80-2.72)
Clashscore	112137	4174 (2.80-2.72)
Ramachandran outliers	110173	4103 (2.80-2.72)
Sidechain outliers	110143	4106 (2.80-2.72)
RSRZ outliers	101464	3697 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	583	<div> <div>5%</div> <div> <div></div> <div>65%</div> <div>30%</div> <div>• •</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	IOD	A	2006	-	-	X	-
8	IOD	A	2007	-	-	-	X

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 5088 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lactoperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	583	Total	C	N	O	S	0	1	0
			4701	2994	835	847	25			

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 3 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 4 is a polymer of unknown type called SUGAR (4-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 5 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 6 is CARBONATE ION (three-letter code: CO3) (formula: CO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	1	3		


- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Ca 1 1	0	0

- Molecule 8 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	8	Total I 8 8	0	0

- Molecule 9 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	183	Total O 183 183	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lactoperoxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.24Å 80.47Å 77.35Å 90.00° 102.71° 90.00°	Depositor
Resolution (Å)	20.00 – 2.75 25.27 – 2.73	Depositor EDS
% Data completeness (in resolution range)	96.0 (20.00-2.75) 95.4 (25.27-2.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.72Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.182 , 0.216 0.184 , 0.216	Depositor DCC
R_{free} test set	798 reflections (5.15%)	DCC
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.711	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 63.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5088	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CO3, SCN, NAG, CA, BMA, HEM, IOD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/4827	0.72	5/6544 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	CYS	C-N-CD	-8.60	101.68	120.60
1	A	355	PRO	CA-N-CD	-7.25	101.35	111.50
1	A	154	PRO	CA-N-CD	-7.25	101.35	111.50
1	A	154	PRO	CA-C-N	-6.28	103.39	117.20
1	A	158	SER	N-CA-C	5.87	126.85	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4701	0	4618	131	0
2	A	56	0	50	0	0
3	A	39	0	34	1	0
4	A	50	0	43	1	0
5	A	3	0	0	0	0
6	A	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	1	0	0	0	0
8	A	8	0	0	5	0
9	A	43	0	30	2	0
10	A	183	0	0	10	0
All	All	5088	0	4775	135	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 135 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:211:LEU:HD23	1:A:254:LEU:HD22	1.44	1.00
1:A:109:GLU:HB3	1:A:112:LYS:HE3	1.48	0.94
1:A:153:CYS:HB2	1:A:154:PRO:HD2	1.47	0.93
1:A:153:CYS:CB	1:A:154:PRO:HD2	1.99	0.93
8:A:2005:IOD:I	10:A:2191:HOH:O	2.61	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	582/583 (100%)	528 (91%)	40 (7%)	14 (2%)	7 20

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	GLU
1	A	107	SER
1	A	108	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	152	VAL
1	A	153	CYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	510/509 (100%)	467 (92%)	43 (8%)	13	32

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	252	LEU
1	A	318	ASN
1	A	526	ASP
1	A	259	ARG
1	A	260	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	349	ASN
1	A	421	ASN
1	A	529	GLN
1	A	318	ASN
1	A	504	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

11 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	581	1,2	14,14,15	0.50	0	15,19,21	1.60	1 (6%)
2	NAG	A	582	2	14,14,15	0.56	0	15,19,21	1.30	1 (6%)
2	NAG	A	583	1,2	14,14,15	0.48	0	15,19,21	0.96	0
2	NAG	A	584	2	14,14,15	0.63	0	15,19,21	1.09	1 (6%)
3	NAG	A	585	1,3	14,14,15	0.46	0	15,19,21	0.84	0
3	NAG	A	586	3	14,14,15	0.64	0	15,19,21	0.96	2 (13%)
3	MAN	A	587	3	11,11,12	0.56	0	13,15,17	0.41	0
4	NAG	A	588	1,4	14,14,15	0.73	0	15,19,21	1.48	1 (6%)
4	NAG	A	589	4	14,14,15	0.71	0	15,19,21	0.86	0
4	BMA	A	590	4	11,11,12	0.98	1 (9%)	13,15,17	2.08	3 (23%)
4	MAN	A	591	4	11,11,12	1.05	0	13,15,17	1.06	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	581	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	582	2	-	0/6/23/26	0/1/1/1
2	NAG	A	583	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	584	2	-	0/6/23/26	0/1/1/1
3	NAG	A	585	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	586	3	-	0/6/23/26	0/1/1/1
3	MAN	A	587	3	-	0/2/19/22	0/1/1/1
4	NAG	A	588	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	589	4	-	0/6/23/26	0/1/1/1
4	BMA	A	590	4	-	0/2/19/22	0/1/1/1
4	MAN	A	591	4	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	590	BMA	O5-C5	2.16	1.48	1.43

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	581	NAG	C2-N2-C7	-4.94	115.74	122.94
2	A	582	NAG	C2-N2-C7	-4.19	116.83	122.94
4	A	591	MAN	C1-C2-C3	-2.59	106.36	109.65
4	A	591	MAN	C1-O5-C5	-2.18	109.16	112.17
3	A	586	NAG	O5-C1-C2	-2.15	108.48	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	586	NAG	1	0
3	A	587	MAN	1	0
4	A	589	NAG	1	0
4	A	590	BMA	1	0

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 9 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	CO3	A	1001	-	0,3,3	0.00	-	0,3,3	0.00	-
5	SCN	A	1502	-	1,2,2	1.57	0	0,1,1	0.00	-
9	HEM	A	605	1	28,50,50	2.06	7 (25%)	17,82,82	4.53	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CO3	A	1001	-	-	0/0/0/0	0/0/0/0
5	SCN	A	1502	-	-	0/0/0/0	0/0/0/0
9	HEM	A	605	1	-	0/6/54/54	0/0/8/8

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	605	HEM	C3B-C2B	-2.52	1.37	1.40
9	A	605	HEM	C1D-ND	2.34	1.41	1.36
9	A	605	HEM	CMC-C2C	2.39	1.56	1.51
9	A	605	HEM	CAA-C2A	2.64	1.56	1.52
9	A	605	HEM	C3C-CAC	4.15	1.56	1.47

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	605	HEM	CBA-CAA-C2A	-3.02	106.72	112.48
9	A	605	HEM	CMD-C2D-C1D	2.28	131.96	128.46
9	A	605	HEM	CMC-C2C-C3C	3.75	131.85	124.89
9	A	605	HEM	CBD-CAD-C3D	9.22	130.05	112.47
9	A	605	HEM	CAD-CBD-CGD	14.79	137.94	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	605	HEM	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	583/583 (100%)	-0.15	31 (5%) 27 21	8, 26, 52, 85	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	578	ARG	7.9
1	A	108	ASN	7.8
1	A	158	SER	6.0
1	A	109	GLU	5.9
1	A	157	GLN	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	583	14/15	0.91	0.20	0.67	45,49,51,53	0
3	NAG	A	585	14/15	0.94	0.17	0.19	39,40,44,45	0
2	NAG	A	581	14/15	0.87	0.17	-0.12	49,51,55,60	0
3	MAN	A	587	11/12	0.61	0.34	-	46,47,48,48	11
4	NAG	A	588	14/15	0.84	0.27	-	45,49,51,51	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	MAN	A	591	11/12	0.59	0.55	-	48,49,49,49	11
3	NAG	A	586	14/15	0.82	0.35	-	47,50,51,51	0
2	NAG	A	582	14/15	0.86	0.43	-	63,66,67,67	0
4	BMA	A	590	11/12	0.66	0.49	-	49,49,49,49	11
4	NAG	A	589	14/15	0.67	0.44	-	48,49,50,50	14
2	NAG	A	584	14/15	0.80	0.48	-	56,57,58,58	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
8	IOD	A	2007	1/1	0.97	0.13	2.12	10,10,10,10	1
9	HEM	A	605	43/43	0.95	0.14	0.01	13,19,23,25	0
7	CA	A	1503	1/1	0.99	0.10	-0.95	14,14,14,14	0
8	IOD	A	2004	1/1	0.97	0.12	-1.03	33,33,33,33	1
5	SCN	A	1502	3/3	0.94	0.12	-1.79	14,14,14,15	0
8	IOD	A	2001	1/1	0.97	0.06	-2.18	47,47,47,47	1
8	IOD	A	2006	1/1	0.93	0.12	-2.30	37,37,37,37	1
8	IOD	A	2005	1/1	0.92	0.10	-3.55	37,37,37,37	1
8	IOD	A	2002	1/1	0.99	0.05	-4.94	42,42,42,42	1
8	IOD	A	2008	1/1	0.92	0.12	-	24,24,24,24	1
6	CO3	A	1001	4/4	0.95	0.20	-	31,32,33,33	0
8	IOD	A	2003	1/1	0.97	0.07	-	47,47,47,47	1

6.5 Other polymers [i](#)

There are no such residues in this entry.