



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 07:36 pm GMT

PDB ID : 3GJ7
Title : Crystal structure of human RanGDP-Nup153ZnF12 complex
Authors : Partridge, J.R.; Schwartz, T.U.
Deposited on : 2009-03-07
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3233 (1.96-1.92)
Clashscore	112137	3430 (1.96-1.92)
Ramachandran outliers	110173	3395 (1.96-1.92)
Sidechain outliers	110143	3395 (1.96-1.92)
RSRZ outliers	101464	3250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	
1	C	221	
2	B	98	
2	D	98	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	306	-	-	-	X
3	MG	C	303	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	201	Total 1609	C 1039	N 279	O 285	S 6	0	0	0
1	C	202	Total 1614	C 1042	N 280	O 286	S 6	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP P62826
A	-3	PRO	-	EXPRESSION TAG	UNP P62826
A	-2	HIS	-	EXPRESSION TAG	UNP P62826
A	-1	MET	-	EXPRESSION TAG	UNP P62826
A	0	ALA	-	EXPRESSION TAG	UNP P62826
A	1	SER	-	EXPRESSION TAG	UNP P62826
A	35	SER	PHE	ENGINEERED	UNP P62826
C	-4	GLY	-	EXPRESSION TAG	UNP P62826
C	-3	PRO	-	EXPRESSION TAG	UNP P62826
C	-2	HIS	-	EXPRESSION TAG	UNP P62826
C	-1	MET	-	EXPRESSION TAG	UNP P62826
C	0	ALA	-	EXPRESSION TAG	UNP P62826
C	1	SER	-	EXPRESSION TAG	UNP P62826
C	35	SER	PHE	ENGINEERED	UNP P62826

- Molecule 2 is a protein called Nuclear pore complex protein Nup153.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	27	Total 201	C 124	N 33	O 40	S 4	0	0	0
2	D	27	Total 201	C 124	N 33	O 40	S 4	0	0	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	653	GLY	-	EXPRESSION TAG	UNP P49791
B	654	PRO	-	EXPRESSION TAG	UNP P49791
B	655	LEU	-	EXPRESSION TAG	UNP P49791
B	656	GLY	-	EXPRESSION TAG	UNP P49791
B	657	SER	-	EXPRESSION TAG	UNP P49791
D	653	GLY	-	EXPRESSION TAG	UNP P49791
D	654	PRO	-	EXPRESSION TAG	UNP P49791
D	655	LEU	-	EXPRESSION TAG	UNP P49791
D	656	GLY	-	EXPRESSION TAG	UNP P49791
D	657	SER	-	EXPRESSION TAG	UNP P49791

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Mg 3 3	0	0
3	C	3	Total Mg 3 3	0	0

- Molecule 4 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O P 28 10 5 11 2	0	0
4	C	1	Total C N O P 28 10 5 11 2	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total 1	Zn 1	0	0
5	D	1	Total 1	Zn 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	107	Total 107	O 107	0	0
6	B	4	Total 4	O 4	0	0
6	C	169	Total 169	O 169	0	0
6	D	14	Total 14	O 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GTP-binding nuclear protein Ran




- Molecule 1: GTP-binding nuclear protein Ran



- Molecule 2: Nuclear pore complex protein Nup153



- Molecule 2: Nuclear pore complex protein Nup153



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.52Å 61.13Å 80.22Å 90.00° 93.71° 90.00°	Depositor
Resolution (Å)	29.75 – 1.93 29.75 – 1.93	Depositor EDS
% Data completeness (in resolution range)	91.9 (29.75-1.93) 91.9 (29.75-1.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.93Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.186 , 0.229 0.179 , 0.227	Depositor DCC
R_{free} test set	1992 reflections (5.08%)	DCC
Wilson B-factor (Å ²)	37.6	Xtrriage
Anisotropy	0.363	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 52.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3983	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1649	0.49	0/2235
1	C	0.39	0/1654	0.54	1/2242 (0.0%)
2	B	0.24	0/205	0.45	0/280
2	D	0.31	0/205	0.45	0/280
All	All	0.36	0/3713	0.51	1/5037 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	50	LEU	CA-CB-CG	5.55	128.06	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1609	0	1626	21	0
1	C	1614	0	1631	9	0
2	B	201	0	191	12	0
2	D	201	0	191	6	0
3	A	3	0	0	0	0
3	C	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	28	0	12	0	0
4	C	28	0	12	0	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
6	A	107	0	0	3	0
6	B	4	0	0	0	0
6	C	169	0	0	1	0
6	D	14	0	0	0	0
All	All	3983	0	3663	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:60:LYS:HE2	1:C:62:ASN:OD1	1.70	0.92
1:A:60:LYS:HE3	1:A:62:ASN:OD1	1.84	0.77
1:A:81:ILE:HA	6:A:297:HOH:O	1.98	0.63
2:B:734:ASN:HD21	2:B:742:VAL:HG12	1.68	0.59
2:B:734:ASN:HD21	2:B:742:VAL:H	1.51	0.58
2:D:725:TRP:NE1	2:D:748:LYS:HE2	2.18	0.57
2:B:724:THR:HG22	2:B:735:LYS:HA	1.89	0.54
2:D:723:GLY:C	2:D:736:PRO:HG3	2.29	0.53
2:D:725:TRP:CE2	2:D:748:LYS:HE2	2.44	0.53
2:B:734:ASN:ND2	2:B:742:VAL:HG12	2.24	0.53
2:D:735:LYS:HB3	2:D:737:GLU:HG2	1.90	0.53
1:A:111:VAL:CG1	6:A:297:HOH:O	2.57	0.52
2:B:723:GLY:O	2:B:736:PRO:HB3	2.11	0.51
1:A:24:THR:CG2	1:A:28:LYS:HE2	2.40	0.51
1:A:86:ALA:CB	1:A:108:LEU:HD21	2.41	0.51
1:A:81:ILE:HG21	2:B:743:ALA:HA	1.91	0.50
2:B:725:TRP:CE2	2:B:748:LYS:HE2	2.46	0.50
1:C:49:PRO:HB3	1:C:60:LYS:HE3	1.94	0.49
1:A:80:TYR:HB2	1:A:111:VAL:HG11	1.94	0.49
2:B:724:THR:HB	2:B:734:ASN:O	2.13	0.49
2:B:735:LYS:HE2	2:B:737:GLU:OE2	2.13	0.49
2:B:724:THR:HA	2:B:736:PRO:HD3	1.97	0.47
1:A:111:VAL:HG13	6:A:297:HOH:O	2.14	0.47
1:A:139:HIS:HA	1:A:144:LEU:HB2	1.97	0.47
1:A:112:CYS:O	1:A:113:GLU:HB2	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:HIS:HB2	1:A:144:LEU:O	2.16	0.45
1:C:123:LYS:HB3	1:C:126:ILE:HD12	1.98	0.45
1:C:159:LYS:HB2	1:C:160:PRO:HD3	1.97	0.45
2:B:734:ASN:ND2	2:B:742:VAL:H	2.12	0.45
1:A:85:CYS:HB2	1:A:164:LEU:HD22	1.99	0.44
1:A:153:SER:O	1:A:154:ASN:HB2	2.18	0.43
1:A:12:LYS:HE3	1:A:64:TRP:CE2	2.53	0.43
1:A:124:VAL:HG22	1:A:149:ILE:O	2.18	0.43
1:A:159:LYS:HB2	1:A:160:PRO:HD3	2.00	0.43
1:A:105:HIS:CE1	1:A:109:VAL:HG21	2.54	0.42
2:D:744:CYS:O	2:D:745:GLU:HB2	2.19	0.42
1:C:153:SER:HA	1:C:185:PRO:HB3	2.02	0.42
1:A:199:HIS:O	1:A:203:VAL:HG23	2.20	0.41
1:A:80:TYR:CB	1:A:111:VAL:HG11	2.51	0.41
1:C:105:HIS:CE1	1:C:142:LYS:HE2	2.55	0.41
1:A:117:ILE:HB	1:A:144:LEU:HD23	2.03	0.41
1:C:95:ARG:HG2	1:C:99:LYS:NZ	2.36	0.40
2:D:737:GLU:HG2	2:D:737:GLU:H	1.76	0.40
1:A:142:LYS:O	1:A:144:LEU:HG	2.22	0.40
1:C:103:ASN:ND2	6:C:364:HOH:O	2.54	0.40
1:C:202:GLU:OE1	1:C:202:GLU:HA	2.21	0.40
2:B:748:LYS:HA	2:B:749:PRO:HD3	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	199/221 (90%)	192 (96%)	6 (3%)	1 (0%)	32 19
1	C	200/221 (90%)	196 (98%)	4 (2%)	0	100 100
2	B	25/98 (26%)	24 (96%)	1 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	25/98 (26%)	23 (92%)	2 (8%)	0	100	100
All	All	449/638 (70%)	435 (97%)	13 (3%)	1 (0%)	51	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/188 (93%)	173 (99%)	1 (1%)	89	87
1	C	174/188 (93%)	173 (99%)	1 (1%)	89	87
2	B	24/79 (30%)	23 (96%)	1 (4%)	34	19
2	D	24/79 (30%)	24 (100%)	0	100	100
All	All	396/534 (74%)	393 (99%)	3 (1%)	85	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	LYS
2	B	737	GLU
1	C	137	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	HIS
1	A	100	ASN
1	A	103	ASN
1	A	105	HIS
2	B	734	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	48	HIS
1	C	100	ASN
1	C	103	ASN
1	C	105	HIS
1	C	139	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 8 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GDP	A	302	3	25,30,30	1.12	2 (8%)	26,47,47	1.89	6 (23%)
4	GDP	C	302	3	25,30,30	1.07	2 (8%)	26,47,47	1.96	6 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GDP	A	302	3	-	0/12/32/32	0/3/3/3
4	GDP	C	302	3	-	0/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	302	GDP	C5-C4	2.84	1.46	1.40
4	A	302	GDP	C5-C4	3.20	1.47	1.40
4	C	302	GDP	C6-C5	3.41	1.47	1.41
4	A	302	GDP	C6-C5	3.61	1.48	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	302	GDP	C6-C5-C4	-4.28	116.59	120.84
4	A	302	GDP	C5-C6-N1	-3.92	117.90	123.48
4	C	302	GDP	C5-C6-N1	-3.88	117.96	123.48
4	A	302	GDP	C6-C5-C4	-3.57	117.29	120.84
4	C	302	GDP	N3-C2-N1	-3.22	122.75	127.46
4	A	302	GDP	N3-C2-N1	-2.93	123.19	127.46
4	A	302	GDP	C4-C5-N7	-2.57	106.93	109.41
4	C	302	GDP	C4-C5-N7	-2.28	107.21	109.41
4	C	302	GDP	C2-N3-C4	4.22	120.08	115.16
4	A	302	GDP	C2-N3-C4	4.48	120.39	115.16
4	A	302	GDP	C6-N1-C2	4.48	122.50	116.06
4	C	302	GDP	C6-N1-C2	4.67	122.77	116.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/221 (90%)	0.30	15 (7%) 15 21	31, 50, 100, 124	0
1	C	202/221 (91%)	0.12	11 (5%) 26 34	27, 40, 72, 127	0
2	B	27/98 (27%)	2.32	11 (40%) 0 0	55, 101, 148, 163	0
2	D	27/98 (27%)	0.73	3 (11%) 6 9	36, 63, 96, 114	0
All	All	457/638 (71%)	0.37	40 (8%) 11 16	27, 47, 107, 163	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	723	GLY	9.9
2	B	739	VAL	6.7
2	B	749	PRO	6.3
2	B	738	ALA	5.7
1	A	207	THR	5.1
2	B	736	PRO	5.0
2	B	725	TRP	4.9
2	B	747	PRO	4.3
1	C	7	PRO	3.7
1	A	87	ILE	3.6
1	A	143	ASN	3.5
1	A	8	GLN	3.5
1	C	207	THR	3.4
2	D	749	PRO	3.4
2	B	724	THR	3.4
2	B	748	LYS	3.3
1	C	87	ILE	3.2
1	A	141	LYS	3.2
1	C	141	LYS	3.2
1	A	7	PRO	3.1
2	D	723	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	733	GLN	2.9
1	A	205	GLN	2.7
1	C	114	ASN	2.5
1	C	149	ILE	2.5
2	B	735	LYS	2.5
1	A	149	ILE	2.5
1	A	134	LYS	2.4
1	A	14	VAL	2.4
1	C	208	ALA	2.4
1	A	199	HIS	2.3
1	C	13	LEU	2.3
1	A	16	VAL	2.2
2	D	738	ALA	2.2
1	C	63	VAL	2.2
1	C	15	LEU	2.2
1	A	142	LYS	2.2
1	A	13	LEU	2.1
1	C	16	VAL	2.1
1	A	63	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MG	C	303	1/1	0.95	0.18	6.99	57,57,57,57	0
3	MG	A	306	1/1	1.00	0.16	4.75	43,43,43,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MG	C	305	1/1	0.98	0.12	0.31	40,40,40,40	0
5	ZN	B	300	1/1	1.00	0.09	-0.54	67,67,67,67	0
5	ZN	D	300	1/1	1.00	0.07	-0.87	40,40,40,40	0
4	GDP	A	302	28/28	0.98	0.08	-1.06	33,42,51,55	0
4	GDP	C	302	28/28	0.99	0.08	-1.37	27,35,40,49	0
3	MG	A	304	1/1	0.98	0.11	-	48,48,48,48	0
3	MG	A	301	1/1	0.98	0.16	-	44,44,44,44	0
3	MG	C	301	1/1	0.99	0.12	-	34,34,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.