



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 04:05 am GMT

PDB ID : 1GL7
Title : PLASMID COUPLING PROTEIN TRWB IN COMPLEX WITH THE NON-HYDROLISABLE ATP-ANALOGUE ADPNP.
Authors : Gomis-Ruth, F.X.; Moncalian, G.; De La Cruz, F.; Coll, M.
Deposited on : 2001-08-28
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2037 (3.00-3.00)
Ramachandran outliers	110173	1973 (3.00-3.00)
Sidechain outliers	110143	1976 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	437	
1	B	437	
1	D	437	
1	E	437	
1	F	437	
1	G	437	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 19892 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CONJUGAL TRANSFER PROTEIN TRWB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	0	0
			3233	2046	577	600	10			
1	B	417	Total	C	N	O	S	0	0	0
			3276	2072	587	607	10			
1	D	417	Total	C	N	O	S	0	0	0
			3273	2069	587	607	10			
1	E	414	Total	C	N	O	S	0	0	0
			3254	2059	581	604	10			
1	F	413	Total	C	N	O	S	0	0	0
			3240	2050	579	601	10			
1	G	412	Total	C	N	O	S	0	0	0
			3237	2048	578	601	10			

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	E	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	F	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	G	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		
4	B	29	Total	O	0	0
			29	29		
4	D	39	Total	O	0	0
			39	39		
4	E	53	Total	O	0	0
			53	53		
4	F	23	Total	O	0	0
			23	23		
4	G	18	Total	O	0	0
			18	18		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CONJUGAL TRANSFER PROTEIN TRWB




• Molecule 1: CONJUGAL TRANSFER PROTEIN TRWB






- Molecule 1: CONJUGAL TRANSFER PROTEIN TRWB




- Molecule 1: CONJUGAL TRANSFER PROTEIN TRWB




- Molecule 1: CONJUGAL TRANSFER PROTEIN TRWB





• Molecule 1: CONJUGAL TRANSFER PROTEIN TRWB

Chain G: 51% 39% 6%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	152.70Å 152.70Å 252.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 3.00	Depositor
% Data completeness (in resolution range)	100.0 (50.00-3.00)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.219 , 0.265	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	19892	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3297	0.68	0/4457
1	B	0.46	0/3340	0.68	1/4514 (0.0%)
1	D	0.47	0/3337	0.69	1/4509 (0.0%)
1	E	0.47	0/3318	0.70	0/4485
1	F	0.46	0/3304	0.69	0/4466
1	G	0.47	0/3301	0.68	0/4462
All	All	0.47	0/19897	0.69	2/26893 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	157	VAL	N-CA-C	-5.51	96.12	111.00
1	D	157	VAL	N-CA-C	-5.14	97.11	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3233	0	3261	169	0
1	B	3276	0	3308	165	1
1	D	3273	0	3302	157	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	3254	0	3283	185	0
1	F	3240	0	3270	172	0
1	G	3237	0	3264	180	0
2	A	31	0	13	1	0
2	B	31	0	13	2	0
2	D	31	0	13	1	0
2	E	31	0	13	2	0
2	F	31	0	13	2	0
2	G	31	0	13	1	0
3	A	1	0	0	0	0
4	A	30	0	0	1	0
4	B	29	0	0	1	0
4	D	39	0	0	3	0
4	E	53	0	0	3	0
4	F	23	0	0	0	0
4	G	18	0	0	0	0
All	All	19892	0	19766	989	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 989 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:LEU:HD13	1:B:403:LEU:HD11	1.36	1.05
1:D:413:LEU:HD23	1:D:475:ALA:HB2	1.37	1.03
1:F:463:MET:HB2	1:F:466:GLU:HG3	1.43	1.01
1:B:503:VAL:O	1:B:504:GLU:HB2	1.63	0.98
1:G:154:MET:HE3	1:G:156:ILE:HD11	1.44	0.98

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:LYS:NZ	1:D:241:GLU:OE1[4_456]	2.08	0.12

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/437 (93%)	367 (90%)	36 (9%)	4 (1%)	18	59
1	B	413/437 (94%)	369 (89%)	39 (9%)	5 (1%)	15	53
1	D	413/437 (94%)	373 (90%)	36 (9%)	4 (1%)	18	59
1	E	410/437 (94%)	373 (91%)	34 (8%)	3 (1%)	25	67
1	F	409/437 (94%)	371 (91%)	32 (8%)	6 (2%)	12	48
1	G	408/437 (93%)	369 (90%)	35 (9%)	4 (1%)	18	59
All	All	2460/2622 (94%)	2222 (90%)	212 (9%)	26 (1%)	17	56

5 of 26 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	289	PRO
1	F	293	THR
1	B	125	HIS
1	B	189	ASN
1	D	289	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/364 (94%)	315 (92%)	29 (8%)	13	43
1	B	348/364 (96%)	316 (91%)	32 (9%)	11	38
1	D	347/364 (95%)	314 (90%)	33 (10%)	10	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	346/364 (95%)	315 (91%)	31 (9%)	11	40
1	F	344/364 (94%)	313 (91%)	31 (9%)	11	40
1	G	344/364 (94%)	319 (93%)	25 (7%)	16	50
All	All	2073/2184 (95%)	1892 (91%)	181 (9%)	12	41

5 of 181 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	353	LEU
1	E	240	ARG
1	G	289	PRO
1	D	372	THR
1	D	482	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	494	GLN
1	E	180	GLN
1	G	423	ASN
1	D	499	GLN
1	E	423	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ANP	A	701	-	29,33,33	2.06	10 (34%)	28,52,52	2.22	7 (25%)
2	ANP	B	701	-	29,33,33	2.06	10 (34%)	28,52,52	2.24	6 (21%)
2	ANP	D	701	-	29,33,33	2.10	11 (37%)	28,52,52	2.23	9 (32%)
2	ANP	E	701	-	29,33,33	2.05	11 (37%)	28,52,52	2.25	7 (25%)
2	ANP	F	701	-	29,33,33	1.99	10 (34%)	28,52,52	2.27	8 (28%)
2	ANP	G	701	-	29,33,33	1.98	10 (34%)	28,52,52	2.25	7 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	701	-	-	0/13/38/38	0/3/3/3
2	ANP	B	701	-	-	0/13/38/38	0/3/3/3
2	ANP	D	701	-	-	0/13/38/38	0/3/3/3
2	ANP	E	701	-	-	0/13/38/38	0/3/3/3
2	ANP	F	701	-	-	0/13/38/38	0/3/3/3
2	ANP	G	701	-	-	0/13/38/38	0/3/3/3

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	701	ANP	PG-O3G	-2.97	1.48	1.56
2	E	701	ANP	PG-O3G	-2.95	1.48	1.56
2	E	701	ANP	O4'-C4'	-2.91	1.38	1.45
2	A	701	ANP	O4'-C4'	-2.86	1.38	1.45
2	A	701	ANP	PG-O3G	-2.81	1.49	1.56

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	ANP	O1G-PG-N3B	-7.07	101.22	111.79
2	F	701	ANP	O1G-PG-N3B	-6.92	101.45	111.79
2	D	701	ANP	O1G-PG-N3B	-6.85	101.54	111.79
2	E	701	ANP	O1G-PG-N3B	-6.83	101.58	111.79
2	G	701	ANP	O1G-PG-N3B	-6.78	101.64	111.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	ANP	1	0
2	B	701	ANP	2	0
2	D	701	ANP	1	0
2	E	701	ANP	2	0
2	F	701	ANP	2	0
2	G	701	ANP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.