



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 10:59 pm GMT

PDB ID : 4GPD  
Title : THE STRUCTURE OF LOBSTER APO-D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE AT 3.0 ANGSTROMS RESOLUTION  
Authors : Griffith, J.P.; Song, S.; Rossmann, M.G.  
Deposited on : 1988-01-04  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	333	<div> <div></div> <div>9% 27% 39% 25%</div> </div>
1	2	333	<div> <div></div> <div>6% 33% 38% 23%</div> </div>
1	3	333	<div> <div></div> <div>6% 28% 39% 28%</div> </div>
1	4	333	<div> <div></div> <div>7% 30% 37% 26%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10150 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APO-D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1	333	Total	C	N	O	S	0	0	0
			2507	1591	419	482	15			
1	2	333	Total	C	N	O	S	0	0	0
			2507	1591	419	482	15			
1	3	333	Total	C	N	O	S	0	0	0
			2507	1591	419	482	15			
1	4	333	Total	C	N	O	S	0	0	0
			2507	1591	419	482	15			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	1	23	Total	O	0	0
			23	23		
2	2	28	Total	O	0	0
			28	28		
2	3	27	Total	O	0	0
			27	27		
2	4	44	Total	O	0	0
			44	44		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: APO-D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE




#### • Molecule 1: APO-D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE






• Molecule 1: APO-D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE



• Molecule 1: APO-D-GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.02Å 80.96Å 82.55Å 110.85° 71.47° 116.86°	Depositor
Resolution (Å)	6.00 – 2.80 45.95 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-2.80) 57.9 (45.95-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.54 (at 2.51Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.218 , (Not available) 0.230 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.025 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	10150	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	1	1.55	19/2553 (0.7%)	2.75	238/3451 (6.9%)
1	2	1.61	27/2553 (1.1%)	2.88	223/3451 (6.5%)
1	3	1.63	29/2553 (1.1%)	2.93	257/3451 (7.4%)
1	4	1.61	26/2553 (1.0%)	3.05	259/3451 (7.5%)
All	All	1.60	101/10212 (1.0%)	2.90	977/13804 (7.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	3	0	2

The worst 5 of 101 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	2	296	GLY	N-CA	10.32	1.61	1.46
1	2	205	SER	CB-OG	8.01	1.52	1.42
1	3	50	GLY	N-CA	-7.68	1.34	1.46
1	4	50	GLY	N-CA	-7.47	1.34	1.46
1	1	300	SER	CA-CB	-7.33	1.42	1.52

The worst 5 of 977 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	4	227	MET	CA-CB-CG	39.23	179.99	113.30
1	2	287	ARG	CD-NE-CZ	33.00	169.80	123.60
1	2	10	ARG	NE-CZ-NH1	32.49	136.55	120.30
1	2	10	ARG	NE-CZ-NH2	-30.94	104.83	120.30
1	4	244	ARG	CD-NE-CZ	30.59	166.43	123.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	3	13	ARG	Sidechain
1	3	196	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2507	0	2519	903	2
1	2	2507	0	2523	875	25
1	3	2507	0	2519	901	13
1	4	2507	0	2520	887	13
2	1	23	0	0	12	0
2	2	28	0	0	26	4
2	3	27	0	0	28	2
2	4	44	0	0	32	3
All	All	10150	0	10081	3343	31

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 166.

The worst 5 of 3343 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:4:32:PRO:HA	1:4:74:ASN:ND2	1.32	1.40
1:2:277:VAL:HG21	1:4:41:TYR:CE1	1.58	1.39
1:2:32:PRO:HA	1:2:74:ASN:ND2	1.38	1.34
1:1:220:LEU:HA	1:1:223:LYS:NZ	1.43	1.32
1:1:126:MET:HB2	1:1:215:LYS:NZ	1.45	1.30

The worst 5 of 31 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:2:108:PHE:CE2	1:4:138:LYS:O[1_565]	0.55	1.65

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:2:102:GLU:CB	2:4:351:HOH:O[1_565]	0.85	1.35
1:2:123:ASP:CB	1:4:108:PHE:CE2[1_565]	0.92	1.28
1:2:80:ASN:OD1	1:3:139:ASP:OD2[1_665]	0.98	1.22
1:2:108:PHE:CZ	1:4:138:LYS:O[1_565]	1.03	1.17

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	331/333 (99%)	177 (54%)	70 (21%)	84 (25%)	0	0
1	2	331/333 (99%)	180 (54%)	75 (23%)	76 (23%)	0	0
1	3	331/333 (99%)	175 (53%)	70 (21%)	86 (26%)	0	0
1	4	331/333 (99%)	180 (54%)	73 (22%)	78 (24%)	0	0
All	All	1324/1332 (99%)	712 (54%)	288 (22%)	324 (24%)	0	0

5 of 324 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	1	8	PHE
1	1	10	ARG
1	1	27	VAL
1	1	50	GLY
1	1	51	VAL

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	271/271 (100%)	169 (62%)	102 (38%)	0	0
1	2	271/271 (100%)	182 (67%)	89 (33%)	0	0
1	3	271/271 (100%)	171 (63%)	100 (37%)	0	0
1	4	271/271 (100%)	178 (66%)	93 (34%)	0	0
All	All	1084/1084 (100%)	700 (65%)	384 (35%)	0	0

5 of 384 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	2	289	SER
1	3	80	ASN
1	4	232	PRO
1	2	299	LEU
1	3	16	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	2	201	ASN
1	3	25	GLN
1	4	151	ASN
1	3	6	ASN
1	3	30	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	1	333/333 (100%)	-0.72	1 (0%) 93 92	2, 9, 28, 32	0
1	2	333/333 (100%)	-0.75	1 (0%) 93 92	2, 9, 28, 32	0
1	3	333/333 (100%)	-0.67	3 (0%) 84 79	2, 9, 28, 32	0
1	4	333/333 (100%)	-0.70	1 (0%) 93 92	2, 9, 28, 32	0
All	All	1332/1332 (100%)	-0.71	6 (0%) 90 88	2, 9, 28, 32	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	3	106	ALA	3.0
1	1	110	GLY	2.8
1	2	212	ALA	2.6
1	4	110	GLY	2.5
1	3	132	ASN	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.