



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2017 – 11:50 AM EDT

PDB ID : 2GT4
Title : Crystal Structure of the Y103F mutant of the GDP-mannose mannosyl hydrolase in complex with GDP-mannose and MG+2
Authors : Gabelli, S.B.; Bianchet, M.A.; Azurmendi, H.F.; Mildvan, A.S.; Amzel, L.A.
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	160	
1	B	160	
1	C	160	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BMA	C	202	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GDP-mannose mannosyl hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	160	Total	C	N	O	S	0	0	0
			1302	834	224	241	3			
1	B	160	Total	C	N	O	S	0	0	0
			1302	834	224	241	3			
1	C	160	Total	C	N	O	S	0	0	0
			1302	834	224	241	3			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP P32056
A	103	PHE	TYR	ENGINEERED	UNP P32056
B	1	MET	-	INITIATING METHIONINE	UNP P32056
B	103	PHE	TYR	ENGINEERED	UNP P32056
C	1	MET	-	INITIATING METHIONINE	UNP P32056
C	103	PHE	TYR	ENGINEERED	UNP P32056

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE-ALPHA-D-MANNOSE (three-letter code: GDD) (formula: C₁₆H₂₅N₅O₁₆P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			39	16	5	16	2		
3	B	1	Total	C	N	O	P	0	0
			39	16	5	16	2		
3	C	1	Total	C	N	O	P	0	0
			39	16	5	16	2		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	98	Total	O	0	0
			98	98		
5	B	116	Total	O	0	0
			116	116		
5	C	131	Total	O	0	0
			131	131		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: GDP-mannose mannosyl hydrolase

Chain A:  86% 12% .





- Molecule 1: GDP-mannose mannosyl hydrolase

Chain B:  90% 8% .



- Molecule 1: GDP-mannose mannosyl hydrolase

Chain C:  82% 16% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	137.78Å 93.89Å 66.10Å 90.00° 91.23° 90.00°	Depositor
Resolution (Å)	77.61 – 2.30	Depositor
% Data completeness (in resolution range)	97.5 (77.61-2.30)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.186 , 0.226	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4393	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BMA, GDD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/1336	0.60	1/1811 (0.1%)
1	B	0.52	0/1336	0.63	1/1811 (0.1%)
1	C	0.54	0/1336	0.62	0/1811
All	All	0.51	0/4008	0.62	2/5433 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	82	PHE	CB-CA-C	5.68	121.76	110.40
1	A	135	LEU	CA-CB-CG	5.45	127.84	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1302	0	1249	12	0
1	B	1302	0	1247	25	0
1	C	1302	0	1247	41	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	39	0	23	0	0
3	B	39	0	23	3	0
3	C	39	0	23	4	0
4	B	11	0	10	4	0
4	C	11	0	10	8	0
5	A	98	0	0	3	0
5	B	116	0	0	3	0
5	C	131	0	0	6	0
All	All	4393	0	3832	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:36:LYS:HD2	1:C:118:LEU:CD2	1.65	1.26
1:B:36:LYS:HD2	1:C:118:LEU:HD23	1.26	1.09
1:C:37:ARG:HH22	1:C:41:PRO:HD2	1.21	1.04
1:A:1:MET:N	5:A:444:HOH:O	1.93	1.01
1:B:36:LYS:CD	1:C:118:LEU:HD23	1.92	1.00
1:C:37:ARG:NH1	1:C:41:PRO:O	1.95	0.98
1:C:37:ARG:HH11	1:C:37:ARG:HG2	1.31	0.95
1:B:36:LYS:HD2	1:C:118:LEU:HD21	1.45	0.94
1:B:36:LYS:CD	1:C:118:LEU:CD2	2.45	0.94
1:C:58:THR:HG22	1:C:61:ALA:H	1.32	0.92
5:B:425:HOH:O	1:C:1:MET:SD	2.28	0.91
1:C:111:ARG:HH11	1:C:111:ARG:HG2	1.39	0.85
1:B:37:ARG:CZ	4:B:202:BMA:H61	2.08	0.84
1:C:37:ARG:HH11	1:C:37:ARG:CG	1.97	0.77
1:C:37:ARG:HH22	1:C:41:PRO:CD	1.99	0.76
1:B:154:ARG:HG3	1:B:160:LEU:HD13	1.66	0.76
4:B:202:BMA:H62	3:B:201:GDD:O3B	1.88	0.74
1:B:37:ARG:NH2	3:B:201:GDD:O6A	2.18	0.73
1:C:37:ARG:HD2	4:C:202:BMA:O2	1.91	0.70
1:B:37:ARG:NH2	4:B:202:BMA:H61	2.07	0.69
1:B:37:ARG:HH12	1:B:40:ARG:C	1.98	0.67
1:B:37:ARG:HH22	1:B:41:PRO:HD2	1.60	0.66
1:C:58:THR:HG21	5:C:493:HOH:O	1.94	0.66
1:B:36:LYS:CE	1:C:118:LEU:CD2	2.76	0.64
4:C:202:BMA:O4	5:C:413:HOH:O	2.16	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:115:GLU:H	1:C:115:GLU:CD	2.04	0.61
1:C:37:ARG:HA	4:C:202:BMA:O2	2.00	0.61
1:A:125:ASP:HB2	5:A:439:HOH:O	2.01	0.61
4:C:202:BMA:H62	3:C:201:GDD:O3B	2.02	0.60
4:B:202:BMA:O4	3:B:201:GDD:O2B	2.19	0.59
1:B:93:ASN:ND2	1:B:95:SER:H	1.99	0.59
1:A:29:ARG:NH2	5:A:473:HOH:O	2.36	0.58
1:C:37:ARG:CG	1:C:37:ARG:NH1	2.59	0.58
1:C:38:THR:N	4:C:202:BMA:O2	2.34	0.58
1:B:37:ARG:NH2	5:B:416:HOH:O	2.37	0.57
1:C:111:ARG:HH11	1:C:111:ARG:CG	2.13	0.57
4:C:202:BMA:H62	3:C:201:GDD:PB	2.45	0.56
1:A:2:MET:H	1:A:54:GLN:HE21	1.53	0.54
1:B:37:ARG:HH12	1:B:41:PRO:N	2.05	0.54
1:C:14:ARG:HD3	5:C:472:HOH:O	2.08	0.53
1:C:37:ARG:NH2	4:C:202:BMA:O6	2.42	0.53
4:C:202:BMA:C6	3:C:201:GDD:O3B	2.58	0.52
1:A:16:THR:OG1	1:A:17:PRO:HD2	2.10	0.51
1:C:60:GLU:HG3	1:C:77:ILE:HG21	1.93	0.51
1:A:92:ASP:N	1:A:92:ASP:OD1	2.44	0.50
1:A:88:HIS:HB2	1:A:103:PHE:HB2	1.95	0.49
1:A:26:GLU:HB3	1:A:111:ARG:HG2	1.95	0.48
1:A:64:GLU:OE1	1:A:74:ARG:NH1	2.46	0.48
1:A:1:MET:HG3	1:A:57:GLU:HB2	1.95	0.47
1:A:93:ASN:ND2	1:A:95:SER:H	2.12	0.47
1:C:25:VAL:O	1:C:32:PHE:HA	2.15	0.47
1:B:37:ARG:NH1	1:B:40:ARG:C	2.66	0.47
3:C:201:GDD:H5'1	5:C:413:HOH:O	2.14	0.47
1:C:37:ARG:NH2	1:C:41:PRO:HD2	2.06	0.46
1:B:37:ARG:HD3	1:B:37:ARG:HA	1.81	0.46
1:C:2:MET:H	1:C:54:GLN:HE21	1.64	0.45
1:A:140:ASN:HD22	1:A:140:ASN:HA	1.51	0.45
1:C:125:ASP:HB2	5:C:448:HOH:O	2.17	0.45
1:B:37:ARG:HH12	1:B:41:PRO:CD	2.30	0.45
1:B:36:LYS:CD	1:C:118:LEU:HD21	2.29	0.45
1:C:111:ARG:NH1	1:C:111:ARG:CG	2.79	0.44
1:B:109:ARG:HG2	5:B:449:HOH:O	2.18	0.44
1:B:140:ASN:OD1	1:C:68:MET:O	2.36	0.43
1:B:133:ASP:HB3	1:C:54:GLN:HE22	1.83	0.43
1:C:1:MET:HE1	1:C:65:ARG:HD3	2.00	0.43
1:C:93:ASN:ND2	1:C:95:SER:H	2.16	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:37:ARG:HH12	1:B:41:PRO:HD2	1.84	0.43
1:C:20:SER:OG	1:C:105:VAL:HG22	2.19	0.43
1:B:12:VAL:O	1:B:16:THR:HG22	2.18	0.43
1:C:141:VAL:O	1:C:146:ARG:HD3	2.20	0.42
1:C:37:ARG:HA	1:C:37:ARG:HD2	1.37	0.42
1:C:58:THR:CG2	1:C:60:GLU:HB2	2.50	0.42
1:C:1:MET:HG3	5:C:444:HOH:O	2.19	0.41
1:C:132:SER:HA	1:C:135:LEU:HG	2.00	0.41
1:B:128:ARG:NE	1:C:68:MET:HE2	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/160 (99%)	155 (98%)	3 (2%)	0	100	100
1	B	158/160 (99%)	157 (99%)	1 (1%)	0	100	100
1	C	158/160 (99%)	156 (99%)	2 (1%)	0	100	100
All	All	474/480 (99%)	468 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/137 (100%)	131 (96%)	6 (4%)	33	45
1	B	137/137 (100%)	133 (97%)	4 (3%)	48	64
1	C	137/137 (100%)	131 (96%)	6 (4%)	33	45
All	All	411/411 (100%)	395 (96%)	16 (4%)	37	51

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	19	VAL
1	A	98	ASP
1	A	115	GLU
1	A	135	LEU
1	A	140	ASN
1	B	37	ARG
1	B	39	ASN
1	B	109	ARG
1	B	140	ASN
1	C	37	ARG
1	C	39	ASN
1	C	58	THR
1	C	109	ARG
1	C	111	ARG
1	C	115	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	GLN
1	A	54	GLN
1	A	93	ASN
1	A	124	HIS
1	A	140	ASN
1	B	54	GLN
1	B	93	ASN
1	B	140	ASN
1	C	54	GLN
1	C	81	GLN
1	C	93	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GDD	A	201	2	35,42,42	0.59	0	44,65,65	1.72	8 (18%)
3	GDD	B	201	2	35,42,42	0.66	0	44,65,65	1.66	8 (18%)
4	BMA	B	202	1	11,11,12	0.26	0	13,15,17	0.54	0
3	GDD	C	201	2	35,42,42	0.51	0	44,65,65	1.56	6 (13%)
4	BMA	C	202	1,2	11,11,12	0.27	0	13,15,17	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDD	A	201	2	-	0/19/59/59	0/4/4/4
3	GDD	B	201	2	-	0/19/59/59	0/4/4/4
4	BMA	B	202	1	-	0/2/19/22	0/1/1/1
3	GDD	C	201	2	-	0/19/59/59	0/4/4/4
4	BMA	C	202	1,2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	201	GDD	N3-C2-N1	-5.14	119.95	127.46
3	B	201	GDD	N3-C2-N1	-5.06	120.07	127.46
3	C	201	GDD	N3-C2-N1	-4.86	120.36	127.46
3	A	201	GDD	C5-C6-N1	-3.60	118.36	123.48
3	C	201	GDD	C5-C6-N1	-3.02	119.19	123.48
3	B	201	GDD	C5-C6-N1	-2.94	119.30	123.48
3	A	201	GDD	C4-C5-N7	-2.57	106.92	109.41
3	B	201	GDD	C1'-N9-C4	-2.36	122.55	126.64
3	B	201	GDD	C6-C5-C4	-2.34	118.51	120.84
3	C	201	GDD	C6-C5-C4	-2.22	118.64	120.84
3	A	201	GDD	C61-C51-C41	-2.17	107.93	113.00
3	B	201	GDD	O2B-PB-O3B	2.12	123.26	112.28
3	A	201	GDD	N2-C2-N1	2.12	120.64	117.24
3	B	201	GDD	N2-C2-N1	2.16	120.70	117.24
3	A	201	GDD	C11-O51-C51	2.40	118.24	113.72
3	C	201	GDD	C4'-O4'-C1'	2.45	112.37	109.77
3	B	201	GDD	C6-N1-C2	3.71	121.39	116.06
3	C	201	GDD	C6-N1-C2	3.79	121.51	116.06
3	B	201	GDD	C2-N3-C4	4.10	119.95	115.16
3	A	201	GDD	C6-N1-C2	4.13	122.00	116.06
3	C	201	GDD	C2-N3-C4	4.39	120.29	115.16
3	A	201	GDD	C2-N3-C4	4.44	120.34	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	201	GDD	3	0
4	B	202	BMA	4	0
3	C	201	GDD	4	0
4	C	202	BMA	8	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.