



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:17 am GMT

PDB ID : 4GT2  
Title : Crystal structure of DyP-type peroxidase (SCO3963) from *Streptomyces coelicolor*  
Authors : Lukk, T.; Hetta, A.M.A.; Jones, A.; Solbiati, J.; Majumdar, S.; Cronan, J.E.; Gerlt, J.A.; Nair, S.K.  
Deposited on : 2012-08-28  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	465	<div> <div>77%</div> <div>20%</div> </div>
1	B	465	<div> <div>74%</div> <div>5%</div> <div>21%</div> </div>
1	E	465	<div> <div>75%</div> <div>20%</div> </div>
1	G	465	<div> <div>77%</div> <div>21%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	505	-	-	-	X
3	GOL	G	503	-	-	-	X
4	OXY	E	504	-	-	-	X
4	OXY	G	502	-	-	-	X
5	ACT	B	504	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13783 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein SCO3963.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	369	Total	C	N	O	S	0	2	0
			2822	1746	532	532	12			
1	A	371	Total	C	N	O	S	0	4	0
			2855	1763	542	539	11			
1	E	370	Total	C	N	O	S	0	1	0
			2822	1746	535	530	11			
1	G	369	Total	C	N	O	S	0	2	0
			2823	1746	532	534	11			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	EXPRESSION TAG	UNP Q9ZBW9
B	-18	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
B	-17	SER	-	EXPRESSION TAG	UNP Q9ZBW9
B	-16	SER	-	EXPRESSION TAG	UNP Q9ZBW9
B	-15	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
B	-14	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
B	-13	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
B	-12	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
B	-11	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
B	-10	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
B	-9	SER	-	EXPRESSION TAG	UNP Q9ZBW9
B	-8	SER	-	EXPRESSION TAG	UNP Q9ZBW9
B	-7	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
B	-6	LEU	-	EXPRESSION TAG	UNP Q9ZBW9
B	-5	VAL	-	EXPRESSION TAG	UNP Q9ZBW9
B	-4	PRO	-	EXPRESSION TAG	UNP Q9ZBW9
B	-3	ARG	-	EXPRESSION TAG	UNP Q9ZBW9
B	-2	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
B	-1	SER	-	EXPRESSION TAG	UNP Q9ZBW9
B	0	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
A	-19	MET	-	EXPRESSION TAG	UNP Q9ZBW9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
A	-17	SER	-	EXPRESSION TAG	UNP Q9ZBW9
A	-16	SER	-	EXPRESSION TAG	UNP Q9ZBW9
A	-15	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
A	-14	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
A	-13	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
A	-12	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
A	-11	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
A	-10	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
A	-9	SER	-	EXPRESSION TAG	UNP Q9ZBW9
A	-8	SER	-	EXPRESSION TAG	UNP Q9ZBW9
A	-7	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
A	-6	LEU	-	EXPRESSION TAG	UNP Q9ZBW9
A	-5	VAL	-	EXPRESSION TAG	UNP Q9ZBW9
A	-4	PRO	-	EXPRESSION TAG	UNP Q9ZBW9
A	-3	ARG	-	EXPRESSION TAG	UNP Q9ZBW9
A	-2	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
A	-1	SER	-	EXPRESSION TAG	UNP Q9ZBW9
A	0	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
E	-19	MET	-	EXPRESSION TAG	UNP Q9ZBW9
E	-18	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
E	-17	SER	-	EXPRESSION TAG	UNP Q9ZBW9
E	-16	SER	-	EXPRESSION TAG	UNP Q9ZBW9
E	-15	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
E	-14	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
E	-13	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
E	-12	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
E	-11	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
E	-10	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
E	-9	SER	-	EXPRESSION TAG	UNP Q9ZBW9
E	-8	SER	-	EXPRESSION TAG	UNP Q9ZBW9
E	-7	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
E	-6	LEU	-	EXPRESSION TAG	UNP Q9ZBW9
E	-5	VAL	-	EXPRESSION TAG	UNP Q9ZBW9
E	-4	PRO	-	EXPRESSION TAG	UNP Q9ZBW9
E	-3	ARG	-	EXPRESSION TAG	UNP Q9ZBW9
E	-2	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
E	-1	SER	-	EXPRESSION TAG	UNP Q9ZBW9
E	0	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
G	-19	MET	-	EXPRESSION TAG	UNP Q9ZBW9
G	-18	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
G	-17	SER	-	EXPRESSION TAG	UNP Q9ZBW9

*Continued on next page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	-16	SER	-	EXPRESSION TAG	UNP Q9ZBW9
G	-15	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
G	-14	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
G	-13	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
G	-12	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
G	-11	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
G	-10	HIS	-	EXPRESSION TAG	UNP Q9ZBW9
G	-9	SER	-	EXPRESSION TAG	UNP Q9ZBW9
G	-8	SER	-	EXPRESSION TAG	UNP Q9ZBW9
G	-7	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
G	-6	LEU	-	EXPRESSION TAG	UNP Q9ZBW9
G	-5	VAL	-	EXPRESSION TAG	UNP Q9ZBW9
G	-4	PRO	-	EXPRESSION TAG	UNP Q9ZBW9
G	-3	ARG	-	EXPRESSION TAG	UNP Q9ZBW9
G	-2	GLY	-	EXPRESSION TAG	UNP Q9ZBW9
G	-1	SER	-	EXPRESSION TAG	UNP Q9ZBW9
G	0	HIS	-	EXPRESSION TAG	UNP Q9ZBW9

- # HEM

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	E	1	Total	C	Fe	N	O	
			43	34	1	4	4	
2	G	1	Total	C	Fe	N	O	
			43	34	1	4	4	

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O		
			6	3	3		
3	B	1	Total	C	O		
			6	3	3		
3	A	1	Total	C	O		
			6	3	3		
3	A	1	Total	C	O		
			6	3	3		
3	E	1	Total	C	O		
			6	3	3		
3	E	1	Total	C	O		
			6	3	3		
3	E	1	Total	C	O		
			6	3	3		
3	G	1	Total	C	O		
			6	3	3		

- Molecule 4 is OXYGEN MOLECULE (three-letter code: OXY) (formula:  $O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total O 2 2	0	0
4	A	1	Total O 2 2	0	0
4	E	1	Total O 2 2	0	0
4	G	1	Total O 2 2	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	570	Total	O	0	0
			570	570		
6	A	567	Total	O	0	0
			567	567		
6	E	527	Total	O	0	0
			527	527		
6	G	565	Total	O	0	0
			565	565		



- Molecule 1: Putative uncharacterized protein SCO3963



● Molecule 1: Putative uncharacterized protein SCO3963



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.87Å 203.20Å 76.32Å 90.00° 110.73° 90.00°	Depositor
Resolution (Å)	29.20 – 1.80 29.20 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.20-1.80) 99.7 (29.20-1.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1066)	Depositor
R, $R_{free}$	0.147 , 0.185 0.145 , 0.182	Depositor DCC
$R_{free}$ test set	7140 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	11.8	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 59.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.177 for h,-k,-h-l	Xtriage
Reported twinning fraction	0.010 for h,-k,-h-l	Depositor
Outliers	0 of 141255 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	13783	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT, OXY, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/2917	0.55	0/3943
1	B	0.37	0/2884	0.57	1/3899 (0.0%)
1	E	0.34	0/2884	0.52	0/3899
1	G	0.36	0/2885	0.54	0/3901
All	All	0.36	0/11570	0.55	1/15642 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	441	ARG	NE-CZ-NH1	-9.88	115.36	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2855	0	2753	6	0
1	B	2822	0	2725	15	0
1	E	2822	0	2728	11	0
1	G	2823	0	2722	4	0
2	A	43	0	30	0	0
2	B	43	0	30	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	43	0	30	0	0
2	G	43	0	30	0	0
3	A	12	0	16	0	0
3	B	12	0	16	1	0
3	E	18	0	24	0	0
3	G	6	0	8	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	E	2	0	0	0	0
4	G	2	0	0	0	0
5	B	4	0	3	0	0
6	A	567	0	0	2	0
6	B	570	0	0	5	0
6	E	527	0	0	4	0
6	G	565	0	0	0	0
All	All	13783	0	11115	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 36 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:235:GLY:H	3:B:505:GOL:H31	1.37	0.89
1:A:119[A]:ARG:NH2	6:A:803:HOH:O	2.23	0.71
1:B:97:ARG:NH2	6:B:1099:HOH:O	2.27	0.66
1:E:406:ARG:NH2	6:E:866:HOH:O	2.25	0.61
1:E:265:ARG:NH2	6:E:955:HOH:O	2.35	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/465 (80%)	365 (98%)	6 (2%)	0	100	100
1	B	367/465 (79%)	360 (98%)	7 (2%)	0	100	100
1	E	367/465 (79%)	360 (98%)	7 (2%)	0	100	100
1	G	367/465 (79%)	362 (99%)	5 (1%)	0	100	100
All	All	1472/1860 (79%)	1447 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/345 (83%)	281 (98%)	5 (2%)	66	55
1	B	283/345 (82%)	280 (99%)	3 (1%)	78	72
1	E	282/345 (82%)	279 (99%)	3 (1%)	78	72
1	G	283/345 (82%)	281 (99%)	2 (1%)	87	84
All	All	1134/1380 (82%)	1121 (99%)	13 (1%)	75	72

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	271	GLU
1	A	345	GLU
1	E	416	GLN
1	A	168	GLU
1	E	394	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	285	ASN
1	E	416	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	A	501	1,4	28,50,50	2.14	8 (28%)	17,82,82	1.76	7 (41%)
3	GOL	A	502	-	5,5,5	0.34	0	5,5,5	0.48	0
4	OXY	A	503	2	1,1,1	0.28	0	0,0,0	0.00	-
3	GOL	A	504	-	5,5,5	0.31	0	5,5,5	0.28	0
2	HEM	B	501	1	28,50,50	2.14	6 (21%)	17,82,82	1.69	5 (29%)
3	GOL	B	502	-	5,5,5	0.30	0	5,5,5	0.20	0
4	OXY	B	503	-	1,1,1	0.28	0	0,0,0	0.00	-
5	ACT	B	504	-	1,3,3	1.38	0	0,3,3	0.00	-
3	GOL	B	505	-	5,5,5	0.62	0	5,5,5	0.70	0
2	HEM	E	501	1,4	28,50,50	2.14	7 (25%)	17,82,82	1.54	4 (23%)
3	GOL	E	502	-	5,5,5	0.35	0	5,5,5	0.18	0
3	GOL	E	503	-	5,5,5	0.33	0	5,5,5	0.30	0
4	OXY	E	504	2	1,1,1	0.30	0	0,0,0	0.00	-
3	GOL	E	505	-	5,5,5	0.48	0	5,5,5	0.42	0
2	HEM	G	501	1,4	28,50,50	2.25	8 (28%)	17,82,82	1.78	5 (29%)
4	OXY	G	502	2	1,1,1	0.28	0	0,0,0	0.00	-
3	GOL	G	503	-	5,5,5	0.35	0	5,5,5	0.21	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	1,4	-	0/6/54/54	0/0/8/8
3	GOL	A	502	-	-	0/4/4/4	0/0/0/0
4	OXY	A	503	2	-	0/0/0/0	0/0/0/0
3	GOL	A	504	-	-	0/4/4/4	0/0/0/0
2	HEM	B	501	1	-	0/6/54/54	0/0/8/8
3	GOL	B	502	-	-	0/4/4/4	0/0/0/0
4	OXY	B	503	-	-	0/0/0/0	0/0/0/0
5	ACT	B	504	-	-	0/0/0/0	0/0/0/0
3	GOL	B	505	-	-	0/4/4/4	0/0/0/0
2	HEM	E	501	1,4	-	0/6/54/54	0/0/8/8
3	GOL	E	502	-	-	0/4/4/4	0/0/0/0
3	GOL	E	503	-	-	0/4/4/4	0/0/0/0
4	OXY	E	504	2	-	0/0/0/0	0/0/0/0
3	GOL	E	505	-	-	0/4/4/4	0/0/0/0
2	HEM	G	501	1,4	-	0/6/54/54	0/0/8/8
4	OXY	G	502	2	-	0/0/0/0	0/0/0/0
3	GOL	G	503	-	-	0/4/4/4	0/0/0/0

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	501	HEM	C3B-C2B	-4.33	1.34	1.40
2	G	501	HEM	C3C-C2C	-3.92	1.35	1.40
2	A	501	HEM	C3C-C2C	-3.89	1.35	1.40
2	B	501	HEM	C3B-C2B	-3.77	1.35	1.40
2	B	501	HEM	C3C-C2C	-3.62	1.35	1.40

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	501	HEM	CBD-CAD-C3D	-3.45	105.88	112.47
2	G	501	HEM	C1D-C2D-C3D	-3.03	104.89	107.00
2	E	501	HEM	CBD-CAD-C3D	-2.93	106.87	112.47
2	G	501	HEM	CMA-C3A-C4A	-2.89	124.03	128.46
2	B	501	HEM	CBD-CAD-C3D	-2.75	107.22	112.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	505	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	371/465 (79%)	-0.55	2 (0%) 90 89	6, 11, 27, 43	0
1	B	369/465 (79%)	-0.58	3 (0%) 86 84	6, 10, 29, 59	0
1	E	370/465 (79%)	-0.44	10 (2%) 55 50	7, 13, 33, 57	0
1	G	369/465 (79%)	-0.57	3 (0%) 86 84	6, 11, 26, 51	0
All	All	1479/1860 (79%)	-0.54	18 (1%) 79 77	6, 11, 29, 59	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	445	GLY	7.5
1	B	342	GLY	5.6
1	G	342	GLY	5.3
1	E	342	GLY	5.0
1	G	445	GLY	4.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	505	6/6	0.81	0.17	14.17	17,27,32,34	0
4	OXY	G	502	2/2	0.92	0.14	4.87	21,21,21,23	0
4	OXY	E	504	2/2	0.95	0.13	3.43	26,26,26,28	0
3	GOL	G	503	6/6	0.95	0.12	3.39	15,19,22,23	0
5	ACT	B	504	4/4	0.82	0.14	2.05	22,23,29,31	0
3	GOL	B	502	6/6	0.96	0.10	1.08	11,13,16,18	0
3	GOL	A	502	6/6	0.96	0.09	0.68	18,20,21,23	0
3	GOL	E	503	6/6	0.96	0.08	0.67	16,17,25,29	0
2	HEM	A	501	43/43	0.99	0.09	0.49	4,7,10,13	0
3	GOL	E	502	6/6	0.96	0.08	0.10	9,16,18,22	0
2	HEM	G	501	43/43	0.99	0.07	0.03	4,6,11,12	0
2	HEM	B	501	43/43	0.99	0.07	-0.13	2,7,9,11	0
2	HEM	E	501	43/43	0.99	0.07	-0.34	5,9,13,15	0
3	GOL	E	505	6/6	0.96	0.07	-0.40	16,16,19,24	0
3	GOL	A	504	6/6	0.96	0.06	-0.43	11,15,16,19	0
4	OXY	B	503	2/2	0.93	0.14	-	26,26,26,29	0
4	OXY	A	503	2/2	0.90	0.16	-	27,27,27,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.