



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 08:54 pm GMT

PDB ID : 1GVE
Title : AFLATOXIN ALDEHYDE REDUCTASE (AKR7A1) FROM RAT LIVER
Authors : Kozma, E.; Brown, E.; Ellis, E.M.; Lapthorn, A.J.
Deposited on : 2002-02-08
Resolution : 1.38 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2266 (1.40-1.36)
Ramachandran outliers	110173	2215 (1.40-1.36)
Sidechain outliers	110143	2214 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	327	
1	B	327	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	1335	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 5567 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AFLATOXIN B1 ALDEHYDE REDUCTASE MEMBER 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	3	0
			2552	1626	437	470	19			
1	B	301	Total	C	N	O	S	0	7	0
			2376	1511	414	434	17			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			13	6	7		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	190	Total	O	0	1
			191	191		
5	B	314	Total	O	0	1
			315	315		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: AFLATOXIN B1 ALDEHYDE REDUCTASE MEMBER 3



• Molecule 1: AFLATOXIN B1 ALDEHYDE REDUCTASE MEMBER 3



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	125.54Å 64.68Å 112.84Å 90.00° 91.03° 90.00°	Depositor
Resolution (Å)	28.75 – 1.38	Depositor
% Data completeness (in resolution range)	73.7 (28.75-1.38)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.158 , 0.180	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5567	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAP, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	2/2632 (0.1%)	0.78	1/3568 (0.0%)
1	B	0.90	2/2467 (0.1%)	0.90	4/3335 (0.1%)
All	All	0.83	4/5099 (0.1%)	0.84	5/6903 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	160[A]	MET	CG-SD	-10.39	1.54	1.81
1	A	160[B]	MET	CG-SD	-10.39	1.54	1.81
1	B	97	ARG	CD-NE	-6.57	1.35	1.46
1	B	178	GLU	CD-OE1	-5.66	1.19	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	97	ARG	NE-CZ-NH2	-12.56	114.02	120.30
1	B	63	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	B	265	ARG	NE-CZ-NH2	-6.71	116.94	120.30
1	B	97	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	180	PHE	CB-CG-CD1	5.35	124.54	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2552	0	2477	28	0
1	B	2376	0	2345	33	0
2	A	48	0	25	4	0
3	A	12	0	16	1	0
3	B	60	0	79	11	0
4	A	13	0	5	0	0
5	A	191	0	0	6	0
5	B	315	0	0	9	1
All	All	5567	0	4947	64	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 64 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:GLU:CD	5:A:2019:HOH:O	2.11	0.87
1:B:274:LYS:HE3	5:B:2248:HOH:O	1.83	0.78
1:A:293:GLN:HE21	2:A:1328:NAP:H62A	1.35	0.75
1:A:165:GLN:OE1	2:A:1328:NAP:H2N	1.91	0.71
1:A:293:GLN:NE2	2:A:1328:NAP:H62A	1.89	0.70

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:2040:HOH:O	5:B:2040:HOH:O[2_656]	0.87	1.33

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/327 (99%)	319 (98%)	5 (2%)	1 (0%)	44	17
1	B	303/327 (93%)	297 (98%)	6 (2%)	0	100	100
All	All	628/654 (96%)	616 (98%)	11 (2%)	1 (0%)	51	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	254	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/273 (98%)	262 (97%)	7 (3%)	51	15
1	B	253/273 (93%)	251 (99%)	2 (1%)	85	63
All	All	522/546 (96%)	513 (98%)	9 (2%)	68	32

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	208	GLN
1	B	285	MET
1	A	285	MET
1	A	160[B]	MET
1	A	274	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	128	HIS
1	B	293	GLN
1	B	237	HIS
1	A	272	GLN
1	B	165	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAP	A	1328	-	44,52,52	1.68	8 (18%)	51,80,80	1.71	8 (15%)
3	GOL	A	1329	-	5,5,5	0.62	0	5,5,5	0.41	0
3	GOL	A	1330	-	5,5,5	0.50	0	5,5,5	1.42	1 (20%)
4	CIT	A	1331	-	3,12,12	0.80	0	3,17,17	1.34	1 (33%)
3	GOL	B	1328	-	5,5,5	0.50	0	5,5,5	0.50	0
3	GOL	B	1329	-	5,5,5	0.87	0	5,5,5	0.28	0
3	GOL	B	1330	-	5,5,5	0.24	0	5,5,5	0.40	0
3	GOL	B	1331	-	5,5,5	0.88	0	5,5,5	0.91	0
3	GOL	B	1332	-	5,5,5	0.45	0	5,5,5	0.41	0
3	GOL	B	1333	-	5,5,5	0.41	0	5,5,5	0.66	0
3	GOL	B	1334	-	5,5,5	0.36	0	5,5,5	0.72	0
3	GOL	B	1335	-	5,5,5	0.29	0	5,5,5	0.69	0
3	GOL	B	1336	-	5,5,5	1.32	1 (20%)	5,5,5	0.61	0
3	GOL	B	1337	-	5,5,5	0.36	0	5,5,5	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	1328	-	-	0/27/67/67	0/5/5/5
3	GOL	A	1329	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1330	-	-	0/4/4/4	0/0/0/0
4	CIT	A	1331	-	-	0/6/16/16	0/0/0/0
3	GOL	B	1328	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1329	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1330	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1331	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1332	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1333	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1334	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1335	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1336	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1337	-	-	0/4/4/4	0/0/0/0

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1328	NAP	C3N-C7N	-4.65	1.43	1.50
3	B	1336	GOL	O2-C2	-2.65	1.35	1.43
2	A	1328	NAP	C6N-N1N	2.46	1.41	1.35
2	A	1328	NAP	C4N-C3N	2.60	1.43	1.39
2	A	1328	NAP	C5N-C4N	3.07	1.44	1.38

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1328	NAP	N3A-C2A-N1A	-8.38	121.56	128.86
2	A	1328	NAP	C6N-C5N-C4N	-2.88	115.09	119.44
2	A	1328	NAP	C1B-N9A-C4A	-2.50	122.31	126.64
2	A	1328	NAP	O2D-C2D-C1D	-2.21	104.69	111.61
2	A	1328	NAP	C4D-O4D-C1D	-2.13	107.51	109.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1328	NAP	4	0
3	A	1330	GOL	1	0
3	B	1328	GOL	1	0
3	B	1331	GOL	3	0
3	B	1332	GOL	1	0
3	B	1333	GOL	1	0
3	B	1335	GOL	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.