



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 7, 2017 – 07:47 AM EST

PDB ID : 5GWJ  
Title : Structure of a Human topoisomerase IIbeta fragment in complex with DNA and E7873S  
Authors : Wang, Y.R.; Chen, S.F.; Wu, C.C.; Chan, N.L.  
Deposited on : unknown  
Resolution : 2.57 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3689 (2.60-2.52)
Clashscore	112137	4096 (2.60-2.52)
Ramachandran outliers	110173	4037 (2.60-2.52)
Sidechain outliers	110143	4037 (2.60-2.52)
RSRZ outliers	101464	3700 (2.60-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	803	<div> <div>4%</div> <div>76%</div> <div>8%</div> <div>16%</div> </div>
1	B	803	<div> <div>2%</div> <div>75%</div> <div>9%</div> <div>16%</div> </div>
2	C	8	<div> <div>13%</div> <div>100%</div> </div>
2	E	8	<div> <div>100%</div> </div>
3	D	12	<div> <div>92%</div> <div>8%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	12	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MG	A	1304	-	-	-	X
6	MG	B	1303	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 12264 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase 2-beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	677	Total	C	N	O	S	0	9	0
			5508	3504	954	1026	24			
1	B	676	Total	C	N	O	S	0	5	0
			5462	3477	947	1014	24			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	419	MET	-	expression tag	UNP Q02880
A	420	ALA	-	expression tag	UNP Q02880
A	421	SER	-	expression tag	UNP Q02880
A	422	TRP	-	expression tag	UNP Q02880
A	423	SER	-	expression tag	UNP Q02880
A	424	HIS	-	expression tag	UNP Q02880
A	425	PRO	-	expression tag	UNP Q02880
A	426	GLN	-	expression tag	UNP Q02880
A	427	PHE	-	expression tag	UNP Q02880
A	428	GLU	-	expression tag	UNP Q02880
A	429	LYS	-	expression tag	UNP Q02880
A	430	GLY	-	expression tag	UNP Q02880
A	431	ALA	-	expression tag	UNP Q02880
A	432	ASP	-	expression tag	UNP Q02880
A	433	ASP	-	expression tag	UNP Q02880
A	434	ASP	-	expression tag	UNP Q02880
A	435	ASP	-	expression tag	UNP Q02880
A	436	LYS	-	expression tag	UNP Q02880
A	437	VAL	-	expression tag	UNP Q02880
A	438	PRO	-	expression tag	UNP Q02880
A	439	ASP	-	expression tag	UNP Q02880
A	440	PRO	-	expression tag	UNP Q02880
A	441	THR	-	expression tag	UNP Q02880
A	442	SER	-	expression tag	UNP Q02880
A	443	VAL	-	expression tag	UNP Q02880

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	444	ASP	-	expression tag	UNP Q02880
A	1202	GLY	-	expression tag	UNP Q02880
A	1203	ALA	-	expression tag	UNP Q02880
A	1204	PRO	-	expression tag	UNP Q02880
A	1205	GLY	-	expression tag	UNP Q02880
A	1206	PHE	-	expression tag	UNP Q02880
A	1207	SER	-	expression tag	UNP Q02880
A	1208	SER	-	expression tag	UNP Q02880
A	1209	ILE	-	expression tag	UNP Q02880
A	1210	SER	-	expression tag	UNP Q02880
A	1211	ALA	-	expression tag	UNP Q02880
A	1212	HIS	-	expression tag	UNP Q02880
A	1213	HIS	-	expression tag	UNP Q02880
A	1214	HIS	-	expression tag	UNP Q02880
A	1215	HIS	-	expression tag	UNP Q02880
A	1216	HIS	-	expression tag	UNP Q02880
A	1217	HIS	-	expression tag	UNP Q02880
A	1218	HIS	-	expression tag	UNP Q02880
A	1219	HIS	-	expression tag	UNP Q02880
A	1220	HIS	-	expression tag	UNP Q02880
A	1221	HIS	-	expression tag	UNP Q02880
B	419	MET	-	expression tag	UNP Q02880
B	420	ALA	-	expression tag	UNP Q02880
B	421	SER	-	expression tag	UNP Q02880
B	422	TRP	-	expression tag	UNP Q02880
B	423	SER	-	expression tag	UNP Q02880
B	424	HIS	-	expression tag	UNP Q02880
B	425	PRO	-	expression tag	UNP Q02880
B	426	GLN	-	expression tag	UNP Q02880
B	427	PHE	-	expression tag	UNP Q02880
B	428	GLU	-	expression tag	UNP Q02880
B	429	LYS	-	expression tag	UNP Q02880
B	430	GLY	-	expression tag	UNP Q02880
B	431	ALA	-	expression tag	UNP Q02880
B	432	ASP	-	expression tag	UNP Q02880
B	433	ASP	-	expression tag	UNP Q02880
B	434	ASP	-	expression tag	UNP Q02880
B	435	ASP	-	expression tag	UNP Q02880
B	436	LYS	-	expression tag	UNP Q02880
B	437	VAL	-	expression tag	UNP Q02880
B	438	PRO	-	expression tag	UNP Q02880
B	439	ASP	-	expression tag	UNP Q02880

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	440	PRO	-	expression tag	UNP Q02880
B	441	THR	-	expression tag	UNP Q02880
B	442	SER	-	expression tag	UNP Q02880
B	443	VAL	-	expression tag	UNP Q02880
B	444	ASP	-	expression tag	UNP Q02880
B	1202	GLY	-	expression tag	UNP Q02880
B	1203	ALA	-	expression tag	UNP Q02880
B	1204	PRO	-	expression tag	UNP Q02880
B	1205	GLY	-	expression tag	UNP Q02880
B	1206	PHE	-	expression tag	UNP Q02880
B	1207	SER	-	expression tag	UNP Q02880
B	1208	SER	-	expression tag	UNP Q02880
B	1209	ILE	-	expression tag	UNP Q02880
B	1210	SER	-	expression tag	UNP Q02880
B	1211	ALA	-	expression tag	UNP Q02880
B	1212	HIS	-	expression tag	UNP Q02880
B	1213	HIS	-	expression tag	UNP Q02880
B	1214	HIS	-	expression tag	UNP Q02880
B	1215	HIS	-	expression tag	UNP Q02880
B	1216	HIS	-	expression tag	UNP Q02880
B	1217	HIS	-	expression tag	UNP Q02880
B	1218	HIS	-	expression tag	UNP Q02880
B	1219	HIS	-	expression tag	UNP Q02880
B	1220	HIS	-	expression tag	UNP Q02880
B	1221	HIS	-	expression tag	UNP Q02880


- Molecule 2 is a DNA chain called DNA (5'-D(P\*AP\*GP\*CP\*CP\*GP\*AP\*GP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	8	Total 165	C 77	N 34	O 46	P 8	0	0	0
2	E	8	Total 165	C 77	N 34	O 46	P 8	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(P\*TP\*GP\*CP\*AP\*GP\*CP\*TP\*CP\*GP\*GP\*CP\*T)-3').


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	12	Total 245	C 116	N 43	O 74	P 12	0	0	0
3	F	12	Total 245	C 116	N 43	O 74	P 12	0	0	0

- Molecule 4 is chloro{4,5-di(amino-kappaN)-N-[9-(4-hydroxy-3,5-dimethoxyphenyl)-8-oxo-5,5a,6,8,8a,9-hexahydro-2H-furo[3',4':6,7]naphtho[2,3-d][1,3]dioxol-5-yl]pentanamide}platinum (three-letter code: N2S) (formula: C<sub>26</sub>H<sub>31</sub>ClN<sub>3</sub>O<sub>8</sub>Pt).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Cl	N	O	Pt		
4	A	1	39	26	1	3	8	1	0	1
4	B	1	39	26	1	3	8	1	0	1

- Molecule 5 is chloro{4,5-di(amino-kappaN)-N-[9-(4-hydroxy-3,5-dimethoxyphenyl)-8-oxo-5,5a,6,8,8a,9-hexahydro-2H-furo[3',4':6,7]naphtho[2,3-d][1,3]dioxol-5-yl]pentanamide}platinum (three-letter code: N2W) (formula: C<sub>26</sub>H<sub>31</sub>ClN<sub>3</sub>O<sub>8</sub>Pt).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	Pt	0	1
			39	26	1	3	8	1		
5	B	1	Total	C	Cl	N	O	Pt	0	1
			39	26	1	3	8	1		

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	Mg	0	0
			2	2		
6	A	2	Total	Mg	0	0
			2	2		
6	F	1	Total	Mg	0	0
			1	1		
6	E	1	Total	Mg	0	0
			1	1		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	130	Total	O	0	0
			130	130		
7	B	144	Total	O	0	0
			144	144		
7	C	7	Total	O	0	0
			7	7		
7	D	14	Total	O	0	0
			14	14		
7	E	8	Total	O	0	0
			8	8		
7	F	9	Total	O	0	0
			9	9		



These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A:
- 
- 4%
- 76%
- 8%
- 16%
- MET  
SER  
TRP  
HIS  
PRO  
GLN  
PHE  
GLU  
LYS  
GLY  
ALA  
ASP  
ASP  
LYS  
VAL  
PRO  
THR  
SER  
VAL  
ASP  
LYS  
ILE  
PHE  
TYR  
SER  
GLN  
LYS  
LEU  
SER  
PHE  
TYR  
SER  
ILE  
PRO  
GLU  
PHE  
ASP  
ASP  
TRP  
LYS  
LYS  
HIS  
ILE  
GLU  
ASN  
GLN  
LYS  
ALA  
TRP  
D495  
D496  
R503  
L529
- GLY  
LEU  
GLY  
T634  
S635  
K638  
E642  
Y643  
R688  
R691  
R692  
L693  
H694  
G695  
L696  
PRO  
GLU  
GLN  
PHE  
LEU  
TYR  
GLY  
THR  
ALA  
THR  
K707  
P732  
E757  
A768  
Y773  
Q778  
W782  
G807  
G813  
S818  
P819  
R820  
F823  
R831  
L832  
L833  
L841  
F844
- L845  
D846  
D847  
D848  
R849  
Q850  
R851  
R852  
E853  
P854  
I858  
M863  
A869  
R834  
E838  
V944  
R945  
V955  
M959  
K983  
M1014  
K1027  
R1042  
R1049  
L1072  
E1082  
H1083  
R1084  
M1092  
W1106  
K1107  
E1108  
A1109  
Q1110  
E1111  
LYS  
ALA  
ALA  
GLU  
GLU  
ASP  
GLU  
THR
- GLN  
ASN  
GLN  
HIS  
ASP  
ASP  
SER  
SER  
SER  
ASP  
SER  
GLY  
THR  
PRO  
PRO  
G1135  
L1146  
I1158  
V1168  
K1172  
D1201  
GLY  
PHE  
PRO  
GLY  
PHE  
SER  
ILE  
SER  
ALA  
HIS  
HIS  
HIS  
HIS  
HIS  
HIS  
HIS

- [illegible]



- Molecule 2: DNA (5'-D(P\*AP\*GP\*CP\*CP\*GP\*AP\*GP\*C)-3')



- Molecule 2: DNA (5'-D(P\*AP\*GP\*CP\*CP\*GP\*AP\*GP\*C)-3')



There are no outlier residues recorded for this chain.

- Molecule 3: DNA (5'-D(P\*TP\*GP\*CP\*AP\*GP\*CP\*TP\*CP\*GP\*GP\*CP\*T)-3')



- Molecule 3: DNA (5'-D(P\*TP\*GP\*CP\*AP\*GP\*CP\*TP\*CP\*GP\*GP\*CP\*T)-3')



There are no outlier residues recorded for this chain.

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.18 Å 176.99 Å 94.41 Å 90.00° 111.50° 90.00°	Depositor
Resolution (Å)	29.68 – 2.57 29.68 – 2.57	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.68-2.57) 99.5 (29.68-2.57)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.48	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.85 (at 2.57 Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.183 , 0.226 0.192 , 0.237	Depositor DCC
$R_{free}$ test set	2005 reflections (2.65%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.9	Xtriage
Anisotropy	0.186	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12264	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: N2S, MG, N2W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/5638	0.49	0/7593
1	B	0.32	0/5580	0.50	0/7516
2	C	0.67	0/185	0.75	0/283
2	E	0.80	0/185	0.82	0/283
3	D	0.78	0/273	0.99	0/419
3	F	0.72	0/273	0.95	0/419
All	All	0.37	0/12134	0.54	0/16513

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5508	0	5561	35	0
1	B	5462	0	5496	44	0
2	C	165	0	89	0	0
2	E	165	0	89	0	0
3	D	245	0	136	1	0
3	F	245	0	136	0	0
4	A	39	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	39	0	0	0	0
5	A	39	0	0	0	0
5	B	39	0	0	0	0
6	A	2	0	0	0	0
6	B	2	0	0	0	0
6	E	1	0	0	0	0
6	F	1	0	0	0	0
7	A	130	0	0	2	0
7	B	144	0	0	4	0
7	C	7	0	0	0	0
7	D	14	0	0	0	0
7	E	8	0	0	0	0
7	F	9	0	0	1	0
All	All	12264	0	11507	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:732:PRO:HG2	1:B:869:ALA:HB1	1.72	0.71
1:B:467:HIS:HB3	1:B:496:ARG:HH12	1.56	0.70
1:A:456:LYS:NZ	3:D:11:DC:OP1	2.26	0.68
1:B:651:ARG:NH1	7:B:1402:HOH:O	2.27	0.68
1:A:732:PRO:HG3	1:A:869:ALA:HB1	1.76	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	678/803 (84%)	655 (97%)	22 (3%)	1 (0%)	55	77
1	B	673/803 (84%)	649 (96%)	24 (4%)	0	100	100
All	All	1351/1606 (84%)	1304 (96%)	46 (3%)	1 (0%)	55	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	852	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	603/704 (86%)	603 (100%)	0	100	100
1	B	593/704 (84%)	592 (100%)	1 (0%)	94	98
All	All	1196/1408 (85%)	1195 (100%)	1 (0%)	94	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1096	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	786	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	N2S	A	1301[A]	1	40,44,44	3.54	15 (37%)	54,65,65	2.86	20 (37%)
5	N2W	A	1302[B]	1	40,44,44	3.56	15 (37%)	54,65,65	2.90	20 (37%)
4	N2S	B	1301[A]	1	40,44,44	3.55	14 (35%)	54,65,65	2.85	20 (37%)
5	N2W	B	1302[B]	1	40,44,44	3.58	14 (35%)	54,65,65	2.88	19 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	N2S	A	1301[A]	1	-	0/17/61/61	0/6/6/6
5	N2W	A	1302[B]	1	-	0/17/61/61	0/6/6/6
4	N2S	B	1301[A]	1	-	0/17/61/61	0/6/6/6
5	N2W	B	1302[B]	1	-	0/17/61/61	0/6/6/6

The worst 5 of 58 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1302[B]	N2W	C16-N17	-4.92	1.36	1.46
4	A	1301[A]	N2S	C16-N17	-4.81	1.36	1.46

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1301[A]	N2S	C16-N17	-4.69	1.36	1.46
5	A	1302[B]	N2W	C16-N17	-4.48	1.37	1.46
5	A	1302[B]	N2W	C34-C06	-3.60	1.49	1.56

The worst 5 of 79 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1302[B]	N2W	C05-C06-C34	-10.91	96.69	113.31
5	A	1302[B]	N2W	C05-C06-C34	-10.72	96.98	113.31
4	A	1301[A]	N2S	C05-C06-C34	-10.15	97.85	113.31
4	B	1301[A]	N2S	C05-C06-C34	-10.09	97.94	113.31
4	B	1301[A]	N2S	C30-O31-C32	-5.30	105.32	110.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	677/803 (84%)	-0.04	30 (4%) 35 32	24, 44, 82, 105	0
1	B	676/803 (84%)	-0.16	19 (2%) 53 51	25, 41, 77, 109	0
2	C	8/8 (100%)	-0.28	1 (12%) 4 3	35, 37, 61, 89	0
2	E	8/8 (100%)	-0.43	0 100 100	31, 35, 55, 85	0
3	D	12/12 (100%)	-0.56	0 100 100	29, 40, 51, 57	0
3	F	12/12 (100%)	-0.38	0 100 100	33, 41, 62, 71	0
All	All	1393/1646 (84%)	-0.11	50 (3%) 43 40	24, 43, 79, 109	0

The worst 5 of 50 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	963	THR	5.0
1	A	635	SER	5.0
1	A	849	ASN	4.5
1	A	848	ASP	4.5
1	A	696	LEU	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	MG	A	1304	1/1	0.94	0.86	26.30	59,59,59,59	0
6	MG	B	1303	1/1	0.96	0.85	23.98	60,60,60,60	0
4	N2S	B	1301[A]	39/39	0.97	0.13	-0.40	30,38,90,107	39
5	N2W	A	1302[B]	39/39	0.96	0.14	-0.47	38,45,91,116	39
4	N2S	A	1301[A]	39/39	0.96	0.14	-0.49	38,45,91,115	39
5	N2W	B	1302[B]	39/39	0.97	0.13	-0.49	30,38,91,107	39
6	MG	B	1304	1/1	0.86	0.24	-	47,47,47,47	0
6	MG	A	1303	1/1	0.89	0.32	-	47,47,47,47	0
6	MG	E	101	1/1	0.95	0.20	-	57,57,57,57	0
6	MG	F	101	1/1	0.96	0.39	-	57,57,57,57	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.