



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 05:49 am GMT

PDB ID : 4H02  
Title : Crystal structure of P. falciparum Lysyl-tRNA synthetase  
Authors : Khan, S.; Garg, A.; Sharma, A.  
Deposited on : 2012-09-07  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	507	<div><div>0%</div><div>50%38%7%</div></div>
1	B	507	<div><div>2%</div><div>52%33%8%7%</div></div>
1	C	507	<div><div>2%</div><div>51%35%6%7%</div></div>
1	D	507	<div><div>4%</div><div>51%35%7%7%</div></div>
1	E	507	<div><div>3%</div><div>54%33%6%7%</div></div>
1	F	507	<div><div>4%</div><div>51%34%8%7%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	507	<div><div>%</div><div><div></div><div>54%</div><div>32%</div><div>6%</div><div>7%</div></div></div>
1	H	507	<div><div>5%</div><div><div></div><div>51%</div><div>36%</div><div>7%</div><div>7%</div></div></div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 30845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysyl-tRNA synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	469	Total	C	N	O	S	0	3	0
			3848	2488	639	704	17			
1	B	471	Total	C	N	O	S	0	1	0
			3852	2490	640	705	17			
1	C	471	Total	C	N	O	S	0	0	0
			3844	2484	639	705	16			
1	D	471	Total	C	N	O	S	0	0	0
			3849	2488	640	705	16			
1	E	470	Total	C	N	O	S	0	0	0
			3833	2477	638	702	16			
1	F	470	Total	C	N	O	S	0	1	0
			3843	2485	638	704	16			
1	G	470	Total	C	N	O	S	0	0	0
			3840	2483	638	703	16			
1	H	471	Total	C	N	O	S	0	0	0
			3839	2481	639	703	16			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	10	Total	O	0	0
			10	10		
2	B	14	Total	O	0	0
			14	14		
2	C	13	Total	O	0	0
			13	13		
2	D	16	Total	O	0	0
			16	16		
2	E	9	Total	O	0	0
			9	9		
2	F	8	Total	O	0	0
			8	8		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	9	Total	O	0	0
			9	9		
2	H	18	Total	O	0	0
			18	18		


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Lysyl-tRNA synthetase




#### • Molecule 1: Lysyl-tRNA synthetase







• Molecule 1: Lysyl-tRNA synthetase




• Molecule 1: Lysyl-tRNA synthetase







• Molecule 1: Lysyl-tRNA synthetase




• Molecule 1: Lysyl-tRNA synthetase







• Molecule 1: Lysyl-tRNA synthetase



• Molecule 1: Lysyl-tRNA synthetase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.33Å 59.14Å 297.22Å 90.00° 97.47° 90.00°	Depositor
Resolution (Å)	48.70 – 2.90 48.70 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.1 (48.70-2.90) 93.4 (48.70-2.90)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.60 (at 2.91Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, $R_{free}$	0.226 , 0.298 0.219 , 0.295	Depositor DCC
$R_{free}$ test set	3113 reflections (3.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	46.8	Xtriage
Anisotropy	0.644	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 45.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.39$ , $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	30845	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/3954	0.63	0/5342
1	B	0.50	0/3952	0.65	0/5339
1	C	0.53	1/3941 (0.0%)	0.68	1/5326 (0.0%)
1	D	0.52	0/3946	0.64	0/5331
1	E	0.53	0/3929	0.65	0/5308
1	F	0.51	0/3943	0.61	0/5327
1	G	0.51	0/3937	0.65	1/5319 (0.0%)
1	H	0.53	0/3935	0.62	0/5316
All	All	0.51	1/31537 (0.0%)	0.64	2/42608 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	D	0	3
1	F	0	1
1	G	0	2
1	H	0	1
All	All	0	9

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	360	TRP	NE1-CE2	-5.37	1.30	1.37

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	483	ARG	NE-CZ-NH1	-5.41	117.59	120.30
1	C	360	TRP	CE2-CD2-CG	-5.12	103.20	107.30

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	416	GLU	Peptide
1	A	418	PRO	Peptide
1	D	227	LEU	Peptide
1	D	416	GLU	Peptide
1	D	418	PRO	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3848	0	3817	145	0
1	B	3852	0	3819	145	1
1	C	3844	0	3804	138	0
1	D	3849	0	3814	150	0
1	E	3833	0	3799	132	0
1	F	3843	0	3811	139	0
1	G	3840	0	3806	121	1
1	H	3839	0	3803	153	0
2	A	10	0	0	1	0
2	B	14	0	0	1	0
2	C	13	0	0	0	0
2	D	16	0	0	0	0
2	E	9	0	0	1	0
2	F	8	0	0	1	0
2	G	9	0	0	1	0
2	H	18	0	0	1	0
All	All	30845	0	30473	1071	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 1071 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:547:LEU:HD12	1:C:548:PRO:HD2	1.33	1.09
1:G:469:VAL:HG13	1:G:470:GLU:HG3	1.35	1.05
1:H:417:GLN:O	1:H:419:PHE:N	1.92	1.01
1:D:308:GLU:HG3	1:D:309:LEU:HD13	1.40	1.01
1:B:235:ARG:NH2	1:B:580:ARG:O	2.00	0.95

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:TYR:OH	1:G:434:GLU:OE2[1_465]	1.96	0.24

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	466/507 (92%)	425 (91%)	35 (8%)	6 (1%)	14	43
1	B	466/507 (92%)	414 (89%)	44 (9%)	8 (2%)	11	36
1	C	465/507 (92%)	408 (88%)	51 (11%)	6 (1%)	14	43
1	D	465/507 (92%)	410 (88%)	49 (10%)	6 (1%)	14	43
1	E	464/507 (92%)	417 (90%)	40 (9%)	7 (2%)	12	39
1	F	465/507 (92%)	420 (90%)	33 (7%)	12 (3%)	6	24
1	G	464/507 (92%)	419 (90%)	33 (7%)	12 (3%)	6	24
1	H	465/507 (92%)	419 (90%)	38 (8%)	8 (2%)	11	36
All	All	3720/4056 (92%)	3332 (90%)	323 (9%)	65 (2%)	11	36

5 of 65 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	437	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	416	GLU
1	B	418	PRO
1	B	423	GLU
1	B	458	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/457 (93%)	382 (90%)	45 (10%)	8	24
1	B	427/457 (93%)	365 (86%)	62 (14%)	4	11
1	C	425/457 (93%)	366 (86%)	59 (14%)	4	12
1	D	426/457 (93%)	363 (85%)	63 (15%)	3	11
1	E	424/457 (93%)	370 (87%)	54 (13%)	5	15
1	F	426/457 (93%)	364 (85%)	62 (15%)	3	11
1	G	425/457 (93%)	372 (88%)	53 (12%)	5	16
1	H	424/457 (93%)	370 (87%)	54 (13%)	5	15
All	All	3404/3656 (93%)	2952 (87%)	452 (13%)	4	13

5 of 452 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	372	LEU
1	E	232	ILE
1	H	228	LYS
1	D	415	LEU
1	E	81	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 40 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	172	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	417	GLN
1	H	162	GLN
1	E	339	ASN
1	F	86	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	469/507 (92%)	-0.09	6 (1%) 77 76	22, 46, 91, 122	0
1	B	471/507 (92%)	-0.05	9 (1%) 67 64	22, 47, 88, 118	0
1	C	471/507 (92%)	-0.01	9 (1%) 67 64	22, 48, 93, 129	0
1	D	471/507 (92%)	0.12	18 (3%) 41 35	22, 49, 91, 121	0
1	E	470/507 (92%)	-0.04	14 (2%) 51 44	26, 49, 92, 126	0
1	F	470/507 (92%)	0.09	22 (4%) 32 28	27, 51, 92, 118	0
1	G	470/507 (92%)	-0.07	5 (1%) 80 79	27, 49, 90, 123	0
1	H	471/507 (92%)	0.08	24 (5%) 29 24	26, 50, 91, 125	0
All	All	3763/4056 (92%)	0.00	107 (2%) 53 48	22, 49, 92, 129	0

The worst 5 of 107 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	145	SER	5.6
1	H	164	LEU	5.2
1	C	448	LEU	4.6
1	D	125	GLY	4.5
1	B	432	ILE	4.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.