



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 25, 2017 – 10:40 AM EDT

PDB ID : 5H00  
Title : The crystal structure of D-2-haloacid dehalogenase  
Authors : Wang, Y.; Xue, S.  
Deposited on : unknown  
Resolution : 2.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1044 (2.66-2.62)
Clashscore	112137	1092 (2.66-2.62)
Ramachandran outliers	110173	1077 (2.66-2.62)
Sidechain outliers	110143	1077 (2.66-2.62)
RSRZ outliers	101464	1047 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	<div> <div></div> <div>78%16% . .</div> </div>
1	B	301	<div> <div>2%</div> <div>83%12% . .</div> </div>
1	C	301	<div> <div></div> <div>85%10% . .</div> </div>
1	D	301	<div> <div>%</div> <div>81%14%5%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9561 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called (R)-2-haloacid dehalogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	S	0	0	0
			2317	1485	405	417	10			
1	B	292	Total	C	N	O	S	0	0	0
			2327	1491	408	418	10			
1	C	289	Total	C	N	O	S	0	0	0
			2300	1474	402	414	10			
1	D	286	Total	C	N	O	S	0	0	0
			2274	1457	396	411	10			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	98	Total	O	0	0
			98	98		
2	B	84	Total	O	0	0
			84	84		
2	C	92	Total	O	0	0
			92	92		
2	D	69	Total	O	0	0
			69	69		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: (R)-2-haloacid dehalogenase




- Molecule 1: (R)-2-haloacid dehalogenase




- Molecule 1: (R)-2-haloacid dehalogenase



- Molecule 1: (R)-2-haloacid dehalogenase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.72Å 109.40Å 138.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.24 – 2.64 45.24 – 2.64	Depositor EDS
% Data completeness (in resolution range)	97.8 (45.24-2.64) 97.9 (45.24-2.64)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.22 (at 2.65Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.193 , 0.251 0.192 , 0.252	Depositor DCC
$R_{free}$ test set	2124 reflections (5.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.3	Xtriage
Anisotropy	0.712	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 43.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9561	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.70 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.5642e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.50	0/2377	0.61	1/3229 (0.0%)
1	B	0.49	0/2388	0.57	0/3244
1	C	0.46	0/2360	0.55	0/3206
1	D	0.44	0/2333	0.54	0/3168
All	All	0.47	0/9458	0.57	1/12847 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	18	GLU	N-CA-C	6.46	128.44	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	228	ASP	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2317	0	2285	39	0
1	B	2327	0	2292	31	0
1	C	2300	0	2266	25	0
1	D	2274	0	2236	31	0
2	A	98	0	0	7	0
2	B	84	0	0	6	0
2	C	92	0	0	1	0
2	D	69	0	0	4	0
All	All	9561	0	9079	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:GLY:O	1:B:28:ARG:NH2	2.09	0.85
1:D:65:GLU:OE1	1:D:68:ARG:NH1	2.08	0.85
1:C:21:ILE:HG22	1:C:23:PRO:HD3	1.64	0.80
1:A:107:LEU:HD23	1:A:110:MET:HE3	1.63	0.80
1:C:284:MET:HE2	1:C:285:LEU:HG	1.66	0.77
1:A:16:VAL:HG22	1:A:17:CYS:H	1.49	0.77
1:B:188:GLN:OE1	1:B:191:ALA:HB3	1.88	0.73
1:C:48:TRP:CZ3	1:C:50:GLY:HA2	2.24	0.72
1:A:65:GLU:OE2	1:A:68:ARG:NH1	2.24	0.70
1:B:48:TRP:NE1	2:B:401:HOH:O	2.10	0.69
1:A:28:ARG:NH2	1:B:111:GLY:O	2.28	0.66
1:B:286:SER:OG	2:B:402:HOH:O	2.13	0.66
1:A:167:ILE:HD11	1:C:126:ILE:HA	1.77	0.65
1:B:126:ILE:HA	1:D:167:ILE:HD11	1.79	0.65
1:A:45:GLY:HA2	1:A:170:ILE:HD11	1.81	0.63
1:B:228:ASP:O	2:B:403:HOH:O	2.15	0.63
1:C:111:GLY:O	1:D:28:ARG:NH2	2.32	0.62
1:A:35:GLU:OE2	1:A:38:ARG:NH1	2.31	0.62
1:B:19:ALA:HB1	1:B:273:GLU:HG2	1.80	0.62
1:B:65:GLU:OE2	1:B:68:ARG:NH1	2.32	0.62
1:A:14:ARG:HG3	1:A:14:ARG:O	2.00	0.61
1:A:10:LEU:N	2:A:407:HOH:O	2.34	0.61
1:B:48:TRP:CZ3	1:B:50:GLY:HA2	2.35	0.61
1:C:113:SER:O	1:C:117:ILE:HD12	2.01	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:48:TRP:NE1	2:C:401:HOH:O	2.07	0.60
1:C:211:ARG:O	1:C:211:ARG:HD2	2.01	0.60
1:C:97:SER:HB3	1:C:231:ALA:HB1	1.83	0.59
1:B:169:GLN:OE1	1:D:169:GLN:NE2	2.34	0.59
1:C:105:ASP:OD1	1:C:108:ARG:NH2	2.36	0.59
1:B:130:GLY:HA3	2:B:401:HOH:O	2.04	0.58
1:A:206:TYR:HD1	1:A:209:MET:HE3	1.68	0.58
1:B:188:GLN:OE1	1:B:188:GLN:HA	2.04	0.57
1:B:255:ARG:NH1	1:B:260:ASP:OD2	2.38	0.56
1:D:196:THR:OG1	2:D:402:HOH:O	2.18	0.56
1:D:224:LYS:NZ	2:D:402:HOH:O	2.34	0.54
1:D:121:ARG:HH21	1:D:300:LEU:HB2	1.71	0.54
1:B:21:ILE:HG22	1:B:23:PRO:HD3	1.88	0.54
1:A:211:ARG:NH2	2:A:413:HOH:O	2.40	0.53
1:C:12:LEU:HD23	1:C:236:ARG:HG2	1.91	0.53
1:D:56:VAL:HG12	1:D:63:PHE:HB2	1.91	0.53
1:A:48:TRP:CZ3	1:A:50:GLY:HA2	2.44	0.52
1:A:52:ILE:HD13	1:A:281:PHE:HB3	1.91	0.52
1:D:69:ARG:HD2	1:D:256:MET:HE1	1.92	0.52
1:A:21:ILE:HG22	1:A:23:PRO:HD3	1.92	0.52
1:A:255:ARG:NH1	1:A:260:ASP:OD2	2.41	0.52
1:A:18:GLU:OE2	1:A:280:LEU:HB2	2.10	0.52
1:D:21:ILE:HG22	1:D:23:PRO:HD3	1.92	0.51
1:A:15:PRO:HB3	1:A:283:TRP:CE2	2.46	0.51
1:B:169:GLN:CD	1:D:169:GLN:HE22	2.13	0.50
1:B:236:ARG:HG3	1:B:286:SER:HB2	1.94	0.49
1:C:121:ARG:HH21	1:C:300:LEU:H	1.61	0.49
1:C:185:THR:HG23	1:D:185:THR:HG23	1.94	0.49
1:C:22:ARG:HD2	1:C:58:TYR:CE2	2.48	0.48
1:D:20:ILE:H	1:D:20:ILE:HD12	1.78	0.48
1:B:229:THR:O	1:B:233:GLN:HG3	2.13	0.48
1:C:185:THR:HG21	1:D:184:GLY:HA3	1.95	0.48
1:D:97:SER:HB3	1:D:231:ALA:HB1	1.96	0.48
1:B:300:LEU:O	2:B:405:HOH:O	2.20	0.47
1:D:69:ARG:HD2	1:D:256:MET:CE	2.44	0.47
1:B:211:ARG:NH2	2:B:416:HOH:O	2.47	0.47
1:D:65:GLU:HG3	1:D:256:MET:CE	2.45	0.47
1:B:53:THR:HG23	1:B:63:PHE:CE1	2.50	0.46
1:D:22:ARG:NH2	1:D:273:GLU:OE2	2.49	0.46
1:D:23:PRO:O	1:D:25:PRO:HD3	2.16	0.46
1:C:53:THR:HG23	1:C:63:PHE:CE1	2.50	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:ASP:O	1:A:242:ARG:HG2	2.16	0.46
1:A:205:ASP:O	1:A:209:MET:HE2	2.16	0.45
1:A:242:ARG:HD2	2:A:441:HOH:O	2.16	0.45
1:A:135:LEU:HA	1:A:285:LEU:HD13	1.99	0.45
1:B:169:GLN:HB2	1:D:169:GLN:NE2	2.31	0.45
1:C:52:ILE:HG23	1:C:53:THR:H	1.81	0.44
1:D:169:GLN:HB3	2:D:457:HOH:O	2.17	0.44
1:A:52:ILE:HD11	1:A:278:ILE:HD12	1.98	0.44
1:A:171:ASP:O	2:A:401:HOH:O	2.21	0.44
1:C:100:ILE:HD13	1:C:297:GLN:HB2	2.00	0.44
1:D:250:LEU:HA	1:D:251:PRO:HD3	1.81	0.44
1:A:202:ILE:HG22	1:A:207:LYS:HG3	1.99	0.44
1:B:101:GLU:OE1	1:B:101:GLU:HA	2.18	0.43
1:B:167:ILE:HD12	1:B:168:CYS:N	2.32	0.43
1:A:28:ARG:NE	2:A:421:HOH:O	2.46	0.43
1:A:28:ARG:NH1	2:A:424:HOH:O	2.51	0.43
1:B:166:PRO:HB2	1:D:125:ASP:HB3	1.99	0.43
1:A:31:GLN:HG2	1:A:31:GLN:O	2.19	0.43
1:B:12:LEU:HD12	1:B:13:PRO:HD2	2.01	0.43
1:B:45:GLY:HA3	1:B:167:ILE:HD11	2.00	0.43
1:A:56:VAL:HG12	1:A:63:PHE:HB2	2.01	0.43
1:A:16:VAL:HG11	1:A:279:SER:HB3	2.00	0.42
1:A:294:HIS:CE1	1:A:298:GLN:HE21	2.37	0.42
1:A:107:LEU:HD23	1:A:110:MET:CE	2.42	0.42
1:C:103:GLN:NE2	1:C:298:GLN:OE1	2.53	0.42
1:C:211:ARG:C	1:C:211:ARG:HD2	2.40	0.42
1:C:167:ILE:HA	1:C:167:ILE:HD12	1.89	0.41
1:C:18:GLU:HG2	1:C:18:GLU:H	1.64	0.41
1:D:259:ASP:OD2	2:D:403:HOH:O	2.21	0.41
1:B:250:LEU:HA	1:B:251:PRO:HD3	1.86	0.41
1:D:108:ARG:HH21	1:D:114:VAL:H	1.67	0.41
1:A:294:HIS:O	1:A:298:GLN:HG3	2.21	0.41
1:A:100:ILE:HD13	1:A:297:GLN:HB2	2.03	0.41
1:B:258:ARG:HB3	1:B:258:ARG:NH1	2.36	0.41
1:D:113:SER:O	1:D:117:ILE:HG13	2.20	0.41
1:D:89:ARG:HG2	1:D:242:ARG:NE	2.36	0.41
1:D:148:GLY:HA2	1:D:257:SER:HB3	2.02	0.41
1:A:11:GLN:C	1:A:12:LEU:HD23	2.40	0.41
1:D:65:GLU:HG3	1:D:256:MET:HE1	2.01	0.41
1:C:101:GLU:OE1	1:C:101:GLU:N	2.53	0.41
1:D:89:ARG:NH2	1:D:93:LEU:HD21	2.36	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ILE:HA	1:A:167:ILE:HD12	1.72	0.40
1:C:22:ARG:HD2	1:C:58:TYR:CZ	2.57	0.40
1:A:65:GLU:O	1:A:69:ARG:HG3	2.22	0.40
1:C:117:ILE:HG22	1:C:121:ARG:NH1	2.37	0.40
1:A:255:ARG:HD3	2:A:437:HOH:O	2.21	0.40
1:A:52:ILE:HD11	1:A:278:ILE:CD1	2.52	0.40
1:D:52:ILE:HG23	1:D:53:THR:H	1.86	0.40
1:B:120:ILE:HG12	1:B:212:TRP:CE3	2.57	0.40
1:B:52:ILE:HG23	1:B:53:THR:H	1.86	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/301 (96%)	276 (96%)	12 (4%)	1 (0%)	44	62
1	B	290/301 (96%)	283 (98%)	7 (2%)	0	100	100
1	C	287/301 (95%)	280 (98%)	7 (2%)	0	100	100
1	D	284/301 (94%)	274 (96%)	9 (3%)	1 (0%)	38	54
All	All	1150/1204 (96%)	1113 (97%)	35 (3%)	2 (0%)	51	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	ALA
1	D	156	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/249 (96%)	230 (96%)	9 (4%)	38	58
1	B	240/249 (96%)	232 (97%)	8 (3%)	43	64
1	C	237/249 (95%)	234 (99%)	3 (1%)	73	87
1	D	234/249 (94%)	229 (98%)	5 (2%)	59	78
All	All	950/996 (95%)	925 (97%)	25 (3%)	51	72

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	31	GLN
1	A	52	ILE
1	A	73	SER
1	A	165	SER
1	A	167	ILE
1	A	171	ASP
1	A	205	ASP
1	A	263	GLN
1	B	9	HIS
1	B	52	ILE
1	B	168	CYS
1	B	185	THR
1	B	188	GLN
1	B	205	ASP
1	B	211	ARG
1	B	286	SER
1	C	52	ILE
1	C	205	ASP
1	C	284	MET
1	D	18	GLU
1	D	48	TRP
1	D	52	ILE
1	D	205	ASP
1	D	211	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	291/301 (96%)	-0.46	0 <b>100</b> <b>100</b>	13, 22, 45, 59	0
1	B	292/301 (97%)	-0.37	6 (2%) 64 60	14, 24, 44, 58	0
1	C	289/301 (96%)	-0.45	1 (0%) <b>93</b> <b>94</b>	16, 27, 51, 65	0
1	D	286/301 (95%)	-0.25	4 (1%) 75 72	18, 30, 53, 80	0
All	All	1158/1204 (96%)	-0.38	11 (0%) <b>84</b> <b>83</b>	13, 26, 49, 80	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	16	VAL	5.2
1	B	100	ILE	3.3
1	B	101	GLU	3.2
1	D	19	ALA	3.0
1	C	101	GLU	2.7
1	D	18	GLU	2.3
1	B	99	VAL	2.3
1	B	300	LEU	2.1
1	B	222	ALA	2.1
1	B	10	LEU	2.0
1	D	15	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.