



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2017 – 10:33 AM EDT

PDB ID : 5H0I
Title : Name to be released when published
Authors : Yang, R.; Wong, Y.H.; Lescar, J.; Wu, B.
Deposited on : 2016-10-04
Resolution : 2.56 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029077
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029077


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3689 (2.60-2.52)
Clashscore	112137	4096 (2.60-2.52)
Ramachandran outliers	110173	4037 (2.60-2.52)
Sidechain outliers	110143	4037 (2.60-2.52)
RSRZ outliers	101464	3700 (2.60-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div> <div>6%</div> <div> <div></div> <div>77%</div> <div>10%</div> <div>•</div> <div>11%</div> </div> </div>
1	B	448	<div> <div>6%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>•</div> <div>11%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Asparaginyl endopeptidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	400	Total	C	N	O	S	0	0	0
			3101	1954	531	599	17			
1	B	400	Total	C	N	O	S	0	0	0
			3101	1956	530	598	17			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	138	Total	O	0	0
			138	138		
2	B	154	Total	O	0	0
			154	154		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Asparaginyl endopeptidase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	145.73Å 70.21Å 118.28Å 90.00° 117.14° 90.00°	Depositor
Resolution (Å)	71.72 – 2.56 71.72 – 2.56	Depositor EDS
% Data completeness (in resolution range)	99.6 (71.72-2.56) 99.6 (71.72-2.56)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.55Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.186 , 0.224 0.194 , 0.224	Depositor DCC
R_{free} test set	530 reflections (1.56%)	DCC
Wilson B-factor (Å ²)	50.5	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 69.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6494	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/3176	0.73	0/4309
1	B	0.52	0/3175	0.73	0/4307
All	All	0.52	0/6351	0.73	0/8616

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3101	0	2970	27	0
1	B	3101	0	2977	26	0
2	A	138	0	0	0	0
2	B	154	0	0	0	0
All	All	6494	0	5947	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:174:ASP:CG	1:B:175:HIS:H	1.53	1.11

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:174:ASP:CG	1:A:175:HIS:H	1.49	1.08
1:A:174:ASP:CG	1:A:175:HIS:N	2.26	0.87
1:B:174:ASP:CG	1:B:175:HIS:N	2.30	0.81
1:A:394:GLY:HA3	1:A:407:LEU:HD11	1.70	0.73
1:B:140:ASN:HD21	1:B:153:THR:H	1.37	0.73
1:A:140:ASN:HD21	1:A:153:THR:H	1.38	0.70
1:A:246:TRP:CD1	1:A:301:ALA:HB1	2.29	0.68
1:B:246:TRP:CD1	1:B:301:ALA:HB1	2.30	0.67
1:B:78:CYS:HB2	1:B:123:VAL:HG13	1.79	0.65
1:B:251:PRO:HD2	1:B:353:LEU:HD21	1.79	0.64
1:A:438:LEU:HA	1:A:442:GLY:HA3	1.82	0.61
1:A:345:VAL:HG21	1:A:353:LEU:HD13	1.83	0.60
1:A:241:THR:HG22	1:A:242:THR:HG23	1.86	0.57
1:A:65:SER:HB3	1:A:174:ASP:OD1	2.06	0.55
1:A:441:TYR:O	1:A:444:ARG:HG2	2.07	0.55
1:B:78:CYS:CB	1:B:123:VAL:HG13	2.38	0.53
1:B:281:ASN:HB3	1:B:284:TYR:HB2	1.89	0.53
1:A:306:GLN:HB3	1:A:310:LEU:HD22	1.91	0.53
1:A:281:ASN:HB3	1:A:284:TYR:HB2	1.90	0.53
1:B:397:LEU:HD13	1:B:465:ALA:HB2	1.91	0.52
1:A:407:LEU:HB3	1:A:448:THR:HG23	1.92	0.52
1:B:191:ALA:HB1	1:B:226:ILE:HD13	1.93	0.51
1:A:393:ILE:O	1:A:397:LEU:HB2	2.11	0.51
1:A:397:LEU:HD13	1:A:465:ALA:HB2	1.92	0.51
1:A:281:ASN:HB3	1:A:284:TYR:CB	2.42	0.50
1:A:397:LEU:HD21	1:A:457:ILE:HG21	1.94	0.50
1:B:281:ASN:HB3	1:B:284:TYR:CB	2.42	0.50
1:B:393:ILE:O	1:B:397:LEU:HB2	2.12	0.49
1:A:253:GLN:HG3	1:A:344:VAL:HA	1.94	0.49
1:A:74:GLN:OE1	1:A:99:TYR:O	2.31	0.49
1:B:397:LEU:HD21	1:B:457:ILE:HG21	1.95	0.49
1:B:447:ARG:HG2	1:B:447:ARG:HH11	1.80	0.47
1:A:439:SER:O	1:A:440:GLU:HB2	2.16	0.45
1:A:235:ALA:HB3	1:A:307:TYR:HB2	1.99	0.45
1:B:235:ALA:HB3	1:B:307:TYR:HB2	2.00	0.44
1:A:247:CYS:HA	1:A:267:ASP:HA	2.00	0.44
1:A:86:ARG:HH11	1:A:324:ASN:ND2	2.17	0.43
1:B:438:LEU:HA	1:B:442:GLY:HA3	2.00	0.43
1:B:243:GLU:OE1	1:B:302:SER:HA	2.19	0.42
1:B:282:SER:HB3	1:B:318:PHE:HB3	2.00	0.42
1:B:65:SER:HB3	1:B:174:ASP:OD1	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:118:PRO:HD3	1:B:249:TYR:CG	2.54	0.42
1:A:118:PRO:HD3	1:A:249:TYR:CD1	2.55	0.42
1:A:253:GLN:HG2	1:A:254:GLU:H	1.85	0.42
1:A:438:LEU:HD12	1:A:442:GLY:HA3	2.01	0.42
1:A:282:SER:HB3	1:A:318:PHE:HB3	2.00	0.41
1:B:191:ALA:HB3	1:B:421:TRP:CH2	2.55	0.41
1:B:286:THR:HA	1:B:316:GLY:HA2	2.01	0.41
1:B:110:ARG:HD3	1:B:261:TYR:OH	2.21	0.41
1:B:345:VAL:HG21	1:B:353:LEU:HD13	2.03	0.40
1:B:59:ALA:HA	1:B:169:PHE:O	2.21	0.40
1:B:181:ILE:HG21	1:B:214:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/448 (88%)	375 (95%)	17 (4%)	4 (1%)	18	34
1	B	396/448 (88%)	376 (95%)	17 (4%)	3 (1%)	22	41
All	All	792/896 (88%)	751 (95%)	34 (4%)	7 (1%)	20	37

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	174	ASP
1	A	254	GLU
1	A	401	GLU
1	B	174	ASP
1	A	440	GLU
1	B	254	GLU
1	B	440	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/372 (88%)	313 (95%)	16 (5%)	29	50
1	B	329/372 (88%)	316 (96%)	13 (4%)	36	60
All	All	658/744 (88%)	629 (96%)	29 (4%)	33	56

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	98	MET
1	A	212	PHE
1	A	241	THR
1	A	253	GLN
1	A	268	LEU
1	A	299	SER
1	A	312	LEU
1	A	344	VAL
1	A	353	LEU
1	A	364	GLU
1	A	392	LEU
1	A	400	ILE
1	A	418	VAL
1	A	437	SER
1	A	438	LEU
1	A	439	SER
1	B	98	MET
1	B	123	VAL
1	B	212	PHE
1	B	253	GLN
1	B	299	SER
1	B	310	LEU
1	B	353	LEU
1	B	400	ILE
1	B	402	LYS
1	B	403	CYS
1	B	405	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	408	ASN
1	B	418	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	140	ASN
1	A	202	HIS
1	A	293	HIS
1	A	324	ASN
1	B	140	ASN
1	B	290	GLN
1	B	293	HIS
1	B	303	HIS
1	B	408	ASN
1	B	461	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	400/448 (89%)	0.56	26 (6%)	20 17	34, 70, 117, 144	0
1	B	400/448 (89%)	0.59	29 (7%)	16 14	31, 60, 116, 162	0
All	All	800/896 (89%)	0.58	55 (6%)	18 15	31, 64, 116, 162	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	282	SER	8.5
1	B	343	ILE	8.2
1	B	402	LYS	7.7
1	B	284	TYR	7.5
1	B	472	SER	7.4
1	A	252	ALA	7.2
1	A	403	CYS	7.1
1	A	283	TRP	6.7
1	B	281	ASN	6.6
1	B	407	LEU	6.1
1	B	403	CYS	6.1
1	B	283	TRP	5.9
1	B	471	ALA	5.5
1	A	284	TYR	4.8
1	B	253	GLN	4.0
1	B	438	LEU	3.9
1	A	396	LEU	3.9
1	A	317	LEU	3.8
1	A	471	ALA	3.2
1	A	356	TRP	3.2
1	B	386	ILE	3.0
1	B	436	GLY	2.9
1	B	469	ALA	2.8
1	A	106	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	324	ASN	2.8
1	B	462	MET	2.7
1	A	253	GLN	2.6
1	A	365	GLY	2.6
1	B	392	LEU	2.6
1	A	438	LEU	2.6
1	B	246	TRP	2.6
1	A	468	GLN	2.5
1	A	434	HIS	2.4
1	A	246	TRP	2.4
1	B	468	GLN	2.4
1	A	248	TYR	2.4
1	B	255	ASN	2.4
1	B	252	ALA	2.4
1	A	318	PHE	2.3
1	A	407	LEU	2.3
1	A	390	ILE	2.3
1	A	398	PHE	2.2
1	B	350	ALA	2.2
1	B	410	VAL	2.2
1	A	392	LEU	2.2
1	B	390	ILE	2.1
1	B	470	CYS	2.1
1	B	356	TRP	2.1
1	A	279	VAL	2.1
1	A	231	LEU	2.1
1	B	396	LEU	2.0
1	A	256	PRO	2.0
1	B	344	VAL	2.0
1	B	249	TYR	2.0
1	A	402	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.