



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 23, 2018 – 01:50 PM EST

PDB ID : 5H2P  
Title : A three dimensional movie of structural changes in bacteriorhodopsin: structure obtained 657 us after photoexcitation  
Authors : Royant, A.; Nango, E.; Nakane, T.; Tanaka, T.; Arima, T.; Neutze, R.; Iwata, S.  
Deposited on : 2016-10-15  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	<div> <div>19%</div> <div> <div></div> <div>88%</div> <div>5%</div> <div>7%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
11	C14	A	314	-	-	-	X
3	L2P	A	302	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	L2P	A	310	-	-	-	X
3	L2P	A	311	-	-	-	X
4	TRD	A	303	-	-	-	X
7	OCT	A	307	-	-	-	X
8	MYS	A	312	-	-	-	X

## 2 Entry composition [i](#)


There are 12 unique types of molecules in this entry. The entry contains 5552 atoms, of which 2935 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bacteriorhodopsin.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	231	4751	1564	2420	359	396	12	0	69	0

- Molecule 2 is RETINAL (three-letter code: RET) (formula: C<sub>20</sub>H<sub>28</sub>O).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	H		
2	A	1	96	40	56	0	1

- Molecule 3 is 2,3-DI-PHYTANYL-GLYCEROL (three-letter code: L2P) (formula: C<sub>43</sub>H<sub>88</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			71	23	46	2		
3	A	1	Total	C	H	O	0	0
			65	21	43	1		
3	A	1	Total	C	H	O	0	0
			134	43	88	3		

- Molecule 4 is TRIDECANE (three-letter code: TRD) (formula: C<sub>13</sub>H<sub>28</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	H	0	0
			41	13	28		

- Molecule 5 is DECANE (three-letter code: D10) (formula:  $C_{10}H_{22}$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	H	0	0
			32	10	22		

- Molecule 6 is HEPTANE (three-letter code: HP6) (formula:  $C_7H_{16}$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	H	0	0
			23	7	16		

- Molecule 7 is N-OCTANE (three-letter code: OCT) (formula:  $C_8H_{18}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	H	0	0
			26	8	18		
7	A	1	Total	C	H	0	0
			26	8	18		
7	A	1	Total	C	H	0	0
			26	8	18		

- Molecule 8 is PENTADECANE (three-letter code: MYS) (formula: C<sub>15</sub>H<sub>32</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	H	0	0
			47	15	32		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	H	0	0
			47	15	32		


- Molecule 9 is UNDECANE (three-letter code: UND) (formula:  $C_{11}H_{24}$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	A	1	Total	C	H	0	0
			35	11	24		
9	A	1	Total	C	H	0	0
			35	11	24		


- Molecule 10 is nonane (three-letter code: DD9) (formula:  $C_9H_{20}$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	H	0	0
			29	9	20		

- Molecule 11 is TETRADECANE (three-letter code: C14) (formula: C<sub>14</sub>H<sub>30</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	H	0	0
			44	14	30		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	22	Total 24	O 24	0	12

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Bacteriorhodopsin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.50 Å 62.50 Å 112.00 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	21.91 – 2.10 21.91 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (21.91-2.10) 100.0 (21.91-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.95 (at 2.09 Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.160 , 0.178 0.157 , 0.177	Depositor DCC
$R_{free}$ test set	719 reflections (4.97%)	DCC
Wilson B-factor (Å <sup>2</sup> )	44.7	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 91.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.073 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5552	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DD9, D10, MYS, HP6, RET, TRD, L2P, UND, OCT, C14

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/2391	0.43	0/3267

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2331	2420	2415	8	0
2	A	40	56	54	11	0
3	A	93	177	172	3	0
4	A	13	28	28	0	0
5	A	10	22	22	0	0
6	A	7	16	16	0	0
7	A	24	54	54	0	0
8	A	30	64	64	0	0
9	A	22	48	48	0	0
10	A	9	20	20	0	0
11	A	14	30	30	0	0
12	A	24	0	0	1	0
All	All	2617	2935	2923	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:301[B]:RET:H8	2:A:301[B]:RET:H161	1.71	0.73
2:A:301[A]:RET:H8	2:A:301[A]:RET:H171	1.81	0.62
3:A:302:L2P:H552	3:A:302:L2P:H591	1.86	0.58
2:A:301[B]:RET:C8	2:A:301[B]:RET:H161	2.39	0.53
1:A:185:TYR:HB3	2:A:301[B]:RET:H171	1.92	0.51
1:A:83[B]:TYR:OH	1:A:189[B]:TRP:NE1	2.43	0.51
1:A:82[B]:ARG:NH2	12:A:401[B]:HOH:O	2.40	0.48
1:A:186:PRO:HB3	2:A:301[B]:RET:H183	1.94	0.48
2:A:301[B]:RET:H7	2:A:301[B]:RET:H181	1.66	0.47
2:A:301[B]:RET:H171	2:A:301[B]:RET:C8	2.45	0.46
1:A:186:PRO:HB3	2:A:301[A]:RET:H183	1.98	0.46
2:A:301[A]:RET:H171	2:A:301[A]:RET:C8	2.44	0.45
2:A:301[A]:RET:H8	2:A:301[A]:RET:H161	1.98	0.45
1:A:156:PHE:HB3	1:A:171:PHE:CZ	2.52	0.45
1:A:153:PHE:CE2	1:A:179[B]:VAL:HG21	2.53	0.43
3:A:311:L2P:H172	3:A:311:L2P:H592	2.02	0.42
1:A:90[A]:THR:N	1:A:91:PRO:HD2	2.35	0.42
2:A:301[B]:RET:H171	2:A:301[B]:RET:H8	2.03	0.41
3:A:311:L2P:H541	3:A:311:L2P:H151	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	298/248 (120%)	291 (98%)	6 (2%)	1 (0%)	44 44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/194 (126%)	244 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	RET	A	301[A]	1	19,20,21	0.66	0	27,27,28	1.70	9 (33%)
2	RET	A	301[B]	1	19,20,21	0.74	0	27,27,28	2.07	8 (29%)
3	L2P	A	302	-	24,24,45	0.79	0	26,27,53	0.79	1 (3%)
4	TRD	A	303	-	12,12,12	0.30	0	11,11,11	0.79	0
5	D10	A	304	-	9,9,9	0.31	0	8,8,8	0.68	0
6	HP6	A	305	-	6,6,6	0.31	0	5,5,5	0.63	0
7	OCT	A	306	-	7,7,7	0.27	0	6,6,6	0.76	0
7	OCT	A	307	-	7,7,7	0.30	0	6,6,6	0.72	0
8	MYS	A	308	-	14,14,14	0.30	0	13,13,13	0.77	0
9	UND	A	309	-	10,10,10	0.30	0	9,9,9	0.77	0
3	L2P	A	310	-	21,21,45	0.67	0	23,24,53	0.83	0
3	L2P	A	311	-	45,45,45	0.76	0	49,53,53	0.78	2 (4%)
8	MYS	A	312	-	14,14,14	0.30	0	13,13,13	0.81	0
10	DD9	A	313	-	8,8,8	0.30	0	7,7,7	0.75	0
11	C14	A	314	-	13,13,13	0.30	0	12,12,12	0.80	0
7	OCT	A	315	-	7,7,7	0.31	0	6,6,6	0.64	0
9	UND	A	316	-	10,10,10	0.31	0	9,9,9	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RET	A	301[A]	1	-	0/13/30/31	0/1/1/1
2	RET	A	301[B]	1	-	0/13/30/31	0/1/1/1
3	L2P	A	302	-	-	0/25/25/51	0/0/0/0
4	TRD	A	303	-	-	0/10/10/10	0/0/0/0
5	D10	A	304	-	-	0/7/7/7	0/0/0/0
6	HP6	A	305	-	-	0/4/4/4	0/0/0/0
7	OCT	A	306	-	-	0/5/5/5	0/0/0/0
7	OCT	A	307	-	-	0/5/5/5	0/0/0/0
8	MYS	A	308	-	-	0/12/12/12	0/0/0/0
9	UND	A	309	-	-	0/8/8/8	0/0/0/0
3	L2P	A	310	-	-	0/22/22/51	0/0/0/0
3	L2P	A	311	-	-	0/51/51/51	0/0/0/0
8	MYS	A	312	-	-	0/12/12/12	0/0/0/0
10	DD9	A	313	-	-	0/6/6/6	0/0/0/0
11	C14	A	314	-	-	0/11/11/11	0/0/0/0
7	OCT	A	315	-	-	0/5/5/5	0/0/0/0
9	UND	A	316	-	-	0/8/8/8	0/0/0/0



There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	A	301[B]	RET	C17-C1-C6	-5.97	100.63	110.31
2	A	301[A]	RET	C3-C4-C5	-3.01	108.60	113.78
2	A	301[B]	RET	C11-C10-C9	-2.92	123.14	127.31
2	A	301[A]	RET	C10-C11-C12	-2.68	115.01	123.23
2	A	301[B]	RET	C18-C5-C6	-2.67	121.52	124.51
2	A	301[B]	RET	C7-C8-C9	-2.28	122.79	126.21
2	A	301[A]	RET	C8-C9-C10	-2.16	115.63	118.94
2	A	301[B]	RET	C20-C13-C14	-2.15	117.21	123.66
2	A	301[A]	RET	C11-C10-C9	-2.14	124.25	127.31
3	A	311	L2P	C26-C25-C23	-2.12	108.78	115.73
3	A	302	L2P	C51-C52-C53	-2.06	108.97	115.73
2	A	301[A]	RET	C2-C3-C4	-2.03	106.52	111.34
2	A	301[A]	RET	C18-C5-C6	-2.02	122.25	124.51
2	A	301[A]	RET	C19-C9-C10	-2.01	120.11	122.92
3	A	311	L2P	C16-C15-C13	-2.00	109.17	115.73
2	A	301[B]	RET	C1-C6-C7	2.00	121.36	115.73
2	A	301[A]	RET	C1-C6-C7	2.02	121.41	115.73
2	A	301[B]	RET	C19-C9-C8	2.81	122.57	118.10
2	A	301[B]	RET	C2-C1-C6	3.60	116.11	110.48
2	A	301[A]	RET	C19-C9-C8	3.87	124.27	118.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301[A]	RET	4	0
2	A	301[B]	RET	7	0
3	A	302	L2P	1	0
3	A	311	L2P	2	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	231/248 (93%)	1.14	47 (20%) <b>1</b> <b>1</b>	35, 45, 84, 116	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	162	SER	6.0
1	A	5	THR	5.8
1	A	4	ILE	5.7
1	A	74	GLU	5.7
1	A	158	SER	5.4
1	A	73	GLY	5.0
1	A	159	LYS	4.8
1	A	227[A]	ARG	4.7
1	A	35	SER	4.5
1	A	6	GLY	4.1
1	A	161	GLU	4.1
1	A	233	ALA	4.1
1	A	33	GLY	3.8
1	A	34	VAL	3.6
1	A	82[A]	ARG	3.4
1	A	157	THR	3.3
1	A	97	LEU	3.2
1	A	208[A]	PHE	3.2
1	A	163	MET	3.1
1	A	86[A]	TRP	3.1
1	A	198[A]	ILE	3.0
1	A	160	ALA	2.9
1	A	122	GLY	2.9
1	A	189[A]	TRP	2.9
1	A	36	ASP	2.8
1	A	37	PRO	2.7
1	A	72	GLY	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	10	TRP	2.7
1	A	83[A]	TYR	2.6
1	A	70	PRO	2.6
1	A	32	MET	2.5
1	A	154	PHE	2.5
1	A	223[A]	LEU	2.5
1	A	153	PHE	2.4
1	A	69	VAL	2.4
1	A	81[A]	ALA	2.3
1	A	100	LEU	2.3
1	A	165	PRO	2.3
1	A	93[A]	LEU	2.3
1	A	199[A]	VAL	2.3
1	A	156	PHE	2.2
1	A	75	GLN	2.1
1	A	94[A]	LEU	2.1
1	A	71	PHE	2.1
1	A	79	TYR	2.1
1	A	53	ALA	2.0
1	A	96	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	L2P	A	310	22/46	0.64	0.26	7.42	73,94,115,124	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	L2P	A	302	25/46	0.77	0.28	4.28	59,81,104,138	0
3	L2P	A	311	46/46	0.69	0.28	3.24	60,93,123,127	0
11	C14	A	314	14/14	0.53	0.23	2.35	65,85,110,115	0
7	OCT	A	307	8/8	0.69	0.23	2.26	67,94,105,106	0
8	MYS	A	312	15/15	0.59	0.23	2.21	70,93,106,106	0
4	TRD	A	303	13/13	0.44	0.24	2.09	71,91,118,121	0
9	UND	A	309	11/11	0.67	0.17	2.00	64,83,100,101	0
7	OCT	A	306	8/8	0.71	0.26	1.86	67,84,93,93	0
2	RET	A	301[A]	20/21	0.88	0.28	1.76	34,42,52,66	48
2	RET	A	301[B]	20/21	0.88	0.28	1.73	34,42,52,66	48
5	D10	A	304	10/10	0.73	0.20	1.72	69,92,103,109	0
9	UND	A	316	11/11	0.55	0.20	0.91	70,92,105,115	0
8	MYS	A	308	15/15	0.79	0.20	0.86	75,99,111,125	0
10	DD9	A	313	9/9	0.65	0.17	0.32	77,95,108,110	0
6	HP6	A	305	7/7	0.75	0.16	-0.15	63,82,89,89	0
7	OCT	A	315	8/8	0.53	0.27	-	70,94,110,113	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.