



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 26, 2017 – 12:13 PM EDT

PDB ID : 3H4Z
Title : Crystal Structure of an MBP-Der p 7 fusion protein
Authors : Pedersen, L.C.; Mueller, G.A.; London, R.E.
Deposited on : unknown
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1522 (2.38-2.34)
Clashscore	112137	1626 (2.38-2.34)
Ramachandran outliers	110173	1605 (2.38-2.34)
Sidechain outliers	110143	1606 (2.38-2.34)
RSRZ outliers	101464	1528 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	568	<div> <div>6%</div> <div>70%29%</div> <div>.</div> </div>
1	B	568	<div> <div>7%</div> <div>69%29%</div> <div>..</div> </div>
1	C	568	<div> <div>14%</div> <div>54%42%</div> <div>..</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12999 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein fused with Allergen DERP7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	568	Total	C	N	O	S	0	0	0
			4257	2721	696	828	12			
1	B	563	Total	C	N	O	S	0	0	0
			4220	2700	691	817	12			
1	C	552	Total	C	N	O	S	0	0	0
			4130	2642	674	803	11			

There are 45 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	82	ALA	ASP	ENGINEERED	UNP P0AEX9
A	83	ALA	LYS	ENGINEERED	UNP P0AEX9
A	172	ALA	GLU	ENGINEERED	UNP P0AEX9
A	173	ALA	ASN	ENGINEERED	UNP P0AEX9
A	239	ALA	LYS	ENGINEERED	UNP P0AEX9
A	359	ALA	GLU	ENGINEERED	UNP P0AEX9
A	362	ALA	-	SEE REMARK 999	UNP P0AEX9
A	363	ALA	-	SEE REMARK 999	UNP P0AEX9
A	364	ALA	-	SEE REMARK 999	UNP P0AEX9
A	365	GLN	-	SEE REMARK 999	UNP P0AEX9
A	366	THR	-	SEE REMARK 999	UNP P0AEX9
A	367	ASN	-	SEE REMARK 999	UNP P0AEX9
A	368	ALA	-	SEE REMARK 999	UNP P0AEX9
A	369	ALA	-	SEE REMARK 999	UNP P0AEX9
A	370	ALA	-	SEE REMARK 999	UNP P0AEX9
B	82	ALA	ASP	ENGINEERED	UNP P0AEX9
B	83	ALA	LYS	ENGINEERED	UNP P0AEX9
B	172	ALA	GLU	ENGINEERED	UNP P0AEX9
B	173	ALA	ASN	ENGINEERED	UNP P0AEX9
B	239	ALA	LYS	ENGINEERED	UNP P0AEX9
B	359	ALA	GLU	ENGINEERED	UNP P0AEX9
B	362	ALA	-	SEE REMARK 999	UNP P0AEX9
B	363	ALA	-	SEE REMARK 999	UNP P0AEX9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	364	ALA	-	SEE REMARK 999	UNP P0AEX9
B	365	GLN	-	SEE REMARK 999	UNP P0AEX9
B	366	THR	-	SEE REMARK 999	UNP P0AEX9
B	367	ASN	-	SEE REMARK 999	UNP P0AEX9
B	368	ALA	-	SEE REMARK 999	UNP P0AEX9
B	369	ALA	-	SEE REMARK 999	UNP P0AEX9
B	370	ALA	-	SEE REMARK 999	UNP P0AEX9
C	82	ALA	ASP	ENGINEERED	UNP P0AEX9
C	83	ALA	LYS	ENGINEERED	UNP P0AEX9
C	172	ALA	GLU	ENGINEERED	UNP P0AEX9
C	173	ALA	ASN	ENGINEERED	UNP P0AEX9
C	239	ALA	LYS	ENGINEERED	UNP P0AEX9
C	359	ALA	GLU	ENGINEERED	UNP P0AEX9
C	362	ALA	-	SEE REMARK 999	UNP P0AEX9
C	363	ALA	-	SEE REMARK 999	UNP P0AEX9
C	364	ALA	-	SEE REMARK 999	UNP P0AEX9
C	365	GLN	-	SEE REMARK 999	UNP P0AEX9
C	366	THR	-	SEE REMARK 999	UNP P0AEX9
C	367	ASN	-	SEE REMARK 999	UNP P0AEX9
C	368	ALA	-	SEE REMARK 999	UNP P0AEX9
C	369	ALA	-	SEE REMARK 999	UNP P0AEX9
C	370	ALA	-	SEE REMARK 999	UNP P0AEX9

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			11	6	5		
2	C	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	146	Total	O	0	0
			146	146		
4	B	130	Total	O	0	0
			130	130		
4	C	46	Total	O	0	0
			46	46		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein fused with Allergen DERP7




- Molecule 1: Maltose-binding periplasmic protein fused with Allergen DERP7





● Molecule 1: Maltose-binding periplasmic protein fused with Allergen DERP7



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	193.35Å 117.90Å 92.44Å 90.00° 114.06° 90.00°	Depositor
Resolution (Å)	28.91 – 2.35 28.91 – 2.35	Depositor EDS
% Data completeness (in resolution range)	77.3 (28.91-2.35) 77.3 (28.91-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.19 (at 2.36Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.249 , 0.293 0.252 , 0.294	Depositor DCC
R_{free} test set	3079 reflections (5.05%)	DCC
Wilson B-factor (Å ²)	41.3	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12999	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/4347	0.52	0/5925
1	B	0.27	0/4311	0.49	0/5877
1	C	0.27	0/4220	0.48	0/5759
All	All	0.27	0/12878	0.50	0/17561

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4257	0	4092	124	0
1	B	4220	0	4047	135	0
1	C	4130	0	3936	203	1
2	A	23	0	21	0	0
2	B	23	0	21	0	0
2	C	23	0	21	3	0
3	A	1	0	0	0	0
4	A	146	0	0	8	0
4	B	130	0	0	7	0
4	C	46	0	0	3	0
All	All	12999	0	12138	454	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 454 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1020:ILE:HG22	1:C:1021:HIS:H	1.27	0.95
1:C:139:LEU:HB3	1:C:144:LYS:HB2	1.53	0.91
1:B:1058:PHE:HB3	1:B:1185:VAL:HG21	1.56	0.87
1:A:9:ILE:HD11	1:A:61:PHE:CE1	2.10	0.87
1:A:1144:LEU:HD21	1:A:1210:LEU:HD13	1.57	0.86

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:141:ALA:O	1:C:141:ALA:O[2_555]	1.89	0.31

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	566/568 (100%)	524 (93%)	40 (7%)	2 (0%)	38	44
1	B	559/568 (98%)	504 (90%)	48 (9%)	7 (1%)	14	12
1	C	546/568 (96%)	489 (90%)	48 (9%)	9 (2%)	11	9
All	All	1671/1704 (98%)	1517 (91%)	136 (8%)	18 (1%)	17	16

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	146	ALA
1	C	186	ALA
1	C	1108	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1020	ILE
1	B	1093	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/461 (92%)	422 (99%)	4 (1%)	82	90
1	B	421/461 (91%)	419 (100%)	2 (0%)	91	95
1	C	412/461 (89%)	410 (100%)	2 (0%)	91	95
All	All	1259/1383 (91%)	1251 (99%)	8 (1%)	89	94

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1190	PHE
1	C	1190	PHE
1	B	1176	ASP
1	A	358	ASP
1	B	258	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	365	GLN
1	B	1054	HIS
1	C	282	ASN
1	B	241	ASN
1	C	1030	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	A	671	2	11,11,12	0.46	0	13,15,17	0.66	0
2	GLC	A	672	2	12,12,12	0.37	0	17,17,17	0.40	0
2	GLC	B	673	2	11,11,12	0.44	0	13,15,17	0.69	1 (7%)
2	GLC	B	674	2	12,12,12	0.40	0	17,17,17	0.35	0
2	GLC	C	675	2	11,11,12	0.41	0	13,15,17	0.63	0
2	GLC	C	676	2	12,12,12	0.35	0	17,17,17	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	671	2	-	0/2/19/22	0/1/1/1
2	GLC	A	672	2	-	0/2/22/22	0/1/1/1
2	GLC	B	673	2	-	0/2/19/22	0/1/1/1
2	GLC	B	674	2	-	0/2/22/22	0/1/1/1
2	GLC	C	675	2	-	0/2/19/22	0/1/1/1
2	GLC	C	676	2	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	673	GLC	C1-O5-C5	2.02	114.95	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	675	GLC	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	568/568 (100%)	0.40	34 (5%)	23 32	19, 42, 75, 90	0
1	B	563/568 (99%)	0.55	42 (7%)	15 22	22, 47, 73, 96	0
1	C	552/568 (97%)	1.00	77 (13%)	3 5	35, 61, 81, 96	0
All	All	1683/1704 (98%)	0.65	153 (9%)	10 15	19, 50, 78, 96	0

The worst 5 of 153 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1124	ASP	7.4
1	C	31	THR	7.0
1	C	1069	GLY	6.9
1	C	33	ILE	5.9
1	A	1123	GLY	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GLC	C	675	11/12	0.94	0.26	0.78	40,43,46,46	0
2	GLC	A	672	12/12	0.96	0.24	0.71	26,31,34,37	0
2	GLC	A	671	11/12	0.95	0.23	0.50	27,31,36,37	0
2	GLC	C	676	12/12	0.92	0.21	-0.21	44,48,50,51	0
2	GLC	B	674	12/12	0.92	0.17	-0.29	31,33,36,36	0
2	GLC	B	673	11/12	0.96	0.17	-0.31	35,38,39,40	0
3	NA	A	1216	1/1	0.89	0.21	-	46,46,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.