



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 05:59 pm GMT

PDB ID : 3H59  
Title : Hepatitis C virus polymerase NS5B with thiazine inhibitor 2  
Authors : Harris, S.F.; Ghatge, M.  
Deposited on : 2009-04-21  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	576	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 88%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>88%</span> <span>9% . .</span> </div> </div>
1	B	576	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 88%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>88%</span> <span>8% . .</span> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9582 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	557	Total	C	N	O	S	0	7	0
			4387	2767	775	810	35			
1	B	557	Total	C	N	O	S	0	6	0
			4381	2764	774	809	34			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	EXPRESSION TAG	UNP P26663
A	-4	HIS	-	EXPRESSION TAG	UNP P26663
A	-3	HIS	-	EXPRESSION TAG	UNP P26663
A	-2	HIS	-	EXPRESSION TAG	UNP P26663
A	-1	HIS	-	EXPRESSION TAG	UNP P26663
A	0	HIS	-	EXPRESSION TAG	UNP P26663
A	1	HIS	-	EXPRESSION TAG	UNP P26663
B	-5	MET	-	EXPRESSION TAG	UNP P26663
B	-4	HIS	-	EXPRESSION TAG	UNP P26663
B	-3	HIS	-	EXPRESSION TAG	UNP P26663
B	-2	HIS	-	EXPRESSION TAG	UNP P26663
B	-1	HIS	-	EXPRESSION TAG	UNP P26663
B	0	HIS	-	EXPRESSION TAG	UNP P26663
B	1	HIS	-	EXPRESSION TAG	UNP P26663

- Molecule 2 is N-{3-[(5S)-5-(1,1-DIMETHYLPROPYL)-1-(4-FLUORO-3-METHYLBENZYL)-4-HYDROXY-2-OXO-2,5-DIHYDRO-1H-PYRROL-3-YL]-1,1-DIOXIDO-4H-1,4-BENZOTHAZIN-7-YL}METHANESULFONAMIDE (three-letter code: H59) (formula: C<sub>26</sub>H<sub>30</sub>FN<sub>3</sub>O<sub>6</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			38	26	1	3	6	2		
2	B	1	Total	C	F	N	O	S	0	0
			38	26	1	3	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	349	Total	O	0	0
			349	349		
3	B	389	Total	O	0	0
			389	389		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: RNA-directed RNA polymerase



#### • Molecule 1: RNA-directed RNA polymerase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.86Å 106.42Å 127.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.09 – 2.10 49.09 – 2.10	Depositor EDS
% Data completeness (in resolution range)	93.3 (49.09-2.10) 93.3 (49.09-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.03 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.193 , 0.246 0.196 , 0.245	Depositor DCC
$R_{free}$ test set	3255 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtriage
Anisotropy	0.643	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 53.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9582	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.72 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.7893e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: H59

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.59	1/4482 (0.0%)	0.67	0/6081
1	B	0.60	1/4476 (0.0%)	0.68	1/6073 (0.0%)
All	All	0.60	2/8958 (0.0%)	0.67	1/12154 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	311	CYS	CB-SG	-7.09	1.70	1.82
1	A	311	CYS	CB-SG	-6.95	1.70	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	321	VAL	CB-CA-C	-5.51	100.93	111.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4387	0	4403	31	0
1	B	4381	0	4399	26	0
2	A	38	0	29	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	38	0	29	1	0
3	A	349	0	0	6	1
3	B	389	0	0	5	1
All	All	9582	0	8860	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:MET:SD	3:A:661:HOH:O	2.23	0.94
1:B:329:THR:HG23	3:B:638:HOH:O	1.68	0.93
1:B:82:LEU:HD13	1:B:249:ALA:HB2	1.58	0.86
1:A:182:LEU:HD23	1:A:182:LEU:C	1.96	0.86
1:A:403:THR:HG22	3:A:720:HOH:O	1.81	0.80
1:A:82:LEU:HD13	1:A:249:ALA:HB2	1.68	0.74
1:A:182:LEU:HD23	1:A:182:LEU:O	1.90	0.71
1:B:201:VAL:HG22	1:B:384:LEU:HG	1.75	0.69
1:B:82:LEU:CD1	1:B:249:ALA:HB2	2.23	0.68
1:B:440:GLU:OE2	1:B:457:LEU:HD12	1.93	0.68
1:A:527:ASN:HD21	1:A:534:LEU:H	1.41	0.68
1:B:154:ARG:N	3:B:645:HOH:O	2.26	0.68
2:B:571:H59:H37A	2:B:571:H59:H31A	1.78	0.65
2:A:571:H59:H37A	2:A:571:H59:H31A	1.79	0.65
1:B:461:GLN:H	1:B:461:GLN:HE21	1.46	0.63
1:A:12:THR:O	1:A:139:MET:HE3	1.98	0.63
1:A:201:VAL:HG22	1:A:384:LEU:HG	1.80	0.62
1:B:103:TYR:OH	1:B:118:HIS:HD2	1.85	0.60
1:A:182:LEU:CD2	1:A:182:LEU:C	2.69	0.59
1:A:103:TYR:OH	1:A:118:HIS:HD2	1.85	0.59
1:B:306:ALA:HB3	1:B:308:LEU:HD13	1.85	0.59
1:A:178:VAL:HG23	3:A:657:HOH:O	2.03	0.58
1:B:197:PRO:O	1:B:201:VAL:HG23	2.05	0.57
1:A:201:VAL:HG22	1:A:384:LEU:CD1	2.35	0.57
1:B:313:MET:HG2	1:B:322:VAL:HG22	1.87	0.56
1:A:36[B]:MET:SD	1:A:491:LYS:O	2.64	0.55
1:A:306:ALA:HB3	1:A:308:LEU:HD13	1.90	0.54
1:A:197:PRO:O	1:A:201:VAL:HG23	2.08	0.54
2:A:571:H59:H36A	3:A:822:HOH:O	2.08	0.53
1:A:217:PHE:CE2	1:A:336:LEU:HD21	2.44	0.52

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:119:ILE:HD13	1:B:169:VAL:HG11	1.92	0.51
1:B:306:ALA:CB	1:B:308:LEU:HD13	2.41	0.51
1:A:523:LYS:HG3	1:A:534:LEU:HD23	1.91	0.50
1:A:80:ALA:HB3	1:A:245:LEU:CD2	2.41	0.49
1:A:46:GLY:HA2	1:A:49:GLN:HE21	1.77	0.49
1:A:12:THR:O	1:A:139:MET:CE	2.60	0.49
1:B:182:LEU:HD13	1:B:243:CYS:SG	2.53	0.49
1:A:296:TYR:CD2	1:A:313:MET:CE	2.96	0.48
1:B:262[A]:ILE:HD11	3:B:946:HOH:O	2.14	0.47
1:B:440:GLU:OE2	1:B:457:LEU:CD1	2.60	0.47
1:A:201:VAL:HG22	1:A:384:LEU:CG	2.43	0.47
1:A:77:THR:O	1:B:24:ASN:HB2	2.16	0.46
1:B:372:VAL:HG13	3:B:766:HOH:O	2.16	0.45
1:B:532:THR:HG23	3:B:585:HOH:O	2.16	0.45
1:B:527:ASN:HD21	1:B:534:LEU:H	1.64	0.45
1:A:217:PHE:CD2	1:A:336:LEU:HD21	2.53	0.44
1:B:523:LYS:HG3	1:B:534:LEU:HD23	1.99	0.44
1:A:503:ARG:O	1:A:507:VAL:HG23	2.19	0.42
1:B:411:ASN:O	1:B:415:TYR:HB2	2.20	0.42
1:A:22:PRO:HG2	1:A:401:ARG:HG3	2.01	0.42
1:A:306:ALA:CB	1:A:308:LEU:HD13	2.50	0.42
1:A:412:ILE:O	1:A:416:ALA:HB2	2.20	0.42
1:B:374:HIS:O	1:B:474:LEU:HA	2.20	0.42
1:B:445:CYS:SG	1:B:454:ILE:HD12	2.60	0.42
1:B:52:VAL:HG12	1:B:223[B]:CYS:SG	2.61	0.41
1:A:154:ARG:N	3:A:731:HOH:O	2.53	0.41
1:A:222:ARG:HD3	3:A:764:HOH:O	2.20	0.41
1:A:144:VAL:HB	1:A:394:ARG:HG2	2.04	0.40
1:B:36[A]:MET:O	1:B:147:VAL:HG13	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:737:HOH:O	3:B:853:HOH:O[4_445]	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	560/576 (97%)	552 (99%)	8 (1%)	0	100	100
1	B	559/576 (97%)	545 (98%)	14 (2%)	0	100	100
All	All	1119/1152 (97%)	1097 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	482/491 (98%)	465 (96%)	17 (4%)	41	42
1	B	481/491 (98%)	466 (97%)	15 (3%)	45	48
All	All	963/982 (98%)	931 (97%)	32 (3%)	42	45

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LEU
1	A	82	LEU
1	A	90	LYS
1	A	159	LEU
1	A	182	LEU
1	A	311	CYS
1	A	321	VAL
1	A	355	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	381	VAL
1	A	439	LEU
1	A	461	GLN
1	A	473	SER
1	A	510	ARG
1	A	527	ASN
1	A	532	THR
1	A	536	LEU
1	A	547	LEU
1	B	57	LEU
1	B	82	LEU
1	B	159	LEU
1	B	222	ARG
1	B	321	VAL
1	B	381	VAL
1	B	419	LEU
1	B	439	LEU
1	B	461	GLN
1	B	492	LEU
1	B	527	ASN
1	B	532	THR
1	B	536	LEU
1	B	545	LEU
1	B	547	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	49	GLN
1	A	118	HIS
1	A	273	ASN
1	A	374	HIS
1	A	461	GLN
1	A	483	ASN
1	A	527	ASN
1	B	35	ASN
1	B	49	GLN
1	B	118	HIS
1	B	446	GLN
1	B	461	GLN
1	B	527	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	H59	A	571	-	39,41,41	1.46	6 (15%)	49,65,65	2.05	13 (26%)
2	H59	B	571	-	39,41,41	1.62	4 (10%)	49,65,65	2.22	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	H59	A	571	-	-	0/18/57/57	0/3/4/4
2	H59	B	571	-	-	0/18/57/57	0/3/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	571	H59	O22-S20	-2.58	1.41	1.44

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	571	H59	C21-C18	-2.09	1.34	1.35
2	A	571	H59	C26-S25	2.52	1.81	1.75
2	B	571	H59	C26-S25	2.56	1.81	1.75
2	A	571	H59	C12-S20	3.32	1.80	1.75
2	A	571	H59	C9-N10	3.43	1.49	1.46
2	A	571	H59	C18-N19	4.08	1.40	1.35
2	A	571	H59	C21-S20	4.32	1.76	1.71
2	B	571	H59	C18-N19	5.03	1.41	1.35
2	B	571	H59	C21-S20	6.67	1.78	1.71

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	571	H59	C26-S25-N24	-6.86	98.38	106.64
2	B	571	H59	C18-C21-S20	-5.22	119.93	123.29
2	B	571	H59	C1-C31-N10	-4.99	104.49	113.47
2	A	571	H59	C1-C31-N10	-4.87	104.69	113.47
2	A	571	H59	C18-C21-S20	-4.13	120.63	123.29
2	A	571	H59	C5-C4-C3	-3.95	118.45	123.95
2	B	571	H59	C5-C4-C3	-3.44	119.16	123.95
2	A	571	H59	O30-C11-C7	-3.28	122.60	130.41
2	B	571	H59	O30-C11-C7	-2.67	124.03	130.41
2	A	571	H59	C26-S25-N24	-2.46	103.68	106.64
2	A	571	H59	O23-S20-C12	-2.29	105.97	109.00
2	A	571	H59	O27-S25-N24	-2.14	102.72	107.13
2	A	571	H59	O23-S20-O22	2.05	117.39	113.93
2	A	571	H59	O28-S25-O27	2.22	122.15	118.78
2	A	571	H59	C6-C5-C4	2.36	121.52	119.01
2	A	571	H59	C16-C17-C12	2.61	122.00	119.25
2	B	571	H59	O28-S25-N24	2.87	113.06	107.13
2	B	571	H59	O23-S20-O22	2.88	118.78	113.93
2	A	571	H59	C2-C3-C4	5.54	119.85	116.00
2	B	571	H59	C2-C3-C4	5.87	120.08	116.00
2	A	571	H59	C31-N10-C9	6.78	128.45	121.57
2	B	571	H59	C31-N10-C9	7.00	128.68	121.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	571	H59	2	0
2	B	571	H59	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	557/576 (96%)	-0.22	4 (0%)	87 89	13, 24, 35, 42	0
1	B	557/576 (96%)	-0.23	3 (0%)	90 92	13, 22, 34, 47	0
All	All	1114/1152 (96%)	-0.23	7 (0%)	89 91	13, 23, 34, 47	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	ARG	3.0
1	B	542	ALA	2.5
1	B	95	HIS	2.4
1	A	102	GLY	2.4
1	A	309	GLN	2.2
1	A	502	HIS	2.1
1	B	531	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	H59	A	571	38/38	0.98	0.11	0.48	12,18,22,26	0
2	H59	B	571	38/38	0.97	0.09	-0.40	11,17,20,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.