



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2017 – 06:56 PM EST

PDB ID : 5H6T  
Title : Crystal structure of Hydrazidase from Microbacterium sp. strain HM58-2  
Authors : Akiyama, T.; Ishii, M.; Takuwa, A.; Oinuma, K.; Sasaki, Y.; Takaya, N.; Yajima, S.  
Deposited on : 2016-11-15  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029077
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029077


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 10px; left: 0; width: 100%;"></div> <div style="position: absolute; left: 0; top: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 91%, yellow 91%, yellow 95%, grey 95%, grey 100%);"></div> <div style="position: absolute; left: 91%; top: 0; width: 4%; height: 100%; background-color: yellow;"></div> <div style="position: absolute; left: 95%; top: 0; width: 5%; height: 100%; background-color: grey;"></div> </div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> <span>1%</span> <span>91%</span> <span>5%</span> </div> </div>
1	B	481	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 10px; left: 0; width: 100%;"></div> <div style="position: absolute; left: 0; top: 0; width: 100%; height: 100%; background: linear-gradient(to right, red 0%, red 2%, orange 2%, orange 95%, yellow 95%, yellow 100%);"></div> </div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> <span>2%</span> <span>95%</span> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	1	0
			3439	2142	620	663	14			
1	B	479	Total	C	N	O	S	0	3	0
			3621	2251	661	694	15			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	GLY	-	expression tag	UNP A0A170QJP8
A	-19	SER	-	expression tag	UNP A0A170QJP8
A	-18	SER	-	expression tag	UNP A0A170QJP8
A	-17	HIS	-	expression tag	UNP A0A170QJP8
A	-16	HIS	-	expression tag	UNP A0A170QJP8
A	-15	HIS	-	expression tag	UNP A0A170QJP8
A	-14	HIS	-	expression tag	UNP A0A170QJP8
A	-13	HIS	-	expression tag	UNP A0A170QJP8
A	-12	HIS	-	expression tag	UNP A0A170QJP8
A	-11	SER	-	expression tag	UNP A0A170QJP8
A	-10	SER	-	expression tag	UNP A0A170QJP8
A	-9	GLY	-	expression tag	UNP A0A170QJP8
A	-8	LEU	-	expression tag	UNP A0A170QJP8
A	-7	VAL	-	expression tag	UNP A0A170QJP8
A	-6	PRO	-	expression tag	UNP A0A170QJP8
A	-5	ARG	-	expression tag	UNP A0A170QJP8
A	-4	GLY	-	expression tag	UNP A0A170QJP8
A	-3	SER	-	expression tag	UNP A0A170QJP8
A	-2	HIS	-	expression tag	UNP A0A170QJP8
A	-1	MET	-	expression tag	UNP A0A170QJP8
A	0	ALA	-	expression tag	UNP A0A170QJP8
A	1	SER	-	expression tag	UNP A0A170QJP8
B	-20	GLY	-	expression tag	UNP A0A170QJP8
B	-19	SER	-	expression tag	UNP A0A170QJP8
B	-18	SER	-	expression tag	UNP A0A170QJP8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	HIS	-	expression tag	UNP A0A170QJP8
B	-16	HIS	-	expression tag	UNP A0A170QJP8
B	-15	HIS	-	expression tag	UNP A0A170QJP8
B	-14	HIS	-	expression tag	UNP A0A170QJP8
B	-13	HIS	-	expression tag	UNP A0A170QJP8
B	-12	HIS	-	expression tag	UNP A0A170QJP8
B	-11	SER	-	expression tag	UNP A0A170QJP8
B	-10	SER	-	expression tag	UNP A0A170QJP8
B	-9	GLY	-	expression tag	UNP A0A170QJP8
B	-8	LEU	-	expression tag	UNP A0A170QJP8
B	-7	VAL	-	expression tag	UNP A0A170QJP8
B	-6	PRO	-	expression tag	UNP A0A170QJP8
B	-5	ARG	-	expression tag	UNP A0A170QJP8
B	-4	GLY	-	expression tag	UNP A0A170QJP8
B	-3	SER	-	expression tag	UNP A0A170QJP8
B	-2	HIS	-	expression tag	UNP A0A170QJP8
B	-1	MET	-	expression tag	UNP A0A170QJP8
B	0	ALA	-	expression tag	UNP A0A170QJP8
B	1	SER	-	expression tag	UNP A0A170QJP8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	330	Total O 330 330	0	0
2	B	338	Total O 338 338	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Amidase



- Molecule 1: Amidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.16Å 128.07Å 159.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.97 – 1.60 33.97 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (33.97-1.60) 99.7 (33.97-1.60)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.81 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.165 , 0.178 0.178 , 0.190	Depositor DCC
$R_{free}$ test set	8192 reflections (5.28%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.4	Xtriage
Anisotropy	0.307	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 46.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7728	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.42	0/3498	0.73	0/4758
1	B	0.43	0/3694	0.76	4/5023 (0.1%)
All	All	0.42	0/7192	0.74	4/9781 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	208	ARG	NE-CZ-NH1	7.13	123.86	120.30
1	B	7	ARG	NE-CZ-NH1	7.07	123.84	120.30
1	B	208	ARG	NE-CZ-NH2	-6.93	116.84	120.30
1	B	193	ARG	NE-CZ-NH1	5.11	122.85	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3439	0	3419	7	0
1	B	3621	0	3590	20	0
2	A	330	0	0	2	0
2	B	338	0	0	0	0
All	All	7728	0	7009	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:299:LEU:CD2	1:B:370[A]:LEU:HD11	2.00	0.92
1:B:370[A]:LEU:HD23	1:B:371:MET:HG3	1.58	0.85
1:B:208:ARG:HG2	1:B:208:ARG:HH11	1.51	0.76
1:B:370[A]:LEU:CD2	1:B:371:MET:HG3	2.20	0.71
1:B:256:ILE:HD11	1:B:432:ARG:HH21	1.57	0.70
1:B:299:LEU:HD22	1:B:370[A]:LEU:HD11	1.73	0.69
1:B:256:ILE:HD11	1:B:432:ARG:NH2	2.10	0.66
1:B:208:ARG:CG	1:B:208:ARG:HH11	2.12	0.62
1:B:299:LEU:HD23	1:B:370[A]:LEU:HD11	1.81	0.62
1:B:370[A]:LEU:HD23	1:B:371:MET:CG	2.33	0.56
1:B:370[A]:LEU:HD23	1:B:371:MET:N	2.21	0.55
1:A:4:GLU:HG2	2:A:809:HOH:O	2.07	0.53
1:B:208:ARG:CG	1:B:208:ARG:NH1	2.72	0.52
1:B:174:GLY:O	1:B:179:SER:HB2	2.10	0.52
1:A:245:THR:O	1:B:433:LYS:NZ	2.45	0.49
1:B:-14:HIS:HE1	1:B:-7:VAL:O	1.96	0.49
1:A:411:PRO:HA	1:A:414:ASN:HD22	1.78	0.47
1:A:334:ARG:NH1	2:A:503:HOH:O	2.45	0.47
1:B:299:LEU:CD2	1:B:370[A]:LEU:CD1	2.85	0.45
1:B:83:TYR:O	1:B:122:HIS:HE1	1.98	0.45
1:A:174:GLY:O	1:A:179:SER:HB2	2.17	0.44
1:B:299:LEU:HD23	1:B:370[A]:LEU:CD1	2.46	0.44
1:A:185:SER:HB3	1:A:426:GLY:HA2	2.00	0.43
1:A:150:ARG:HD2	1:A:386:ALA:O	2.19	0.43
1:B:411:PRO:HA	1:B:414:ASN:HD22	1.83	0.42
1:B:181:ARG:NH2	1:B:414:ASN:HD21	2.18	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/481 (95%)	439 (96%)	15 (3%)	1 (0%)	51	27
1	B	480/481 (100%)	467 (97%)	12 (2%)	1 (0%)	51	27
All	All	935/962 (97%)	906 (97%)	27 (3%)	2 (0%)	51	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	381	VAL
1	A	381	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/377 (95%)	353 (99%)	5 (1%)	71	52
1	B	379/377 (100%)	370 (98%)	9 (2%)	54	26
All	All	737/754 (98%)	723 (98%)	14 (2%)	64	37

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	HIS
1	A	179	SER
1	A	208	ARG
1	A	257	GLU
1	A	384	MET
1	B	45	PHE
1	B	125	ILE
1	B	179	SER
1	B	208	ARG
1	B	365	MET
1	B	370[A]	LEU
1	B	370[B]	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	384	MET
1	B	435	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	ASN
1	A	266	HIS
1	A	414	ASN
1	B	-14	HIS
1	B	122	HIS
1	B	266	HIS
1	B	394	HIS
1	B	414	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	456/481 (94%)	-0.16	7 (1%) 74 75	7, 13, 22, 40	2 (0%)
1	B	479/481 (99%)	-0.14	11 (2%) 61 61	8, 13, 24, 37	2 (0%)
All	All	935/962 (97%)	-0.15	18 (1%) 67 67	7, 13, 23, 40	4 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	208	ARG	3.9
1	B	64	PHE	3.8
1	B	208	ARG	3.7
1	A	207	GLY	3.7
1	B	-17	HIS	3.6
1	B	62	ARG	3.4
1	A	64	PHE	3.2
1	B	58	ARG	2.9
1	A	209	SER	2.9
1	B	-16	HIS	2.6
1	B	209	SER	2.4
1	B	63	GLU	2.4
1	A	19	GLU	2.4
1	B	-15	HIS	2.3
1	A	210	ARG	2.3
1	A	327	GLU	2.2
1	B	257	GLU	2.2
1	B	55	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.