



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:53 am GMT

PDB ID : 1H7K
Title : Formation of a tyrosyl radical intermediate in *Proteus mirabilis* catalase by directed mutagenesis and consequences for nucleotide reactivity
Authors : Andreoletti, P.; Gambarelli, S.; Gaillard, J.; Sainz, G.; Stojanoff, V.; Jouve, H.M.
Deposited on : 2001-07-08
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	483	<div> <div>3%</div> <div>82%</div> <div>15%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	700	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4083 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CATALASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	476	3863	2447	685	716	15	0	0	1

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	215	TYR	PHE	CONFLICT	UNP P42321
A	53	OMT	MET	MODIFIED RESIDUE	UNP P42321

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	168	Total 168	O 168	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CATALASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	110.00Å 110.00Å 251.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.26 – 2.40 29.82 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.26-2.40) 99.6 (29.82-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.47 (at 2.39Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.237 , 0.240 0.236 , 0.230	Depositor DCC
R_{free} test set	1411 reflections (4.10%)	DCC
Wilson B-factor (Å ²)	44.5	Xtriage
Anisotropy	0.456	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4083	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, OMT, SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	14/3963 (0.4%)	0.90	18/5363 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	99	ARG	NE-CZ	17.48	1.55	1.33
1	A	99	ARG	CZ-NH1	-14.31	1.14	1.33
1	A	472	GLU	CD-OE2	13.10	1.40	1.25
1	A	357	PRO	N-CA	11.24	1.66	1.47
1	A	364	TYR	CB-CG	-9.98	1.36	1.51
1	A	349	HIS	CG-CD2	-9.31	1.20	1.35
1	A	18	ASN	C-N	-9.06	1.13	1.34
1	A	364	TYR	C-N	8.36	1.53	1.34
1	A	18	ASN	C-O	8.09	1.38	1.23
1	A	356	ALA	C-N	-7.63	1.19	1.34
1	A	153	THR	CA-CB	-7.25	1.34	1.53
1	A	349	HIS	CG-ND1	5.83	1.51	1.38
1	A	103	ASP	CB-CG	5.74	1.63	1.51
1	A	103	ASP	CG-OD2	5.23	1.37	1.25

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	ARG	NE-CZ-NH2	-28.45	106.07	120.30
1	A	99	ARG	NH1-CZ-NH2	15.83	136.82	119.40
1	A	364	TYR	CB-CG-CD2	-13.06	113.16	121.00
1	A	364	TYR	CB-CG-CD1	12.58	128.55	121.00
1	A	103	ASP	CB-CG-OD2	-11.85	107.64	118.30
1	A	357	PRO	CA-N-CD	-8.08	100.18	111.50
1	A	413	ARG	NE-CZ-NH2	7.38	123.99	120.30
1	A	18	ASN	CA-C-O	-6.88	105.65	120.10
1	A	99	ARG	NE-CZ-NH1	-6.41	117.09	120.30
1	A	54	HIS	O-C-N	-6.40	112.47	122.70
1	A	317	ASN	O-C-N	6.14	132.53	122.70
1	A	6	LEU	CA-CB-CG	6.13	129.41	115.30
1	A	365	HIS	N-CA-CB	-6.02	99.76	110.60
1	A	153	THR	CA-CB-CG2	5.96	120.75	112.40
1	A	6	LEU	CB-CG-CD1	-5.61	101.46	111.00
1	A	349	HIS	CB-CG-ND1	-5.44	109.60	123.20
1	A	153	THR	CA-CB-OG1	5.37	120.29	109.00
1	A	364	TYR	C-N-CA	-5.25	108.56	121.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	103	ASP	Sidechain
1	A	18	ASN	Mainchain
1	A	349	HIS	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3863	0	3677	25	0
2	A	43	0	30	3	0
3	A	4	0	3	0	0
4	A	5	0	0	0	0
5	A	168	0	0	0	0
All	All	4083	0	3710	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:600:HEM:HBB2	2:A:600:HEM:HMB2	1.59	0.84
1:A:464:HIS:HD2	1:A:466:GLU:H	1.41	0.68
1:A:423:ARG:HD3	1:A:463:VAL:O	1.95	0.66
2:A:600:HEM:CMB	2:A:600:HEM:HBB2	2.27	0.64
1:A:56:LYS:HE3	1:A:101:ALA:O	1.99	0.63
1:A:189:ARG:HD3	1:A:243:ASP:OD1	2.01	0.61
1:A:54:HIS:CE1	1:A:95:VAL:HG22	2.41	0.56
1:A:132:PHE:HB3	1:A:278:LEU:HD13	1.88	0.56
1:A:278:LEU:HD12	1:A:329:MET:HG3	1.88	0.55
1:A:359:CYS:HB2	1:A:360:PRO:CD	2.37	0.55
1:A:310:GLN:HB3	1:A:352:ILE:HD13	1.89	0.54
1:A:54:HIS:HA	1:A:94:THR:O	2.08	0.54
1:A:179:MET:HG3	1:A:441:ILE:HG12	1.91	0.52
1:A:436:ARG:O	1:A:440:ARG:HG3	2.10	0.52
1:A:433:GLU:O	1:A:437:MET:HG2	2.12	0.49
1:A:131:VAL:HG23	1:A:174:GLN:HE21	1.77	0.49
1:A:73:TYR:HE2	1:A:290:MET:HE1	1.79	0.46
1:A:359:CYS:HB2	1:A:360:PRO:HD2	1.98	0.45
2:A:600:HEM:HBC2	2:A:600:HEM:CMC	2.46	0.44
1:A:254:PRO:HG2	1:A:297:LEU:HB2	2.00	0.43
1:A:398:GLU:HB2	1:A:399:PRO:HD2	2.00	0.43
1:A:5:LYS:HB2	1:A:5:LYS:HE3	1.90	0.42
1:A:173:HIS:HA	1:A:422:PRO:HB3	2.00	0.42
1:A:200:SER:OG	1:A:323:SER:HB3	2.21	0.41
1:A:371:ARG:HA	1:A:371:ARG:HD2	1.83	0.41
1:A:437:MET:O	1:A:441:ILE:HG13	2.21	0.40
1:A:253:TYR:HA	1:A:254:PRO:HD2	1.96	0.40
1:A:128:ASN:HA	1:A:192:HIS:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	473/483 (98%)	454 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	407/413 (98%)	379 (93%)	28 (7%)	18	28

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	VAL
1	A	30	LEU
1	A	37	LEU
1	A	47	VAL
1	A	52	ARG
1	A	113	LEU
1	A	117	THR
1	A	129	THR
1	A	147	VAL
1	A	172	LEU
1	A	191	VAL
1	A	201	PHE
1	A	237	ARG
1	A	242	ARG
1	A	244	LEU
1	A	270	THR
1	A	271	VAL
1	A	278	LEU
1	A	284	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	297	LEU
1	A	330	LEU
1	A	334	LEU
1	A	363	ASN
1	A	366	ARG
1	A	414	GLU
1	A	429	LEU
1	A	459	LEU
1	A	467	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	HIS
1	A	121	ASN
1	A	127	ASN
1	A	174	GLN
1	A	259	GLN
1	A	300	ASN
1	A	341	HIS
1	A	363	ASN
1	A	464	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	OMT	A	53	1	9,9,10	2.00	3 (33%)	8,12,14	2.40	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OMT	A	53	1	-	0/6/8/10	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	53	OMT	CG-SD	-4.14	1.72	1.78
1	A	53	OMT	CE-SD	-2.92	1.62	1.75
1	A	53	OMT	CA-C	2.49	1.53	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	53	OMT	OD2-SD-OD1	-4.08	107.08	116.93
1	A	53	OMT	OD1-SD-CG	-3.15	106.17	108.36
1	A	53	OMT	OD1-SD-CE	2.85	111.94	108.93

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	600	1	28,50,50	1.86	7 (25%)	17,82,82	1.43	4 (23%)
3	ACT	A	700	-	1,3,3	3.01	1 (100%)	0,3,3	0.00	-
4	SO4	A	800	-	4,4,4	0.88	0	6,6,6	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	600	1	-	0/6/54/54	0/0/8/8
3	ACT	A	700	-	-	0/0/0/0	0/0/0/0
4	SO4	A	800	-	-	0/0/0/0	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	HEM	C3C-C2C	-4.04	1.35	1.40
3	A	700	ACT	CH3-C	-3.01	1.44	1.48
2	A	600	HEM	C3B-C2B	-2.02	1.37	1.40
2	A	600	HEM	C1B-NB	2.00	1.39	1.36
2	A	600	HEM	C3B-CAB	2.85	1.53	1.47
2	A	600	HEM	C3C-CAC	3.14	1.54	1.47
2	A	600	HEM	CBB-CAB	3.57	1.54	1.28
2	A	600	HEM	CBC-CAC	3.72	1.55	1.28

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	HEM	CMA-C3A-C4A	-2.67	124.36	128.46
2	A	600	HEM	CMD-C2D-C1D	-2.46	124.69	128.46
2	A	600	HEM	CMB-C2B-C3B	2.38	129.32	124.89
2	A	600	HEM	CMC-C2C-C3C	2.58	129.68	124.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	HEM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	475/483 (98%)	0.04	16 (3%) 46 44	19, 40, 54, 59	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	PRO	3.9
1	A	431	ASP	3.4
1	A	284	HIS	3.2
1	A	81	GLU	2.9
1	A	227	ASP	2.8
1	A	450	LYS	2.7
1	A	362	HIS	2.7
1	A	395	ASP	2.6
1	A	451	GLU	2.6
1	A	414	GLU	2.4
1	A	359	CYS	2.4
1	A	361	PHE	2.3
1	A	252	ASP	2.1
1	A	479	GLU	2.1
1	A	205	ASP	2.0
1	A	404	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	OMT	A	53	10/11	0.93	0.14	-	26,31,37,38	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ACT	A	700	4/4	0.83	0.37	9.47	74,74,74,74	0
2	HEM	A	600	43/43	0.97	0.12	-0.69	18,23,24,27	0
4	SO4	A	800	5/5	0.94	0.12	-	90,90,90,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.