



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2017 – 01:50 am GMT

PDB ID : 3H7L
Title : CRYSTAL STRUCTURE OF ENDOGLUCANASE-RELATED PROTEIN FROM *Vibrio parahaemolyticus*
Authors : Patskovsky, Y.; Toro, R.; Morano, C.; Rutter, M.; Chang, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2009-04-27
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	586	<div> <div>4%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div>••</div> </div> </div>
1	B	586	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>••</div> </div> </div>
1	C	586	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	586	-	-	-	X
2	GOL	B	587	-	-	-	X
2	GOL	C	587	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDOGLUCANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	576	Total	C	N	O	S	0	2	0
			4617	2924	789	886	18			
1	B	575	Total	C	N	O	S	0	4	0
			4622	2926	792	886	18			
1	C	575	Total	C	N	O	S	0	6	0
			4635	2934	793	890	18			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	expression tag	UNP Q87LX4
A	1	SER	-	expression tag	UNP Q87LX4
A	87	THR	TYR	engineered	UNP Q87LX4
A	132	ALA	VAL	engineered	UNP Q87LX4
A	368	THR	GLY	engineered	UNP Q87LX4
A	579	GLY	-	expression tag	UNP Q87LX4
A	580	HIS	-	expression tag	UNP Q87LX4
A	581	HIS	-	expression tag	UNP Q87LX4
A	582	HIS	-	expression tag	UNP Q87LX4
A	583	HIS	-	expression tag	UNP Q87LX4
A	584	HIS	-	expression tag	UNP Q87LX4
A	585	HIS	-	expression tag	UNP Q87LX4
B	0	MET	-	expression tag	UNP Q87LX4
B	1	SER	-	expression tag	UNP Q87LX4
B	87	THR	TYR	engineered	UNP Q87LX4
B	132	ALA	VAL	engineered	UNP Q87LX4
B	368	THR	GLY	engineered	UNP Q87LX4
B	579	GLY	-	expression tag	UNP Q87LX4
B	580	HIS	-	expression tag	UNP Q87LX4
B	581	HIS	-	expression tag	UNP Q87LX4
B	582	HIS	-	expression tag	UNP Q87LX4
B	583	HIS	-	expression tag	UNP Q87LX4
B	584	HIS	-	expression tag	UNP Q87LX4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	585	HIS	-	expression tag	UNP Q87LX4
C	0	MET	-	expression tag	UNP Q87LX4
C	1	SER	-	expression tag	UNP Q87LX4
C	87	THR	TYR	engineered	UNP Q87LX4
C	132	ALA	VAL	engineered	UNP Q87LX4
C	368	THR	GLY	engineered	UNP Q87LX4
C	579	GLY	-	expression tag	UNP Q87LX4
C	580	HIS	-	expression tag	UNP Q87LX4
C	581	HIS	-	expression tag	UNP Q87LX4
C	582	HIS	-	expression tag	UNP Q87LX4
C	583	HIS	-	expression tag	UNP Q87LX4
C	584	HIS	-	expression tag	UNP Q87LX4
C	585	HIS	-	expression tag	UNP Q87LX4

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	203	Total 203	O 203	0	0
3	B	334	Total 334	O 334	0	0
3	C	373	Total 373	O 373	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ENDOGLUCANASE



• Molecule 1: ENDOGLUCANASE



• Molecule 1: ENDOGLUCANASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	166.12Å 166.12Å 343.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.30 31.39 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.30) 99.9 (31.39-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.20	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, R_{free}	0.185 , 0.232 0.186 , 0.231	Depositor DCC
R_{free} test set	3732 reflections (3.10%)	DCC
Wilson B-factor (Å ²)	25.4	Xtriage
Anisotropy	0.388	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14814	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/4741	0.61	0/6434
1	B	0.51	0/4752	0.61	0/6448
1	C	0.51	0/4768	0.62	0/6469
All	All	0.50	0/14261	0.61	0/19351

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4617	0	4362	19	0
1	B	4622	0	4370	23	0
1	C	4635	0	4382	21	0
2	A	6	0	8	0	0
2	B	12	0	16	0	0
2	C	12	0	16	1	0
3	A	203	0	0	0	0
3	B	334	0	0	1	0
3	C	373	0	0	0	0
All	All	14814	0	13154	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:246:GLY:H	1:B:249:GLN:HE21	1.34	0.74
1:B:246:GLY:H	1:B:249:GLN:NE2	1.91	0.69
1:C:493:GLY:H	1:C:498:ASN:ND2	1.96	0.63
1:C:465:ILE:HA	1:C:468:GLN:HE21	1.63	0.62
1:A:20:ILE:HB	1:A:63:TYR:HB2	1.82	0.62
1:A:21:GLN:HE22	1:A:485:PRO:HA	1.64	0.62
1:A:21:GLN:HE21	1:A:62:PHE:HE1	1.46	0.61
1:A:120:ASP:HA	1:A:123:VAL:HG22	1.83	0.59
1:A:20:ILE:HD11	1:A:80:VAL:HG21	1.85	0.59
1:B:493:GLY:H	1:B:498:ASN:ND2	2.02	0.58
1:A:515:CYS:HA	1:A:548:GLN:HA	1.86	0.57
1:B:549:TRP:HB3	1:B:552:HIS:CD2	2.40	0.56
1:B:557:LEU:HG	1:B:561:MET:HE1	1.88	0.56
1:C:3:LEU:HG	1:C:85:SER:HB3	1.87	0.55
1:C:20:ILE:HD11	1:C:80:VAL:HG21	1.86	0.55
1:C:515:CYS:HA	1:C:548:GLN:HA	1.87	0.55
1:C:166:VAL:O	1:C:170:ILE:HG12	2.07	0.54
1:A:493:GLY:H	1:A:498:ASN:ND2	2.06	0.54
1:A:465:ILE:HA	1:A:468:GLN:HE21	1.73	0.54
1:B:120:ASP:HA	1:B:123:VAL:CG2	2.38	0.53
1:C:549:TRP:HB3	1:C:552:HIS:CD2	2.44	0.53
1:B:295:ASN:HD22	1:B:295:ASN:H	1.56	0.51
1:C:493:GLY:H	1:C:498:ASN:HD21	1.56	0.51
1:C:488:MET:HG3	1:C:497:ASN:O	2.10	0.51
1:B:352:GLN:HE22	1:C:25:PRO:HB2	1.75	0.51
1:C:20:ILE:HB	1:C:63:TYR:HB2	1.93	0.50
1:A:549:TRP:HB3	1:A:552:HIS:CD2	2.46	0.50
1:B:515:CYS:HA	1:B:548:GLN:HA	1.93	0.49
1:A:368:THR:HB	1:A:452:MET:HG3	1.93	0.49
1:B:516:ASN:HA	1:B:552:HIS:NE2	2.27	0.49
1:B:465:ILE:HA	1:B:468:GLN:HE21	1.78	0.48
1:B:173:GLY:HA3	1:B:561:MET:HE3	1.95	0.48
1:C:308:LEU:HD11	1:C:374:TYR:HB2	1.96	0.48
1:A:516:ASN:HA	1:A:552:HIS:NE2	2.29	0.48
1:A:557:LEU:HG	1:A:561:MET:HE1	1.97	0.47
1:A:103:VAL:O	1:A:106:TYR:HB3	2.16	0.46
1:B:403:ASN:HD22	1:B:403:ASN:N	2.15	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:MET:HE1	3:B:836:HOH:O	2.16	0.45
1:B:333:ARG:HD3	1:C:26:HIS:ND1	2.32	0.44
1:A:217:PHE:CZ	1:A:219:LYS:HA	2.52	0.44
1:A:3:LEU:HG	1:A:85:SER:HB3	1.98	0.44
1:C:368:THR:HB	1:C:452:MET:HG3	2.01	0.43
1:B:217:PHE:CZ	1:B:219:LYS:HA	2.54	0.43
1:C:300:ASN:HB2	1:C:350:ALA:O	2.18	0.43
1:C:146:SER:HB3	1:C:216:VAL:O	2.18	0.43
1:A:286:LEU:O	1:A:290:ASN:HB2	2.19	0.42
1:B:516:ASN:ND2	1:B:543:TRP:O	2.52	0.42
1:C:533:PRO:HD2	2:C:587:GOL:H31	2.01	0.42
1:A:322:ARG:O	1:A:326:GLU:HG2	2.20	0.42
1:C:51:GLN:HG3	1:C:408:PRO:HB3	2.02	0.42
1:B:549:TRP:CG	1:B:551:PRO:HD2	2.55	0.42
1:B:550:ILE:HG13	1:B:551:PRO:HD3	2.02	0.42
1:B:33:GLN:HG2	1:B:79:GLN:HB3	2.02	0.42
1:C:324:LEU:O	1:C:328:ARG:HG3	2.20	0.42
1:B:111:ARG:HH22	1:B:193:GLU:HG2	1.85	0.42
1:C:213:TYR:CD1	1:C:229:ILE:HG21	2.55	0.42
1:A:324:LEU:O	1:A:328:ARG:HG3	2.20	0.41
1:B:120:ASP:HA	1:B:123:VAL:HG23	2.01	0.41
1:A:149:LEU:HG	1:A:164:PRO:HG3	2.01	0.41
1:B:24:GLN:HA	1:B:25:PRO:HD2	1.93	0.41
1:C:287:LYS:HD2	1:C:326:GLU:OE2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	576/586 (98%)	551 (96%)	23 (4%)	2 (0%)	44 55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	577/586 (98%)	553 (96%)	22 (4%)	2 (0%)	44	55
1	C	579/586 (99%)	550 (95%)	28 (5%)	1 (0%)	51	63
All	All	1732/1758 (98%)	1654 (96%)	73 (4%)	5 (0%)	44	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	81	GLU
1	C	437	TYR
1	B	81	GLU
1	A	437	TYR
1	B	437	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	483/490 (99%)	460 (95%)	23 (5%)	30	40
1	B	484/490 (99%)	465 (96%)	19 (4%)	37	51
1	C	486/490 (99%)	470 (97%)	16 (3%)	43	59
All	All	1453/1470 (99%)	1395 (96%)	58 (4%)	36	50

All (58) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	23	GLU
1	A	33	GLN
1	A	56	ASN
1	A	64	LEU
1	A	81	GLU
1	A	83	SER
1	A	84	ARG
1	A	92	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	122	GLN
1	A	123	VAL
1	A	131	THR
1	A	174	LEU
1	A	273	GLN
1	A	287	LYS
1	A	321	THR
1	A	383	GLN
1	A	388	LYS
1	A	432	ASN
1	A	483	LEU
1	A	539	MET
1	A	557	LEU
1	A	565	GLN
1	B	33	GLN
1	B	56	ASN
1	B	82	ASP
1	B	83	SER
1	B	122	GLN
1	B	131	THR
1	B	174	LEU
1	B	215	THR
1	B	287	LYS
1	B	295	ASN
1	B	388	LYS
1	B	389	CYS
1	B	397	PHE
1	B	403	ASN
1	B	404	LYS
1	B	496	ARG
1	B	536	LYS
1	B	557	LEU
1	B	569	GLN
1	C	3	LEU
1	C	33	GLN
1	C	56	ASN
1	C	81	GLU
1	C	83	SER
1	C	122[A]	GLN
1	C	122[B]	GLN
1	C	131	THR
1	C	174	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	215	THR
1	C	219	LYS
1	C	287	LYS
1	C	380	ASP
1	C	404	LYS
1	C	496	ARG
1	C	569	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	GLN
1	A	105	HIS
1	A	277	ASN
1	A	295	ASN
1	A	383	GLN
1	A	403	ASN
1	A	468	GLN
1	A	498	ASN
1	A	565	GLN
1	A	569	GLN
1	B	33	GLN
1	B	93	HIS
1	B	98	GLN
1	B	249	GLN
1	B	277	ASN
1	B	292	GLN
1	B	295	ASN
1	B	332	GLN
1	B	343	GLN
1	B	352	GLN
1	B	403	ASN
1	B	468	GLN
1	B	498	ASN
1	C	33	GLN
1	C	277	ASN
1	C	332	GLN
1	C	343	GLN
1	C	468	GLN
1	C	498	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	586	-	5,5,5	0.30	0	5,5,5	0.57	0
2	GOL	B	586	-	5,5,5	0.39	0	5,5,5	0.70	0
2	GOL	B	587	-	5,5,5	0.29	0	5,5,5	0.46	0
2	GOL	C	586	-	5,5,5	0.43	0	5,5,5	0.56	0
2	GOL	C	587	-	5,5,5	0.53	0	5,5,5	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	586	-	-	0/4/4/4	0/0/0/0
2	GOL	B	586	-	-	0/4/4/4	0/0/0/0
2	GOL	B	587	-	-	0/4/4/4	0/0/0/0
2	GOL	C	586	-	-	0/4/4/4	0/0/0/0
2	GOL	C	587	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	587	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	576/586 (98%)	0.00	22 (3%) 41 48	22, 43, 73, 114	0
1	B	575/586 (98%)	-0.47	8 (1%) 75 80	9, 27, 58, 115	0
1	C	575/586 (98%)	-0.53	7 (1%) 79 82	11, 26, 52, 100	0
All	All	1726/1758 (98%)	-0.33	37 (2%) 64 70	9, 32, 66, 115	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	576	ILE	7.5
1	B	574	LYS	5.4
1	B	573	SER	4.7
1	B	575	ASN	4.5
1	A	572	THR	4.4
1	C	575	ASN	3.7
1	A	575	ASN	3.3
1	A	128	ALA	3.3
1	A	574	LYS	3.1
1	B	43	VAL	3.1
1	C	572	THR	3.1
1	A	464	GLU	2.9
1	C	573	SER	2.7
1	A	465	ILE	2.6
1	A	461	ALA	2.6
1	A	225	ALA	2.6
1	B	41	GLN	2.5
1	A	342	GLU	2.5
1	A	250	GLY	2.5
1	A	381	SER	2.5
1	A	39	SER	2.4
1	A	240	PHE	2.4
1	A	573	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	208	GLU	2.3
1	C	379	ASP	2.3
1	A	379	ASP	2.3
1	A	462	SER	2.3
1	A	420	ASN	2.3
1	C	461	ALA	2.2
1	A	129	ASN	2.2
1	C	464	GLU	2.1
1	C	82	ASP	2.1
1	B	461	ALA	2.1
1	A	41	GLN	2.1
1	B	572	THR	2.1
1	A	382	VAL	2.1
1	B	39	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GOL	C	587	6/6	0.91	0.17	8.73	24,30,35,40	0
2	GOL	B	587	6/6	0.95	0.16	5.29	27,41,43,44	0
2	GOL	B	586	6/6	0.95	0.17	3.80	30,35,39,42	0
2	GOL	C	586	6/6	0.98	0.13	1.88	22,29,31,35	0
2	GOL	A	586	6/6	0.94	0.15	1.29	40,54,58,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.