



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:15 pm GMT

PDB ID : 2HCO  
Title : THE STRUCTURE OF HUMAN CARBONMONOXY HAEMOGLOBIN AT  
2.7 ANGSTROMS RESOLUTION  
Authors : Baldwin, J.M.  
Deposited on : 1979-08-07  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	141	
2	B	146	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMOGLOBIN (CARBONMONOXY) (ALPHA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	141	Total	C	N	O	S	0	0	0
			1069	685	187	194	3			

- Molecule 2 is a protein called HEMOGLOBIN (CARBONMONOXY) (BETA CHAIN).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	146	Total	C	N	O	S	0	0	0
			1123	724	195	201	3			

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			2	1	1		
4	B	1	Total	C	O	0	0
			2	1	1		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: HEMOGLOBIN (CARBONMONOXY) (ALPHA CHAIN)



#### • Molecule 2: HEMOGLOBIN (CARBONMONOXY) (BETA CHAIN)



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.70Å 53.70Å 193.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.70	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.70)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REAL-SPACE REFINEMENT	Depositor
R, $R_{free}$	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2282	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	0.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	1/1097 (0.1%)	0.98	3/1491 (0.2%)
2	B	0.72	2/1153 (0.2%)	1.01	3/1566 (0.2%)
All	All	0.69	3/2250 (0.1%)	1.00	6/3057 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	8
All	All	0	10

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	37	TRP	NE1-CE2	-7.71	1.27	1.37
1	A	14	TRP	NE1-CE2	-6.96	1.28	1.37
2	B	15	TRP	NE1-CE2	-6.31	1.29	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	30	ARG	NE-CZ-NH1	9.01	124.80	120.30
2	B	30	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	A	1	VAL	CA-CB-CG2	6.80	121.11	110.90
1	A	1	VAL	CA-CB-CG1	6.70	120.94	110.90
1	A	92	ARG	NE-CZ-NH1	6.53	123.56	120.30
2	B	40	ARG	NE-CZ-NH1	5.25	122.93	120.30

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	116	GLU	Mainchain
1	A	78	ASN	Sidechain
2	B	102	ASN	Sidechain
2	B	127	GLN	Sidechain
2	B	131	GLN	Sidechain
2	B	30	ARG	Sidechain
2	B	77	HIS	Mainchain
2	B	80	ASN	Sidechain,Mainchain
2	B	81	LEU	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1069	0	1073	24	3
2	B	1123	0	1118	19	0
3	A	43	0	30	4	0
3	B	43	0	30	4	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
All	All	2282	0	2251	46	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:LEU:HD11	3:A:150:HEM:HBA2	1.52	0.90
3:A:150:HEM:HHD	3:A:150:HEM:HBC2	1.64	0.77
1:A:17:VAL:HG23	1:A:113:LEU:HD11	1.71	0.72
3:B:150:HEM:HHA	3:B:150:HEM:CBD	2.25	0.67
1:A:17:VAL:CG2	1:A:113:LEU:HD11	2.26	0.66
1:A:76:MET:HB3	1:A:135:VAL:HG21	1.86	0.58

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:LEU:HD11	1:A:105:LEU:HD21	1.86	0.58
1:A:31:ARG:HG2	2:B:124:PRO:HB3	1.88	0.55
3:B:150:HEM:HHA	3:B:150:HEM:HBD2	1.89	0.54
1:A:95:PRO:HA	1:A:98:PHE:HD2	1.73	0.53
2:B:20:VAL:HA	2:B:68:LEU:HD12	1.89	0.53
1:A:35:SER:HB3	2:B:131:GLN:HG3	1.92	0.52
2:B:107:GLY:HA3	2:B:134:VAL:HG13	1.91	0.52
2:B:1:VAL:HG21	2:B:78:LEU:O	2.11	0.51
1:A:42:TYR:C	1:A:44:PRO:HD3	2.31	0.51
2:B:75:LEU:HD12	2:B:133:VAL:HG11	1.93	0.50
1:A:1:VAL:HG22	1:A:2:LEU:H	1.76	0.49
1:A:47:ASP:C	1:A:49:SER:H	2.15	0.48
2:B:24:GLY:HA2	2:B:68:LEU:HG	1.96	0.48
1:A:106:LEU:HD21	1:A:125:LEU:HB3	1.95	0.48
1:A:27:GLU:O	1:A:31:ARG:HG3	2.14	0.47
1:A:83:LEU:HA	1:A:83:LEU:HD13	1.74	0.47
1:A:139:LYS:HD3	1:A:139:LYS:O	2.15	0.47
1:A:85:ASP:O	1:A:89:HIS:HB2	2.14	0.47
1:A:3:SER:HB2	1:A:4:PRO:HD2	1.97	0.46
2:B:57:ASN:HB3	2:B:60:VAL:HB	1.98	0.46
2:B:42:PHE:O	2:B:45:PHE:HB2	2.16	0.45
1:A:42:TYR:CD2	3:A:150:HEM:HBC1	2.51	0.45
2:B:105:LEU:O	2:B:109:VAL:HG13	2.17	0.45
3:B:150:HEM:HHA	3:B:150:HEM:HBD1	1.96	0.45
1:A:34:LEU:HA	1:A:34:LEU:HD12	1.74	0.44
1:A:125:LEU:HA	1:A:125:LEU:HD12	1.73	0.44
1:A:105:LEU:HA	1:A:105:LEU:HD12	1.80	0.43
1:A:56:LYS:HE3	1:A:56:LYS:HB2	1.75	0.43
2:B:99:ASP:HA	2:B:100:PRO:HD3	1.88	0.43
2:B:24:GLY:CA	2:B:68:LEU:HG	2.48	0.43
2:B:14:LEU:HA	2:B:14:LEU:HD23	1.74	0.42
2:B:29:GLY:HA3	2:B:55:MET:SD	2.59	0.42
2:B:15:TRP:O	2:B:18:VAL:HB	2.20	0.42
1:A:26:ALA:HB2	1:A:59:GLY:HA3	2.02	0.42
2:B:3:LEU:HB3	2:B:8:LYS:HG2	2.02	0.42
2:B:144:LYS:HD3	2:B:144:LYS:HA	1.83	0.41
1:A:101:LEU:O	1:A:105:LEU:HB2	2.21	0.41
2:B:91:LEU:O	2:B:95:LYS:HB2	2.20	0.41
2:B:42:PHE:HE1	3:B:150:HEM:HBC2	1.85	0.41
3:A:150:HEM:CBC	3:A:150:HEM:HHD	2.44	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:ASP:OD2	1:A:68:ASN:ND2[5_645]	1.60	0.60
1:A:47:ASP:CG	1:A:68:ASN:ND2[5_645]	1.91	0.29
1:A:47:ASP:OD1	1:A:68:ASN:ND2[5_645]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	128 (92%)	9 (6%)	2 (1%)	13	33
2	B	144/146 (99%)	137 (95%)	5 (4%)	2 (1%)	13	33
All	All	283/287 (99%)	265 (94%)	14 (5%)	4 (1%)	13	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	HIS
1	A	18	GLY
2	B	77	HIS
2	B	3	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	105 (93%)	8 (7%)	17	39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	118/118 (100%)	99 (84%)	19 (16%)	3	7
All	All	231/231 (100%)	204 (88%)	27 (12%)	6	14

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	VAL
1	A	17	VAL
1	A	48	LEU
1	A	49	SER
1	A	81	SER
1	A	96	VAL
1	A	109	LEU
1	A	125	LEU
2	B	3	LEU
2	B	8	LYS
2	B	11	VAL
2	B	14	LEU
2	B	20	VAL
2	B	26	GLU
2	B	30	ARG
2	B	50	THR
2	B	66	LYS
2	B	68	LEU
2	B	73	ASP
2	B	87	THR
2	B	91	LEU
2	B	101	GLU
2	B	104	ARG
2	B	108	ASN
2	B	109	VAL
2	B	110	LEU
2	B	146	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	ASN
1	A	97	ASN
2	B	102	ASN
2	B	108	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	HEM	A	150	1,4	28,50,50	1.45	4 (14%)	17,82,82	1.07	1 (5%)
4	CMO	A	151	3	0,1,1	0.00	-	0,0,0	0.00	-
3	HEM	B	150	2,4	28,50,50	1.28	3 (10%)	17,82,82	1.22	0
4	CMO	B	151	3	0,1,1	0.00	-	0,0,0	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	150	1,4	-	0/6/54/54	0/0/8/8
4	CMO	A	151	3	-	0/0/0/0	0/0/0/0
3	HEM	B	150	2,4	-	0/6/54/54	0/0/8/8
4	CMO	B	151	3	-	0/0/0/0	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	150	HEM	C3C-C2C	-3.89	1.35	1.40
3	B	150	HEM	C3C-C2C	-3.32	1.36	1.40
3	A	150	HEM	C3B-C2B	-3.16	1.36	1.40
3	B	150	HEM	C3B-C2B	-2.22	1.37	1.40
3	B	150	HEM	C1B-NB	2.50	1.39	1.36
3	A	150	HEM	C4C-NC	2.52	1.39	1.36
3	A	150	HEM	C4D-ND	2.71	1.40	1.36

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	150	HEM	CMA-C3A-C4A	-2.04	125.33	128.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	150	HEM	4	0
3	B	150	HEM	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.