



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 03:37 am GMT

PDB ID : 5HDH
Title : Crystal structure of human TLR8 with an uncleaved Z-loop
Authors : Tanji, H.; Ohto, U.; Shimizu, T.
Deposited on : 2016-01-05
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	801	<div> <div>5%</div> <div> <div></div> <div>62%</div> <div>28%</div> <div>• 6%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	917	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6371 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Toll-like receptor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	751	6042	3865	1029	1128	20	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	452	ASN	ARG	engineered mutation	UNP Q9NR97
A	453	GLN	LYS	engineered mutation	UNP Q9NR97
A	454	SER	ARG	engineered mutation	UNP Q9NR97
A	455	ASN	ARG	engineered mutation	UNP Q9NR97

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	79	Total	O	0	0
			79	79		

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	171.52Å 171.52Å 301.33Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.02 – 2.60 45.02 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.02-2.60) 100.0 (45.02-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.206 , 0.239 0.212 , 0.243	Depositor DCC
R_{free} test set	2610 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	66.7	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 65.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6371	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, NAG, MES, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	0/6164	0.83	6/8355 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	137	LEU	C-N-CD	6.63	142.32	128.40
1	A	472	ARG	C-N-CD	6.03	141.06	128.40
1	A	57	VAL	C-N-CD	5.92	140.82	128.40
1	A	144	LEU	C-N-CD	5.57	140.10	128.40
1	A	138	PRO	CA-N-CD	-5.25	104.15	111.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	353	TYR	Sidechain
1	A	466	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6042	0	6035	293	0
2	A	168	0	151	7	0
3	A	22	0	17	0	0
4	A	33	0	30	0	0
5	A	15	0	0	0	0
6	A	12	0	13	1	0
7	A	79	0	0	5	0
All	All	6371	0	6246	294	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 294 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:THR:HA	1:A:117:THR:CG2	1.49	1.42
1:A:79:THR:HA	1:A:117:THR:HG21	1.28	1.14
1:A:79:THR:HA	1:A:117:THR:HG22	1.30	1.12
1:A:755:ALA:HA	1:A:786:TRP:HZ3	1.07	1.11
1:A:755:ALA:HA	1:A:786:TRP:CZ3	1.89	1.07

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	739/801 (92%)	675 (91%)	57 (8%)	7 (1%)	20	40

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	PHE
1	A	466	ASN
1	A	627	ASP
1	A	467	PHE
1	A	378	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	698/745 (94%)	671 (96%)	27 (4%)	37	65

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	277	ASN
1	A	466	ASN
1	A	808	ASP
1	A	338	ARG
1	A	117	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	233	GLN
1	A	277	ASN
1	A	595	ASN
1	A	231	ASN
1	A	625	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	901	1	14,14,15	0.29	0	15,19,21	0.57	0
2	NAG	A	902	1,2	14,14,15	0.99	2 (14%)	15,19,21	2.04	3 (20%)
2	NAG	A	903	3,2	14,14,15	1.00	1 (7%)	15,19,21	1.27	2 (13%)
3	BMA	A	904	2,4	11,11,12	0.35	0	13,15,17	0.81	0
4	MAN	A	905	3	11,11,12	0.64	0	13,15,17	0.93	1 (7%)
2	NAG	A	906	1	14,14,15	0.60	0	15,19,21	0.82	0
2	NAG	A	907	1	14,14,15	0.58	0	15,19,21	1.15	1 (6%)
2	NAG	A	908	1,2	14,14,15	0.59	0	15,19,21	1.52	5 (33%)
2	NAG	A	909	2	14,14,15	0.28	0	15,19,21	0.57	0
2	NAG	A	910	1	14,14,15	0.87	1 (7%)	15,19,21	1.51	2 (13%)
2	NAG	A	911	1	14,14,15	0.56	0	15,19,21	0.95	0
2	NAG	A	912	1,2	14,14,15	0.67	0	15,19,21	1.36	2 (13%)
2	NAG	A	913	3,2	14,14,15	0.82	1 (7%)	15,19,21	1.83	3 (20%)
3	BMA	A	914	2,4	11,11,12	0.65	0	13,15,17	1.75	2 (15%)
4	MAN	A	915	3	11,11,12	0.79	0	13,15,17	1.28	2 (15%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MAN	A	916	3	11,11,12	0.42	0	13,15,17	0.92	1 (7%)
2	NAG	A	917	1	14,14,15	0.63	0	15,19,21	1.32	2 (13%)
5	SO4	A	918	-	4,4,4	0.39	0	6,6,6	0.33	0
5	SO4	A	919	-	4,4,4	0.40	0	6,6,6	0.24	0
5	SO4	A	920	-	4,4,4	0.47	0	6,6,6	0.27	0
6	MES	A	921	-	12,12,12	2.21	1 (8%)	14,16,16	2.07	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	901	1	-	0/6/23/26	0/1/1/1
2	NAG	A	902	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	903	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	904	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	905	3	-	0/2/19/22	0/1/1/1
2	NAG	A	906	1	-	0/6/23/26	0/1/1/1
2	NAG	A	907	1	-	0/6/23/26	0/1/1/1
2	NAG	A	908	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	909	2	-	0/6/23/26	0/1/1/1
2	NAG	A	910	1	-	0/6/23/26	0/1/1/1
2	NAG	A	911	1	-	0/6/23/26	0/1/1/1
2	NAG	A	912	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	913	3,2	-	0/6/23/26	0/1/1/1
3	BMA	A	914	2,4	-	0/2/19/22	0/1/1/1
4	MAN	A	915	3	-	0/2/19/22	0/1/1/1
4	MAN	A	916	3	-	0/2/19/22	0/1/1/1
2	NAG	A	917	1	-	0/6/23/26	0/1/1/1
5	SO4	A	918	-	-	0/0/0/0	0/0/0/0
5	SO4	A	919	-	-	0/0/0/0	0/0/0/0
5	SO4	A	920	-	-	0/0/0/0	0/0/0/0
6	MES	A	921	-	-	0/6/14/14	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	921	MES	C8-S	-7.19	1.66	1.77
2	A	903	NAG	O5-C1	-2.63	1.39	1.43
2	A	910	NAG	O5-C1	-2.25	1.40	1.43
2	A	913	NAG	O5-C1	-2.16	1.40	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	902	NAG	O5-C1	-2.09	1.40	1.43

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	902	NAG	O5-C1-C2	-5.97	103.17	111.47
2	A	913	NAG	O5-C1-C2	-4.73	104.89	111.47
2	A	913	NAG	C2-N2-C7	-3.42	117.95	122.94
2	A	903	NAG	O5-C1-C2	-3.39	106.75	111.47
2	A	902	NAG	C1-O5-C5	-3.21	107.74	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	NAG	2	0
2	A	902	NAG	5	0
6	A	921	MES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	751/801 (93%)	0.13	43 (5%) 24 18	37, 73, 136, 171	0

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	64	TYR	6.2
1	A	467	PHE	5.1
1	A	74	PHE	4.8
1	A	808	ASP	4.8
1	A	42	ASN	4.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	917	14/15	0.96	0.23	3.12	64,72,85,87	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	A	907	14/15	0.87	0.27	1.03	81,95,108,116	0
2	NAG	A	906	14/15	0.95	0.15	-0.31	64,73,81,82	0
2	NAG	A	902	14/15	0.96	0.15	-0.60	44,52,62,73	0
2	NAG	A	903	14/15	0.98	0.13	-1.12	52,60,72,74	0
2	NAG	A	908	14/15	0.98	0.13	-1.64	42,47,57,64	0
2	NAG	A	912	14/15	0.99	0.13	-1.66	43,46,51,54	0
2	NAG	A	910	14/15	0.92	0.14	-	77,91,103,104	0
5	SO4	A	920	5/5	0.96	0.08	-	97,98,109,112	0
3	BMA	A	914	11/12	0.93	0.15	-	68,97,107,125	0
2	NAG	A	901	14/15	0.87	0.16	-	83,106,122,122	0
2	NAG	A	911	14/15	0.82	0.23	-	101,115,142,145	0
5	SO4	A	919	5/5	0.90	0.17	-	94,108,132,136	0
3	BMA	A	904	11/12	0.87	0.17	-	68,98,115,130	0
4	MAN	A	916	11/12	0.84	0.17	-	109,119,129,131	0
4	MAN	A	905	11/12	0.71	0.24	-	100,122,138,138	0
5	SO4	A	918	5/5	0.96	0.11	-	71,85,90,96	0
4	MAN	A	915	11/12	0.87	0.14	-	88,103,119,128	0
6	MES	A	921	12/12	0.84	0.25	-	58,78,82,86	12
2	NAG	A	909	14/15	0.91	0.17	-	58,94,121,129	0
2	NAG	A	913	14/15	0.97	0.10	-	42,53,68,75	0

6.5 Other polymers [i](#)

There are no such residues in this entry.