



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 9, 2017 – 11:04 am GMT

PDB ID : 5HDP
Title : Hydrolase StnA mutant - S185A
Authors : Qian, T.
Deposited on : 2016-01-05
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	recalc28906
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28906


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	383	<div> <div>74%</div> <div>8%</div> <div>17%</div> </div>
1	B	383	<div> <div>74%</div> <div>8%</div> <div>17%</div> </div>
1	C	383	<div> <div>73%</div> <div>9%</div> <div>17%</div> </div>
1	D	383	<div> <div>76%</div> <div>7%</div> <div>17%</div> </div>
1	E	383	<div> <div>78%</div> <div>5%</div> <div>17%</div> </div>
1	F	383	<div> <div>76%</div> <div>6%</div> <div>18%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	383	<div><div><div>%</div><div><div></div></div><div>77%</div><div>5%</div><div>17%</div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16999 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	316	Total	C	N	O	S	0	0	0
			2390	1497	416	466	11			
1	A	316	Total	C	N	O	S	0	0	0
			2390	1497	416	466	11			
1	B	316	Total	C	N	O	S	0	0	0
			2390	1497	416	466	11			
1	C	316	Total	C	N	O	S	0	0	0
			2390	1497	416	466	11			
1	E	316	Total	C	N	O	S	0	0	0
			2390	1497	416	466	11			
1	F	314	Total	C	N	O	S	0	0	0
			2380	1491	414	464	11			
1	G	316	Total	C	N	O	S	0	0	0
			2390	1497	416	466	11			

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1	VAL	-	expression tag	UNP L7PIJ2
D	185	ALA	SER	engineered mutation	UNP L7PIJ2
D	376	LEU	-	expression tag	UNP L7PIJ2
D	377	GLU	-	expression tag	UNP L7PIJ2
D	378	HIS	-	expression tag	UNP L7PIJ2
D	379	HIS	-	expression tag	UNP L7PIJ2
D	380	HIS	-	expression tag	UNP L7PIJ2
D	381	HIS	-	expression tag	UNP L7PIJ2
D	382	HIS	-	expression tag	UNP L7PIJ2
D	383	HIS	-	expression tag	UNP L7PIJ2
A	1	VAL	-	expression tag	UNP L7PIJ2
A	185	ALA	SER	engineered mutation	UNP L7PIJ2
A	376	LEU	-	expression tag	UNP L7PIJ2
A	377	GLU	-	expression tag	UNP L7PIJ2
A	378	HIS	-	expression tag	UNP L7PIJ2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	379	HIS	-	expression tag	UNP L7PIJ2
A	380	HIS	-	expression tag	UNP L7PIJ2
A	381	HIS	-	expression tag	UNP L7PIJ2
A	382	HIS	-	expression tag	UNP L7PIJ2
A	383	HIS	-	expression tag	UNP L7PIJ2
B	1	VAL	-	expression tag	UNP L7PIJ2
B	185	ALA	SER	engineered mutation	UNP L7PIJ2
B	376	LEU	-	expression tag	UNP L7PIJ2
B	377	GLU	-	expression tag	UNP L7PIJ2
B	378	HIS	-	expression tag	UNP L7PIJ2
B	379	HIS	-	expression tag	UNP L7PIJ2
B	380	HIS	-	expression tag	UNP L7PIJ2
B	381	HIS	-	expression tag	UNP L7PIJ2
B	382	HIS	-	expression tag	UNP L7PIJ2
B	383	HIS	-	expression tag	UNP L7PIJ2
C	1	VAL	-	expression tag	UNP L7PIJ2
C	185	ALA	SER	engineered mutation	UNP L7PIJ2
C	376	LEU	-	expression tag	UNP L7PIJ2
C	377	GLU	-	expression tag	UNP L7PIJ2
C	378	HIS	-	expression tag	UNP L7PIJ2
C	379	HIS	-	expression tag	UNP L7PIJ2
C	380	HIS	-	expression tag	UNP L7PIJ2
C	381	HIS	-	expression tag	UNP L7PIJ2
C	382	HIS	-	expression tag	UNP L7PIJ2
C	383	HIS	-	expression tag	UNP L7PIJ2
E	1	VAL	-	expression tag	UNP L7PIJ2
E	185	ALA	SER	engineered mutation	UNP L7PIJ2
E	376	LEU	-	expression tag	UNP L7PIJ2
E	377	GLU	-	expression tag	UNP L7PIJ2
E	378	HIS	-	expression tag	UNP L7PIJ2
E	379	HIS	-	expression tag	UNP L7PIJ2
E	380	HIS	-	expression tag	UNP L7PIJ2
E	381	HIS	-	expression tag	UNP L7PIJ2
E	382	HIS	-	expression tag	UNP L7PIJ2
E	383	HIS	-	expression tag	UNP L7PIJ2
F	1	VAL	-	expression tag	UNP L7PIJ2
F	185	ALA	SER	engineered mutation	UNP L7PIJ2
F	376	LEU	-	expression tag	UNP L7PIJ2
F	377	GLU	-	expression tag	UNP L7PIJ2
F	378	HIS	-	expression tag	UNP L7PIJ2
F	379	HIS	-	expression tag	UNP L7PIJ2
F	380	HIS	-	expression tag	UNP L7PIJ2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	381	HIS	-	expression tag	UNP L7PIJ2
F	382	HIS	-	expression tag	UNP L7PIJ2
F	383	HIS	-	expression tag	UNP L7PIJ2
G	1	VAL	-	expression tag	UNP L7PIJ2
G	185	ALA	SER	engineered mutation	UNP L7PIJ2
G	376	LEU	-	expression tag	UNP L7PIJ2
G	377	GLU	-	expression tag	UNP L7PIJ2
G	378	HIS	-	expression tag	UNP L7PIJ2
G	379	HIS	-	expression tag	UNP L7PIJ2
G	380	HIS	-	expression tag	UNP L7PIJ2
G	381	HIS	-	expression tag	UNP L7PIJ2
G	382	HIS	-	expression tag	UNP L7PIJ2
G	383	HIS	-	expression tag	UNP L7PIJ2

- Molecule 2 is methyl 5-amino-6-(7-amino-6-methoxy-5,8-dioxo-5,8-dihydroquinolin-2-yl)-4-(2-hydroxy-3-methoxyphenyl)-3-methylpyridine-2-carboxylate (three-letter code: STM) (formula: C₂₅H₂₂N₄O₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	D	1	Total	C	N	O	0	0
			36	25	4	7		
2	A	1	Total	C	N	O	0	0
			36	25	4	7		
2	B	1	Total	C	N	O	0	0
			36	25	4	7		
2	C	1	Total	C	N	O	0	0
			36	25	4	7		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	E	1	Total	C	N	O	0	0
			36	25	4	7		
2	F	1	Total	C	N	O	0	0
			36	25	4	7		
2	G	1	Total	C	N	O	0	0
			36	25	4	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	5	Total	O	0	0
			5	5		
3	A	2	Total	O	0	0
			2	2		
3	B	9	Total	O	0	0
			9	9		
3	C	2	Total	O	0	0
			2	2		
3	E	4	Total	O	0	0
			4	4		
3	F	3	Total	O	0	0
			3	3		
3	G	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Hydrolase






• Molecule 1: Hydrolase



• Molecule 1: Hydrolase



• Molecule 1: Hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	83.26Å 92.85Å 104.04Å 115.06° 106.09° 97.69°	Depositor
Resolution (Å)	46.07 – 2.90 46.07 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.1 (46.07-2.90) 78.5 (46.07-2.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.244 , 0.298 0.245 , 0.296	Depositor DCC
R_{free} test set	2786 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	52.7	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 11.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	16999	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: STM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2448	0.73	0/3342
1	B	0.52	0/2448	0.72	0/3342
1	C	0.50	0/2448	0.70	0/3342
1	D	0.54	0/2448	0.75	0/3342
1	E	0.49	0/2448	0.68	0/3342
1	F	0.47	0/2438	0.69	1/3328 (0.0%)
1	G	0.48	0/2448	0.70	0/3342
All	All	0.50	0/17126	0.71	1/23380 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	370	ARG	NE-CZ-NH1	6.35	123.47	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2390	0	2286	17	0
1	B	2390	0	2286	19	0
1	C	2390	0	2286	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2390	0	2286	14	0
1	E	2390	0	2286	9	0
1	F	2380	0	2279	9	0
1	G	2390	0	2286	15	0
2	A	36	0	0	2	0
2	B	36	0	0	1	0
2	C	36	0	0	2	0
2	D	36	0	0	3	0
2	E	36	0	0	1	0
2	F	36	0	0	1	0
2	G	36	0	0	4	0
3	A	2	0	0	1	0
3	B	9	0	0	0	0
3	C	2	0	0	0	0
3	D	5	0	0	0	0
3	E	4	0	0	1	0
3	F	3	0	0	0	0
3	G	2	0	0	0	0
All	All	16999	0	15995	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:260:ARG:NH2	1:B:346:ASP:OD1	2.20	0.75
1:G:342:ILE:CD1	1:G:363:ALA:HB1	2.21	0.71
1:A:260:ARG:NH2	1:A:346:ASP:OD1	2.24	0.70
1:D:340:ASP:OD2	1:D:370:ARG:NH2	2.25	0.69
1:A:260:ARG:NH1	1:A:307:ARG:HG3	2.07	0.69

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/383 (82%)	301 (96%)	13 (4%)	0	100	100
1	B	314/383 (82%)	301 (96%)	13 (4%)	0	100	100
1	C	314/383 (82%)	299 (95%)	14 (4%)	1 (0%)	44	77
1	D	314/383 (82%)	301 (96%)	13 (4%)	0	100	100
1	E	314/383 (82%)	299 (95%)	14 (4%)	1 (0%)	44	77
1	F	312/383 (82%)	294 (94%)	17 (5%)	1 (0%)	44	77
1	G	314/383 (82%)	299 (95%)	15 (5%)	0	100	100
All	All	2196/2681 (82%)	2094 (95%)	99 (4%)	3 (0%)	55	85

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	74	GLY
1	F	74	GLY
1	E	175	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/304 (83%)	249 (98%)	4 (2%)	68	90
1	B	253/304 (83%)	251 (99%)	2 (1%)	85	96
1	C	253/304 (83%)	249 (98%)	4 (2%)	68	90
1	D	253/304 (83%)	251 (99%)	2 (1%)	85	96
1	E	253/304 (83%)	250 (99%)	3 (1%)	75	93
1	F	253/304 (83%)	250 (99%)	3 (1%)	75	93
1	G	253/304 (83%)	253 (100%)	0	100	100
All	All	1771/2128 (83%)	1753 (99%)	18 (1%)	80	95

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	90	SER
1	C	103	THR
1	E	277	ASN
1	B	90	SER
1	B	277	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	361	GLN
1	F	361	GLN
1	E	361	GLN
1	A	361	GLN
1	C	361	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	STM	A	401	-	39,39,39	1.37	2 (5%)	49,58,58	3.62	22 (44%)
2	STM	B	401	-	39,39,39	1.39	2 (5%)	49,58,58	1.84	14 (28%)
2	STM	C	401	-	39,39,39	1.47	3 (7%)	49,58,58	2.25	19 (38%)
2	STM	D	401	-	39,39,39	1.49	3 (7%)	49,58,58	2.30	17 (34%)
2	STM	E	401	-	39,39,39	1.49	2 (5%)	49,58,58	2.22	17 (34%)
2	STM	F	401	-	39,39,39	1.41	2 (5%)	49,58,58	2.09	15 (30%)
2	STM	G	401	-	39,39,39	1.45	2 (5%)	49,58,58	2.32	16 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	STM	A	401	-	-	0/18/38/38	0/4/4/4
2	STM	B	401	-	-	0/18/38/38	0/4/4/4
2	STM	C	401	-	-	0/18/38/38	0/4/4/4
2	STM	D	401	-	-	0/18/38/38	0/4/4/4
2	STM	E	401	-	-	0/18/38/38	0/4/4/4
2	STM	F	401	-	-	0/18/38/38	0/4/4/4
2	STM	G	401	-	-	1/18/38/38	0/4/4/4

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	STM	C3-C4	-2.49	1.38	1.41
2	D	401	STM	C2-C3	-2.08	1.39	1.41
2	A	401	STM	O1-C6	4.57	1.43	1.33
2	B	401	STM	O1-C6	4.60	1.43	1.33
2	D	401	STM	C14-C15	5.05	1.47	1.39

The worst 5 of 120 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	STM	O6-C22-C21	-8.89	109.44	124.37
2	C	401	STM	C14-C15-C16	-4.45	118.11	122.26
2	D	401	STM	C14-C15-C16	-4.31	118.23	122.26
2	A	401	STM	O2-C13-C12	-4.09	117.37	122.03
2	E	401	STM	C14-C15-C16	-3.97	118.56	122.26

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	G	401	STM	C7-O1-C6-C5

There are no ring outliers.

7 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	STM	2	0
2	B	401	STM	1	0
2	C	401	STM	2	0
2	D	401	STM	3	0
2	E	401	STM	1	0
2	F	401	STM	1	0
2	G	401	STM	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/383 (82%)	-0.31	0 100 100	28, 44, 64, 108	1 (0%)
1	B	316/383 (82%)	-0.29	1 (0%) 93 93	28, 47, 68, 123	1 (0%)
1	C	316/383 (82%)	-0.26	1 (0%) 93 93	31, 50, 75, 95	1 (0%)
1	D	316/383 (82%)	-0.32	2 (0%) 89 88	24, 40, 61, 101	1 (0%)
1	E	316/383 (82%)	-0.25	1 (0%) 93 93	31, 53, 77, 90	1 (0%)
1	F	314/383 (81%)	-0.15	3 (0%) 82 81	32, 59, 81, 100	1 (0%)
1	G	316/383 (82%)	-0.21	2 (0%) 89 88	39, 59, 76, 83	1 (0%)
All	All	2210/2681 (82%)	-0.26	10 (0%) 90 90	24, 50, 75, 123	7 (0%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	291	ALA	4.2
1	D	74	GLY	4.1
1	F	307	ARG	3.4
1	E	307	ARG	2.8
1	B	237	ARG	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	STM	E	401	36/36	0.94	0.21	0.30	53,64,70,71	0
2	STM	C	401	36/36	0.94	0.18	-0.04	36,39,46,47	0
2	STM	F	401	36/36	0.91	0.18	-0.14	45,59,69,73	0
2	STM	A	401	36/36	0.95	0.16	-0.37	30,34,39,44	0
2	STM	D	401	36/36	0.96	0.16	-0.38	25,31,36,39	0
2	STM	G	401	36/36	0.92	0.16	-0.39	46,55,64,74	0
2	STM	B	401	36/36	0.95	0.16	-0.59	33,36,43,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.