



wwPDB/EMDataBank EM Map/Model Validation Summary Report ⓘ

Sep 23, 2017 – 08:14 AM EDT

PDB ID : 5HI9
EMDB ID: : EMD-6580
Title : Structure of the full-length TRPV2 channel by cryo-electron microscopy
Authors : Huynh, K.W.; Cohen, M.R.; Jiansen, J.; Samanta, A.; Lodowski, D.T.; Zhou, Z.H.; Moiseenkova-Bell, V.Y.
Deposited on : unknown
Resolution : 4.40 Å(reported)

This is a wwPDB/EMDataBank EM Map/Model Validation Summary Report
for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Clashscore | 125131 | 1336 |
| Ramachandran outliers | 121729 | 1120 |
| Sidechain outliers | 121581 | 1026 |

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 770 | |
| 1 | B | 770 | |
| 1 | C | 770 | |
| 1 | D | 770 | |

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 19636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transient Receptor Potential Cation Channel Subfamily V Member 2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 1 | A | 602 | Total | C | N | O | S | 0 | 0 |
| | | | 4909 | 3207 | 825 | 849 | 28 | | |
| 1 | B | 602 | Total | C | N | O | S | 0 | 0 |
| | | | 4909 | 3207 | 825 | 849 | 28 | | |
| 1 | C | 602 | Total | C | N | O | S | 0 | 0 |
| | | | 4909 | 3207 | 825 | 849 | 28 | | |
| 1 | D | 602 | Total | C | N | O | S | 0 | 0 |
| | | | 4909 | 3207 | 825 | 849 | 28 | | |

There are 40 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| A | 151 | PRO | LEU | conflict | UNP Q9WUD2 |
| A | 762 | THR | - | expression tag | UNP Q9WUD2 |
| A | 763 | GLU | - | expression tag | UNP Q9WUD2 |
| A | 764 | THR | - | expression tag | UNP Q9WUD2 |
| A | 765 | SER | - | expression tag | UNP Q9WUD2 |
| A | 766 | GLN | - | expression tag | UNP Q9WUD2 |
| A | 767 | VAL | - | expression tag | UNP Q9WUD2 |
| A | 768 | ALA | - | expression tag | UNP Q9WUD2 |
| A | 769 | PRO | - | expression tag | UNP Q9WUD2 |
| A | 770 | ALA | - | expression tag | UNP Q9WUD2 |
| B | 151 | PRO | LEU | conflict | UNP Q9WUD2 |
| B | 762 | THR | - | expression tag | UNP Q9WUD2 |
| B | 763 | GLU | - | expression tag | UNP Q9WUD2 |
| B | 764 | THR | - | expression tag | UNP Q9WUD2 |
| B | 765 | SER | - | expression tag | UNP Q9WUD2 |
| B | 766 | GLN | - | expression tag | UNP Q9WUD2 |
| B | 767 | VAL | - | expression tag | UNP Q9WUD2 |
| B | 768 | ALA | - | expression tag | UNP Q9WUD2 |
| B | 769 | PRO | - | expression tag | UNP Q9WUD2 |
| B | 770 | ALA | - | expression tag | UNP Q9WUD2 |
| C | 151 | PRO | LEU | conflict | UNP Q9WUD2 |

Continued on next page...


Continued from previous page...

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------------|------------|
| C | 762 | THR | - | expression tag | UNP Q9WUD2 |
| C | 763 | GLU | - | expression tag | UNP Q9WUD2 |
| C | 764 | THR | - | expression tag | UNP Q9WUD2 |
| C | 765 | SER | - | expression tag | UNP Q9WUD2 |
| C | 766 | GLN | - | expression tag | UNP Q9WUD2 |
| C | 767 | VAL | - | expression tag | UNP Q9WUD2 |
| C | 768 | ALA | - | expression tag | UNP Q9WUD2 |
| C | 769 | PRO | - | expression tag | UNP Q9WUD2 |
| C | 770 | ALA | - | expression tag | UNP Q9WUD2 |
| D | 151 | PRO | LEU | conflict | UNP Q9WUD2 |
| D | 762 | THR | - | expression tag | UNP Q9WUD2 |
| D | 763 | GLU | - | expression tag | UNP Q9WUD2 |
| D | 764 | THR | - | expression tag | UNP Q9WUD2 |
| D | 765 | SER | - | expression tag | UNP Q9WUD2 |
| D | 766 | GLN | - | expression tag | UNP Q9WUD2 |
| D | 767 | VAL | - | expression tag | UNP Q9WUD2 |
| D | 768 | ALA | - | expression tag | UNP Q9WUD2 |
| D | 769 | PRO | - | expression tag | UNP Q9WUD2 |
| D | 770 | ALA | - | expression tag | UNP Q9WUD2 |


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transient Receptor Potential Cation Channel Subfamily V Member 2




- Molecule 1: Transient Receptor Potential Cation Channel Subfamily V Member 2






• Molecule 1: Transient Receptor Potential Cation Channel Subfamily V Member 2


Chain C: 58% 18% 22%



• Molecule 1: Transient Receptor Potential Cation Channel Subfamily V Member 2

Chain D: 58% 18% 22%





4 Experimental information

| Property | Value | Source |
|--------------------------------------|---------------------------|-----------|
| Reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, C4 | Depositor |
| Number of particles used | 42550 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | NONE | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 21 | Depositor |
| Minimum defocus (nm) | Not provided | Depositor |
| Maximum defocus (nm) | Not provided | Depositor |
| Magnification | 31000 | Depositor |
| Image detector | GATAN K2 SUMMIT (4k x 4k) | Depositor |

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|-------------|-------------|-----------------|
| | | RMSZ | $\# Z > 2$ | RMSZ | $\# Z > 2$ |
| 1 | A | 0.41 | 0/5031 | 0.69 | 8/6822 (0.1%) |
| 1 | B | 0.41 | 0/5031 | 0.69 | 8/6822 (0.1%) |
| 1 | C | 0.41 | 0/5031 | 0.69 | 8/6822 (0.1%) |
| 1 | D | 0.41 | 0/5031 | 0.69 | 8/6822 (0.1%) |
| All | All | 0.41 | 0/20124 | 0.69 | 32/27288 (0.1%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | A | 0 | 16 |
| 1 | B | 0 | 16 |
| 1 | C | 0 | 16 |
| 1 | D | 0 | 16 |
| All | All | 0 | 64 |

There are no bond length outliers.

The worst 5 of 32 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 1 | B | 307 | LYS | CB-CA-C | 9.96 | 130.32 | 110.40 |
| 1 | A | 307 | LYS | CB-CA-C | 9.95 | 130.30 | 110.40 |
| 1 | D | 307 | LYS | CB-CA-C | 9.95 | 130.30 | 110.40 |
| 1 | C | 307 | LYS | CB-CA-C | 9.93 | 130.27 | 110.40 |
| 1 | A | 493 | GLU | CB-CA-C | -6.84 | 96.72 | 110.40 |

There are no chirality outliers.

5 of 64 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 1 | A | 166 | SER | Peptide |
| 1 | A | 174 | LYS | Peptide |
| 1 | A | 175 | ARG | Peptide |
| 1 | A | 177 | LEU | Peptide |
| 1 | A | 308 | ILE | Peptide |

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 4909 | 0 | 4965 | 236 | 0 |
| 1 | B | 4909 | 0 | 4965 | 243 | 0 |
| 1 | C | 4909 | 0 | 4965 | 241 | 0 |
| 1 | D | 4909 | 0 | 4965 | 234 | 0 |
| All | All | 19636 | 0 | 19860 | 918 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 918 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|----------------|--------------------------|-------------------|
| 1:A:223:TRP:HE3 | 1:A:270:MET:SD | 1.54 | 1.30 |
| 1:D:223:TRP:HE3 | 1:D:270:MET:SD | 1.54 | 1.30 |
| 1:C:223:TRP:HE3 | 1:C:270:MET:SD | 1.54 | 1.28 |
| 1:B:223:TRP:HE3 | 1:B:270:MET:SD | 1.54 | 1.28 |
| 1:A:223:TRP:CE3 | 1:A:270:MET:SD | 2.34 | 1.20 |

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|----|
| 1 | A | 596/770 (77%) | 545 (91%) | 46 (8%) | 5 (1%) | 22 | 67 |
| 1 | B | 596/770 (77%) | 545 (91%) | 46 (8%) | 5 (1%) | 22 | 67 |
| 1 | C | 596/770 (77%) | 545 (91%) | 46 (8%) | 5 (1%) | 22 | 67 |
| 1 | D | 596/770 (77%) | 545 (91%) | 46 (8%) | 5 (1%) | 22 | 67 |
| All | All | 2384/3080 (77%) | 2180 (91%) | 184 (8%) | 20 (1%) | 27 | 67 |

5 of 20 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 167 | ALA |
| 1 | A | 464 | TRP |
| 1 | B | 167 | ALA |
| 1 | B | 464 | TRP |
| 1 | C | 167 | ALA |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|----|
| 1 | A | 535/683 (78%) | 530 (99%) | 5 (1%) | 82 | 91 |
| 1 | B | 535/683 (78%) | 530 (99%) | 5 (1%) | 82 | 91 |
| 1 | C | 535/683 (78%) | 530 (99%) | 5 (1%) | 82 | 91 |
| 1 | D | 535/683 (78%) | 530 (99%) | 5 (1%) | 82 | 91 |
| All | All | 2140/2732 (78%) | 2120 (99%) | 20 (1%) | 83 | 91 |

5 of 20 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 694 | ARG |
| 1 | C | 152 | LEU |
| 1 | D | 300 | LYS |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 373 | VAL |
| 1 | B | 479 | GLN |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | B | 396 | ASN |
| 1 | C | 129 | GLN |
| 1 | D | 396 | ASN |
| 1 | B | 452 | GLN |
| 1 | C | 127 | ASN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.