



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 01:14 am GMT

PDB ID : 2HLO
Title : Crystal Structure of Fragment D-dimer from Human Fibrin Complexed with Gly-hydroxyPro-Arg-Pro-amide
Authors : Doolittle, R.F.; Kollman, J.M.; Chen, A.; Pandi, L.
Deposited on : 2006-07-08
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	87	
1	D	87	
2	B	328	
2	E	328	
3	C	324	
3	F	324	

Continued on next page...

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 10750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fibrin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	67	547	337	103	104	3	0	0	0
1	D	54	441	269	84	85	3	0	0	0

- Molecule 2 is a protein called Fibrin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	303	2428	1515	429	462	22	0	0	0
2	E	296	2377	1484	420	451	22	0	0	0

- Molecule 3 is a protein called Fibrin, gamma polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	292	2343	1485	396	451	11	0	0	0
3	F	285	2287	1453	384	439	11	0	0	0

- Molecule 4 is a protein called GLY-HYP-ARG-PRO-AMIDE PEPTIDE LIGAND.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	G	5	31	18	8	5	0	0	1
4	H	5	31	18	8	5	0	0	1

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 6 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	E	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	2	Total	Ca	0	0
			2	2		
7	C	2	Total	Ca	0	0
			2	2		
7	F	1	Total	Ca	0	0
			1	1		
7	E	2	Total	Ca	0	0
			2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	8	Total	O	0	0
			8	8		
8	B	55	Total	O	0	0
			55	55		
8	C	44	Total	O	0	0
			44	44		
8	D	13	Total	O	0	0
			13	13		
8	E	46	Total	O	0	0
			46	46		
8	F	33	Total	O	0	0
			33	33		
8	G	1	Total	O	0	0
			1	1		
8	H	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fibrin alpha chain



- Molecule 1: Fibrin alpha chain




- Molecule 2: Fibrin beta chain




- Molecule 2: Fibrin beta chain






• Molecule 3: Fibrin, gamma polypeptide



• Molecule 3: Fibrin, gamma polypeptide



• Molecule 4: GLY-HYP-ARG-PRO-AMIDE PEPTIDE LIGAND



• Molecule 4: GLY-HYP-ARG-PRO-AMIDE PEPTIDE LIGAND





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.69Å 149.48Å 232.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.60 29.12 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.0 (30.00-2.60) 93.7 (29.12-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	34.39 (at 2.61Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.210 , 0.268 0.210 , 0.267	Depositor DCC
R_{free} test set	2825 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	45.6	Xtrriage
Anisotropy	0.113	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	10750	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HYP, CA, NAG, NDG, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/548	0.50	0/731
1	D	0.30	0/441	0.51	0/587
2	B	0.35	0/2490	0.62	0/3364
2	E	0.35	0/2438	0.61	0/3291
3	C	0.37	0/2408	0.60	0/3257
3	F	0.36	0/2351	0.58	0/3180
4	G	0.60	0/21	0.71	0/25
4	H	0.57	0/21	0.60	0/25
All	All	0.35	0/10718	0.60	0/14460

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	547	0	573	32	0
1	D	441	0	458	19	0
2	B	2428	0	2295	78	0
2	E	2377	0	2244	103	0
3	C	2343	0	2188	79	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	2287	0	2136	58	0
4	G	31	0	32	4	0
4	H	31	0	32	2	0
5	B	28	0	25	2	0
6	E	28	0	25	3	0
7	B	2	0	0	0	0
7	C	2	0	0	0	0
7	E	2	0	0	0	0
7	F	1	0	0	0	0
8	A	8	0	0	2	0
8	B	55	0	0	2	0
8	C	44	0	0	2	0
8	D	13	0	0	1	0
8	E	46	0	0	4	0
8	F	33	0	0	6	0
8	G	1	0	0	0	0
8	H	2	0	0	0	0
All	All	10750	0	10008	348	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 348 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:ARG:HH21	1:A:189:ILE:HD13	1.25	1.02
3:C:338:LYS:N	3:C:339:CYS:HA	1.85	0.89
1:A:136:LEU:HD22	2:B:168:LEU:HD21	1.56	0.85
2:B:160:ASN:HB2	2:B:161:ILE:HD13	1.59	0.84
1:D:175:LEU:H	1:D:175:LEU:HD23	1.46	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	65/87 (75%)	60 (92%)	2 (3%)	3 (5%)	3	3
1	D	52/87 (60%)	50 (96%)	2 (4%)	0	100	100
2	B	301/328 (92%)	274 (91%)	23 (8%)	4 (1%)	14	29
2	E	294/328 (90%)	272 (92%)	19 (6%)	3 (1%)	18	37
3	C	290/324 (90%)	266 (92%)	19 (7%)	5 (2%)	11	21
3	F	283/324 (87%)	262 (93%)	17 (6%)	4 (1%)	13	26
4	G	2/5 (40%)	1 (50%)	0	1 (50%)	0	0
4	H	2/5 (40%)	2 (100%)	0	0	100	100
All	All	1289/1488 (87%)	1187 (92%)	82 (6%)	20 (2%)	11	23

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	127	ILE
1	A	191	LYS
2	B	281	ASP
2	B	458	PHE
3	C	104	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	62/82 (76%)	60 (97%)	2 (3%)	44	72
1	D	50/82 (61%)	46 (92%)	4 (8%)	14	27
2	B	261/286 (91%)	247 (95%)	14 (5%)	26	49
2	E	254/286 (89%)	245 (96%)	9 (4%)	41	68
3	C	246/270 (91%)	239 (97%)	7 (3%)	49	76
3	F	239/270 (88%)	230 (96%)	9 (4%)	38	66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	G	2/2 (100%)	2 (100%)	0	100	100
4	H	2/2 (100%)	2 (100%)	0	100	100
All	All	1116/1280 (87%)	1071 (96%)	45 (4%)	36	64

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	317	ASN
1	D	175	LEU
3	F	285	ASP
3	C	325	ASN
1	D	182	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	319	ASN
2	E	174	ASN
3	F	307	HIS
1	D	181	GLN
2	E	189	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HYP	G	2	4	7,8,9	1.02	0	5,10,12	1.33	1 (20%)
4	HYP	H	2	4	7,8,9	1.07	1 (14%)	5,10,12	1.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HYP	G	2	4	-	0/0/11/13	0/1/1/1
4	HYP	H	2	4	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	2	HYP	CA-C	2.30	1.53	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	2	HYP	CG-CB-CA	-2.25	100.96	103.93

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	B	470	2,5	14,14,15	0.75	0	15,19,21	1.15	2 (13%)
5	NAG	B	471	5	14,14,15	0.60	0	15,19,21	0.66	0
6	NAG	E	470	2,6	14,14,15	0.55	0	15,19,21	0.64	0
6	NDG	E	471	6	14,14,15	0.67	0	15,19,21	0.80	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	470	2,5	-	0/6/23/26	0/1/1/1
5	NAG	B	471	5	-	0/6/23/26	0/1/1/1
6	NAG	E	470	2,6	-	0/6/23/26	0/1/1/1
6	NDG	E	471	6	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	470	NAG	O5-C1-C2	-2.17	108.46	111.47
6	E	471	NDG	C2-N2-C7	-2.05	119.95	122.94
5	B	470	NAG	C3-C4-C5	2.23	114.14	110.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	470	NAG	2	0
5	B	471	NAG	2	0
6	E	470	NAG	3	0
6	E	471	NDG	3	0

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	67/87 (77%)	0.17	7 (10%) 7 4	33, 69, 124, 158	0
1	D	54/87 (62%)	-0.12	4 (7%) 15 11	29, 65, 111, 139	0
2	B	303/328 (92%)	-0.26	9 (2%) 51 43	23, 41, 103, 166	0
2	E	296/328 (90%)	-0.25	12 (4%) 38 30	24, 40, 99, 142	0
3	C	292/324 (90%)	-0.36	3 (1%) 82 79	24, 44, 79, 129	0
3	F	285/324 (87%)	-0.38	4 (1%) 75 71	26, 45, 78, 123	0
4	G	3/5 (60%)	-0.64	0 100 100	48, 48, 51, 69	0
4	H	3/5 (60%)	-0.67	0 100 100	42, 42, 46, 61	0
All	All	1303/1488 (87%)	-0.28	39 (2%) 51 43	23, 44, 99, 166	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	459	PRO	9.5
2	E	459	PRO	8.1
2	B	458	PHE	7.1
2	E	390	PRO	6.7
1	A	128	GLU	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	HYP	G	2	8/9	0.98	0.11	-	37,47,50,52	0
4	HYP	H	2	8/9	0.98	0.10	-	40,46,50,53	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	NDG	E	471	14/15	0.65	0.41	-	105,118,126,127	0
5	NAG	B	470	14/15	0.61	0.42	-	121,132,143,154	0
6	NAG	E	470	14/15	0.86	0.29	-	65,85,107,114	0
5	NAG	B	471	14/15	0.54	0.48	-	120,154,156,157	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	CA	B	3	1/1	0.96	0.16	2.49	53,53,53,53	0
7	CA	F	1	1/1	0.98	0.15	1.52	44,44,44,44	0
7	CA	E	3	1/1	0.95	0.14	1.11	47,47,47,47	0
7	CA	B	2	1/1	0.95	0.10	-1.63	56,56,56,56	0
7	CA	C	1	1/1	0.93	0.08	-1.70	61,61,61,61	0
7	CA	E	2	1/1	0.95	0.06	-1.96	58,58,58,58	0
7	CA	C	4	1/1	0.96	0.10	-2.06	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.