



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 16, 2018 – 03:35 AM EST

PDB ID : 3HM7  
Title : Crystal structure of allantoinase from *Bacillus halodurans* C-125  
Authors : Patskovsky, Y.; Romero, R.; Rutter, M.; Miller, S.; Wasserman, S.R.; Sauder, J.M.; Raushel, F.M.; Burley, S.K.; Almo, S.C.; New York Structural GenomiX Research Consortium (NYSGXRC); New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2009-05-28  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div> <div>7%</div> <div>82%</div> <div>14%</div> <div>..</div> </div>
1	B	448	<div> <div>5%</div> <div>82%</div> <div>15%</div> <div>..</div> </div>
1	C	448	<div> <div>27%</div> <div>78%</div> <div>18%</div> <div>..</div> </div>
1	D	448	<div> <div>7%</div> <div>81%</div> <div>15%</div> <div>..</div> </div>
1	E	448	<div> <div>3%</div> <div>87%</div> <div>10%</div> <div>..</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	448	 A horizontal bar chart showing the quality of chain F. The bar is divided into four segments: a small red segment at the beginning labeled '2%', a large green segment labeled '84%', a yellow segment labeled '13%', and a small grey segment at the end. Two dots '• •' are located at the far right end of the bar.

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 20901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Allantoinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	4	0
			3425	2166	593	649	17			
1	B	437	Total	C	N	O	S	0	5	0
			3437	2173	594	653	17			
1	C	435	Total	C	N	O	S	0	3	0
			3410	2154	588	651	17			
1	D	436	Total	C	N	O	S	0	2	0
			3415	2157	591	650	17			
1	E	436	Total	C	N	O	S	0	2	0
			3412	2156	590	649	17			
1	F	436	Total	C	N	O	S	0	4	0
			3425	2165	592	651	17			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
A	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
A	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
A	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
A	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
A	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
A	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
B	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
B	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
B	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
B	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
B	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
B	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
C	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
C	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
C	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
C	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
C	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
C	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
D	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
D	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
D	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
D	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
D	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
D	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
E	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
E	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
E	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8
E	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
E	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
E	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	-1	MET	-	EXPRESSION TAG	UNP Q9KAH8
F	0	SER	-	EXPRESSION TAG	UNP Q9KAH8
F	1	LEU	-	EXPRESSION TAG	UNP Q9KAH8
F	439	GLU	-	EXPRESSION TAG	UNP Q9KAH8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	440	GLY	-	EXPRESSION TAG	UNP Q9KAH8
F	441	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	442	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	443	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	444	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	445	HIS	-	EXPRESSION TAG	UNP Q9KAH8
F	446	HIS	-	EXPRESSION TAG	UNP Q9KAH8

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Zn 1	0	0
2	E	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0
2	A	1	Total 1	Zn 1	0	0
2	F	1	Total 1	Zn 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	53	Total 53	O 53	0	0
3	B	69	Total 69	O 69	0	0
3	C	39	Total 39	O 39	0	0
3	D	47	Total 47	O 47	0	0
3	E	74	Total 74	O 74	0	0
3	F	89	Total 89	O 89	0	0




- Molecule 1: Allantoinase

Chain D:

● Molecule 1: Allantoinase

Chain E:

Residue	Conservation
Met	Conserved
Ser	Conserved
Leu	Conserved
Lys	Conserved
R3	Conserved
V14	Conserved
T15	Conserved
E16	Conserved
T17	Conserved
T18	Conserved
D43	Conserved
G50	Conserved
L53	Conserved
P57	Conserved
H65	Conserved
F66	Conserved
N67	Conserved
P96	Conserved
L97	Conserved
N98	Conserved
S99	Conserved
T105	Conserved
R113	Conserved
K119	Conserved
N134	Conserved
I135	Conserved
G144	Conserved
K150	Conserved
G157	Conserved
T158	Conserved
D159	Conserved
D160	Conserved
L169	Conserved
M173	Conserved
K174	Conserved
K175	Conserved
S181	Conserved
Q204	Conserved
S219	Conserved
G241	Conserved
R242	Conserved
L272	Conserved
L275	Conserved
L284	Conserved
Q295	Conserved
H316	Conserved
Q322	Conserved
K323	Conserved
K324	Conserved
T325	Conserved
G338	Conserved
C339	Conserved
M346	Conserved
L365	Conserved
I382	Conserved
Q383	Conserved
F390	Conserved
L401	Conserved
M402	Conserved
A403	Conserved
S404	Conserved
D405	Conserved
L406	Conserved
T427	Conserved
V434	Conserved
Y435	Conserved
H438	Conserved
GLU	Conserved
GLY	Conserved
HIS	Conserved
HIS	Conserved
HIS	Conserved
HIS	Conserved
HIS	Conserved





H433	H437	GLU
Y434		GLY
Y435		HIS
Q436		HIS
	H438	HIS
		HIS
		HIS
		HIS
		HIS

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.66Å 157.66Å 418.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.60 27.39 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (20.00-2.60) 100.0 (27.39-2.60)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.94 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, $R_{free}$	0.240 , 0.267 0.240 , 0.267	Depositor DCC
$R_{free}$ test set	2847 reflections (3.10%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 50.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	20901	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 69.84 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.4571e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3507	0.60	0/4749
1	B	0.44	0/3523	0.61	1/4770 (0.0%)
1	C	0.43	0/3485	0.62	0/4720
1	D	0.42	0/3488	0.59	0/4724
1	E	0.43	0/3488	0.59	0/4724
1	F	0.41	0/3508	0.61	1/4752 (0.0%)
All	All	0.42	0/20999	0.60	2/28439 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	395	LEU	CA-CB-CG	6.19	129.54	115.30
1	F	262	VAL	CB-CA-C	-5.13	101.65	111.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	159	ASP	Peptide
1	A	160	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3425	0	3404	27	0
1	B	3437	0	3411	25	0
1	C	3410	0	3376	32	0
1	D	3415	0	3379	29	0
1	E	3412	0	3380	16	0
1	F	3425	0	3393	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	53	0	0	0	0
3	B	69	0	0	0	0
3	C	39	0	0	0	0
3	D	47	0	0	0	0
3	E	74	0	0	0	0
3	F	89	0	0	0	0
All	All	20901	0	20343	138	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:67:ASN:ND2	1:D:98:ASN:HB2	1.96	0.80
1:F:75:GLU:HG2	1:F:318:PRO:HG3	1.69	0.73
1:B:75:GLU:HG2	1:B:318:PRO:HG3	1.76	0.68
1:E:67:ASN:ND2	1:E:98:ASN:HB2	2.09	0.67
1:F:67:ASN:ND2	1:F:98:ASN:HB2	2.09	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/448 (98%)	410 (94%)	25 (6%)	3 (1%)	25	49
1	B	440/448 (98%)	425 (97%)	13 (3%)	2 (0%)	32	58
1	C	436/448 (97%)	397 (91%)	35 (8%)	4 (1%)	20	40
1	D	436/448 (97%)	416 (95%)	18 (4%)	2 (0%)	32	58
1	E	436/448 (97%)	419 (96%)	15 (3%)	2 (0%)	32	58
1	F	438/448 (98%)	424 (97%)	12 (3%)	2 (0%)	32	58
All	All	2624/2688 (98%)	2491 (95%)	118 (4%)	15 (1%)	28	53

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132	PRO
1	A	242	HIS
1	B	242	HIS
1	C	242	HIS
1	D	242	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/382 (98%)	343 (92%)	32 (8%)	12	24
1	B	377/382 (99%)	351 (93%)	26 (7%)	18	36
1	C	373/382 (98%)	336 (90%)	37 (10%)	9	17

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	373/382 (98%)	340 (91%)	33 (9%)	12	22
1	E	373/382 (98%)	350 (94%)	23 (6%)	21	42
1	F	375/382 (98%)	347 (92%)	28 (8%)	16	31
All	All	2246/2292 (98%)	2067 (92%)	179 (8%)	14	27

5 of 179 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	304	LEU
1	D	113	ARG
1	F	242	HIS
1	C	328	THR
1	C	393	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	418	GLN
1	D	352	HIS
1	F	383	GLN
1	D	162	GLN
1	B	67	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	436/448 (97%)	0.29	32 (7%) 16 11	22, 38, 54, 77	0
1	B	437/448 (97%)	0.09	21 (4%) 31 24	28, 37, 52, 66	0
1	C	435/448 (97%)	1.48	123 (28%) 1 0	22, 39, 51, 74	0
1	D	436/448 (97%)	0.22	31 (7%) 17 12	26, 38, 53, 70	0
1	E	436/448 (97%)	-0.02	12 (2%) 53 46	25, 37, 47, 68	0
1	F	436/448 (97%)	-0.06	8 (1%) 69 63	26, 37, 48, 64	0
All	All	2616/2688 (97%)	0.33	227 (8%) 11 7	22, 37, 52, 77	0

The worst 5 of 227 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	158	THR	9.3
1	C	402	ASN	7.5
1	A	157	GLY	7.4
1	C	156	CYS	6.9
1	C	144	GLY	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	D	447	1/1	0.83	0.20	-0.80	40,40,40,40	1
2	ZN	C	447	1/1	0.93	0.21	-1.82	50,50,50,50	1
2	ZN	B	447	1/1	0.96	0.08	-2.84	39,39,39,39	1
2	ZN	E	447	1/1	0.97	0.08	-4.25	39,39,39,39	1
2	ZN	A	447	1/1	0.97	0.06	-	43,43,43,43	1
2	ZN	F	447	1/1	0.94	0.22	-	39,39,39,39	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.