



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2017 – 08:33 AM EDT

PDB ID : 3HM8
Title : Crystal structure of the C-terminal Hexokinase domain of human HK3
Authors : Nedyalkova, L.; Tong, Y.; Rabeh, W.; Tempel, W.; Landry, R.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Weigelt, J.; Bochkarev, A.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : unknown
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	445	<div> <div></div> <div>85%14%.</div> </div>
1	B	445	<div> <div>%</div> <div>88%11%.</div> </div>
1	C	445	<div> <div>%</div> <div>85%13%.</div> </div>
1	D	445	<div> <div>2%</div> <div>87%11%.</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hexokinase-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	441	Total	C	N	O	S	0	0	0
			3206	2012	555	617	22			
1	B	441	Total	C	N	O	S	0	0	0
			3117	1959	546	590	22			
1	C	436	Total	C	N	O	S	0	0	0
			3116	1962	537	596	21			
1	D	437	Total	C	N	O	S	0	0	0
			3082	1934	534	592	22			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	478	GLY	-	EXPRESSION TAG	UNP P52790
A	479	SER	-	EXPRESSION TAG	UNP P52790
B	478	GLY	-	EXPRESSION TAG	UNP P52790
B	479	SER	-	EXPRESSION TAG	UNP P52790
C	478	GLY	-	EXPRESSION TAG	UNP P52790
C	479	SER	-	EXPRESSION TAG	UNP P52790
D	478	GLY	-	EXPRESSION TAG	UNP P52790
D	479	SER	-	EXPRESSION TAG	UNP P52790

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		
2	C	1	Total	C	O	0	0
			12	6	6		
2	D	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is BETA-D-GLUCOSE-6-PHOSPHATE (three-letter code: BG6) (formula: $C_6H_{13}O_9P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			16	6	9	1		
3	B	1	Total	C	O	P	0	0
			16	6	9	1		
3	C	1	Total	C	O	P	0	0
			16	6	9	1		
3	D	1	Total	C	O	P	0	0
			16	6	9	1		



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	83.90 Å 83.90 Å 323.33 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 20.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	97.2 (20.00-2.80) 97.2 (20.00-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.79 Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.245 , 0.270 0.249 , 0.271	Depositor DCC
R_{free} test set	1502 reflections (2.92%)	DCC
Wilson B-factor (Å ²)	68.9	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 26.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.076 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12633	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, BG6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	1/3249 (0.0%)	0.57	0/4407
1	B	0.55	1/3160 (0.0%)	0.55	0/4295
1	C	0.55	0/3158	0.54	0/4288
1	D	0.52	0/3124	0.54	0/4250
All	All	0.56	2/12691 (0.0%)	0.55	0/17240

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	829	CYS	CB-SG	-6.89	1.70	1.82
1	B	829	CYS	CB-SG	-5.17	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3206	0	3098	48	0
1	B	3117	0	2950	32	0
1	C	3116	0	2967	40	0
1	D	3082	0	2884	31	0
2	A	12	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	12	0	12	0	0
2	C	12	0	12	0	0
2	D	12	0	12	0	0
3	A	16	0	11	0	0
3	B	16	0	11	0	0
3	C	16	0	11	0	0
3	D	16	0	11	0	0
All	All	12633	0	11991	145	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 145 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:702:VAL:HG13	1:A:703:PRO:HD2	1.61	0.83
1:B:702:VAL:HG13	1:B:703:PRO:HD2	1.61	0.83
1:C:702:VAL:HG13	1:C:703:PRO:HD2	1.61	0.81
1:D:702:VAL:HG13	1:D:703:PRO:HD2	1.62	0.79
1:D:562:THR:HG21	1:D:595:GLN:HE22	1.48	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	439/445 (99%)	424 (97%)	14 (3%)	1 (0%)	51 83
1	B	439/445 (99%)	423 (96%)	14 (3%)	2 (0%)	32 67
1	C	432/445 (97%)	420 (97%)	11 (2%)	1 (0%)	51 83
1	D	433/445 (97%)	419 (97%)	13 (3%)	1 (0%)	51 83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	1743/1780 (98%)	1686 (97%)	52 (3%)	5 (0%)	44 77

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	774	PHE
1	A	530	PRO
1	B	530	PRO
1	C	530	PRO
1	D	530	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	321/365 (88%)	318 (99%)	3 (1%)	82 95
1	B	296/365 (81%)	293 (99%)	3 (1%)	80 95
1	C	301/365 (82%)	298 (99%)	3 (1%)	80 95
1	D	293/365 (80%)	290 (99%)	3 (1%)	80 95
All	All	1211/1460 (83%)	1199 (99%)	12 (1%)	80 95

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	898	GLN
1	C	606	PHE
1	D	606	PHE
1	B	715	TRP
1	C	898	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	574	GLN
1	B	698	ASN
1	C	698	ASN
1	A	698	ASN
1	C	574	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	A	1001	-	12,12,12	1.32	3 (25%)	17,17,17	1.04	0
3	BG6	A	1002	-	16,16,16	1.55	3 (18%)	24,24,24	0.91	2 (8%)
2	GLC	B	1001	-	12,12,12	1.22	2 (16%)	17,17,17	0.88	0
3	BG6	B	1002	-	16,16,16	1.59	3 (18%)	24,24,24	0.99	2 (8%)
2	GLC	C	1001	-	12,12,12	1.04	1 (8%)	17,17,17	0.82	0
3	BG6	C	1002	-	16,16,16	1.61	3 (18%)	24,24,24	0.95	2 (8%)
2	GLC	D	1001	-	12,12,12	1.07	1 (8%)	17,17,17	0.72	0
3	BG6	D	1002	-	16,16,16	1.74	5 (31%)	24,24,24	1.02	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	1001	-	-	0/2/22/22	0/1/1/1
3	BG6	A	1002	-	-	0/6/26/26	0/1/1/1
2	GLC	B	1001	-	-	0/2/22/22	0/1/1/1
3	BG6	B	1002	-	-	0/6/26/26	0/1/1/1
2	GLC	C	1001	-	-	0/2/22/22	0/1/1/1
3	BG6	C	1002	-	-	0/6/26/26	0/1/1/1
2	GLC	D	1001	-	-	0/2/22/22	0/1/1/1
3	BG6	D	1002	-	-	0/6/26/26	0/1/1/1

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1002	BG6	P-O2P	2.01	1.63	1.54
3	D	1002	BG6	O5-C5	2.02	1.49	1.44
3	D	1002	BG6	C4-C3	2.05	1.57	1.52
3	D	1002	BG6	P-O2P	2.15	1.63	1.54
2	B	1001	GLC	O5-C1	2.15	1.47	1.43

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1002	BG6	O5-C5-C6	2.06	110.75	106.64
3	C	1002	BG6	O1P-P-O6	2.35	112.99	106.73
3	B	1002	BG6	O1P-P-O6	2.38	113.08	106.73
3	B	1002	BG6	O5-C5-C6	2.46	111.56	106.64
3	A	1002	BG6	O1P-P-O6	2.53	113.45	106.73

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	441/445 (99%)	-0.23	0 100 100	54, 69, 98, 117	0
1	B	441/445 (99%)	-0.11	4 (0%) 84 79	61, 89, 128, 148	0
1	C	436/445 (97%)	-0.09	6 (1%) 75 69	58, 82, 141, 162	0
1	D	437/445 (98%)	-0.13	9 (2%) 64 54	69, 91, 138, 151	0
All	All	1755/1780 (98%)	-0.14	19 (1%) 80 74	54, 83, 131, 162	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	599	GLY	4.5
1	D	598	SER	4.3
1	C	770	LEU	3.6
1	D	486	THR	3.2
1	D	774	PHE	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	BG6	D	1002	16/16	0.94	0.20	0.84	76,79,83,84	0
3	BG6	A	1002	16/16	0.96	0.17	0.46	58,60,65,65	0
2	GLC	C	1001	12/12	0.97	0.17	0.33	60,61,63,64	0
2	GLC	A	1001	12/12	0.96	0.18	0.29	54,55,56,57	0
2	GLC	B	1001	12/12	0.96	0.16	0.03	66,67,71,71	0
3	BG6	B	1002	16/16	0.95	0.17	-0.05	65,69,73,74	0
2	GLC	D	1001	12/12	0.96	0.15	-0.24	69,71,72,72	0
3	BG6	C	1002	16/16	0.96	0.15	-0.35	69,73,76,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.