



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2017 – 06:26 PM EDT

PDB ID : 2HN9
Title : Crystal structure of the uridine phosphorylase from *Salmonella typhimurium* in complex with thymine and phosphate ion at 2.12Å resolution
Authors : Gabdoulkhakov, A.G.; Timofeev, V.I.; Mikhailov, A.M.
Deposited on : unknown
Resolution : 2.12 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4988 (2.14-2.10)
Clashscore	112137	5557 (2.14-2.10)
Ramachandran outliers	110173	5504 (2.14-2.10)
Sidechain outliers	110143	5505 (2.14-2.10)
RSRZ outliers	101464	5021 (2.14-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	<div> <div>5%</div> <div> <div></div> <div>88%</div> <div>10%</div> <div>.</div> </div> </div>
1	B	253	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>.</div> </div> </div>
1	C	253	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div>.</div> </div> </div>
1	D	253	<div> <div>6%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div>..</div> </div> </div>
1	E	253	<div> <div>4%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	253	 87%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	A	1254	-	-	-	X
3	TDR	F	2003	-	-	-	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 11492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			1900	1188	334	365	13			
1	B	243	Total	C	N	O	S	0	0	0
			1819	1140	321	347	11			
1	C	243	Total	C	N	O	S	0	0	0
			1824	1143	322	347	12			
1	D	247	Total	C	N	O	S	0	0	0
			1856	1162	327	355	12			
1	E	247	Total	C	N	O	S	0	0	0
			1857	1163	328	355	11			
1	F	242	Total	C	N	O	S	0	0	0
			1815	1139	320	345	11			

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is THYMINE (three-letter code: TDR) (formula: $C_5H_6N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	5	2	2		
3	B	1	Total	C	N	O	0	0
			9	5	2	2		
3	D	1	Total	C	N	O	0	0
			9	5	2	2		
3	F	1	Total	C	N	O	0	0
			9	5	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	77	Total	O	0	0
			77	77		
4	B	63	Total	O	0	0
			63	63		
4	C	52	Total	O	0	0
			52	52		
4	D	58	Total	O	0	0
			58	58		
4	E	63	Total	O	0	0
			63	63		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	67	Total	O	0	0
			67	67		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase




- Molecule 1: Uridine phosphorylase



● Molecule 1: Uridine phosphorylase



● Molecule 1: Uridine phosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	88.80Å 123.69Å 134.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.12 12.00 – 2.12	Depositor EDS
% Data completeness (in resolution range)	100.0 (10.00-2.12) 100.0 (12.00-2.12)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.03 (at 2.13Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.196 , 0.226 0.195 , 0.226	Depositor DCC
R_{free} test set	4176 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	12.9	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.46 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11492	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TDR, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1930	0.51	0/2613
1	B	0.33	0/1848	0.50	0/2504
1	C	0.32	0/1852	0.49	0/2505
1	D	0.32	0/1885	0.49	0/2552
1	E	0.33	0/1886	0.49	0/2553
1	F	0.32	0/1844	0.50	0/2497
All	All	0.33	0/11245	0.50	0/15224

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1900	0	1914	23	0
1	B	1819	0	1830	10	0
1	C	1824	0	1843	10	0
1	D	1856	0	1868	17	0
1	E	1857	0	1872	10	0
1	F	1815	0	1832	19	0
2	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	9	0	6	1	0
3	B	9	0	6	0	0
3	D	9	0	6	0	0
3	F	9	0	6	0	0
4	A	77	0	0	0	0
4	B	63	0	0	0	0
4	C	52	0	0	0	0
4	D	58	0	0	0	0
4	E	63	0	0	0	0
4	F	67	0	0	0	0
All	All	11492	0	11183	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1016:LEU:HG	1:A:1063:ILE:HD11	1.27	1.14
1:D:3016:LEU:HG	1:D:3063:ILE:HD11	1.46	0.96
1:B:6016:LEU:HG	1:B:6063:ILE:HD11	1.47	0.93
1:F:4063:ILE:HD13	1:F:4086:ILE:HD11	1.55	0.88
1:A:1016:LEU:HG	1:A:1063:ILE:CD1	2.07	0.83
1:A:1162:PHE:HB2	1:F:4076:ILE:HD11	1.68	0.76
1:A:1099:GLN:HB2	1:A:1102:ILE:HD13	1.70	0.74
1:A:1162:PHE:CB	1:F:4076:ILE:HD11	2.18	0.72
1:B:6016:LEU:HG	1:B:6063:ILE:CD1	2.25	0.63
1:D:3240:HIS:O	1:D:3244:ILE:HG12	1.97	0.63
1:A:1049:GLU:HB3	1:F:4049:GLU:HB3	1.80	0.63
1:B:6063:ILE:HD13	1:B:6086:ILE:HD11	1.82	0.61
1:D:3144:ALA:HA	1:D:3244:ILE:HD12	1.82	0.60
1:F:4022:ALA:HB2	1:F:4086:ILE:HD13	1.83	0.60
1:A:1158:SER:HB3	1:A:1200:ALA:HB2	1.82	0.60
1:D:3158:SER:HB3	1:D:3200:ALA:HB2	1.82	0.59
1:C:2178:ARG:HA	1:C:2181:LYS:HE3	1.84	0.59
1:A:1220:ILE:HD11	3:A:2001:TDR:H5M1	1.85	0.59
1:E:5081:LEU:HB3	1:E:5086:ILE:HD13	1.84	0.58
1:F:4158:SER:HB3	1:F:4200:ALA:HB2	1.85	0.58
1:C:2038:MET:HG2	1:C:2057:LEU:HD13	1.84	0.57
1:F:4038:MET:HG2	1:F:4057:LEU:HD13	1.87	0.57
1:D:3016:LEU:HG	1:D:3063:ILE:CD1	2.28	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:5158:SER:HB3	1:E:5200:ALA:HB2	1.86	0.56
1:F:4081:LEU:HB3	1:F:4086:ILE:HD12	1.86	0.56
1:A:1038:MET:HG2	1:A:1057:LEU:HD13	1.86	0.56
1:D:3038:MET:HG2	1:D:3057:LEU:HD13	1.88	0.56
1:F:4016:LEU:HG	1:F:4063:ILE:HD11	1.87	0.55
1:A:1162:PHE:HB3	1:F:4076:ILE:HD11	1.89	0.55
1:D:3099:GLN:HB2	1:D:3102:ILE:HD12	1.88	0.55
1:C:2158:SER:HB3	1:C:2200:ALA:HB2	1.89	0.55
1:E:5144:ALA:HA	1:E:5244:ILE:HD12	1.89	0.54
1:B:6158:SER:HB3	1:B:6200:ALA:HB2	1.88	0.54
1:B:6176:VAL:O	1:B:6181:LYS:HE2	2.08	0.54
1:E:5016:LEU:HD21	1:E:5086:ILE:CD1	2.38	0.53
1:A:1162:PHE:HB3	1:F:4076:ILE:CD1	2.38	0.53
1:A:1060:LYS:HD2	1:A:1253:LEU:HB3	1.91	0.53
1:F:4092:ILE:HD12	1:F:4242:VAL:HA	1.91	0.53
1:E:5016:LEU:HD21	1:E:5086:ILE:HD12	1.93	0.51
1:B:6049:GLU:HB3	1:D:3049:GLU:HB3	1.93	0.51
1:D:3081:LEU:HB3	1:D:3086:ILE:HD13	1.92	0.51
1:C:2038:MET:SD	1:C:2062:VAL:HG21	2.51	0.50
1:A:1044:LEU:HD11	1:A:1054:ARG:HB2	1.93	0.49
1:D:3099:GLN:HB2	1:D:3102:ILE:CD1	2.43	0.49
1:A:1162:PHE:CB	1:F:4076:ILE:CD1	2.91	0.48
1:E:5147:ILE:HG12	1:E:5244:ILE:HD11	1.95	0.48
1:D:3147:ILE:HG12	1:D:3244:ILE:CD1	2.43	0.47
1:A:1072:PRO:O	1:A:1076:ILE:HD12	2.14	0.47
1:C:2140:LEU:HD22	1:C:2216:VAL:HB	1.96	0.47
1:F:4092:ILE:CD1	1:F:4242:VAL:HA	2.45	0.47
1:A:1063:ILE:CD1	1:A:1086:ILE:HD11	2.45	0.47
1:E:5168:ARG:NH1	1:E:5227:GLU:O	2.48	0.46
1:A:1232:GLU:HB2	1:A:1235:LYS:HB2	1.98	0.46
1:D:3147:ILE:HG12	1:D:3244:ILE:HD11	1.98	0.46
1:A:1063:ILE:HD13	1:A:1086:ILE:HD11	1.98	0.46
1:C:2034:ILE:HG13	1:C:2242:VAL:HG13	1.97	0.46
1:D:3057:LEU:HB3	1:D:3253:LEU:HD11	1.97	0.45
1:B:6081:LEU:HB3	1:B:6086:ILE:HD13	1.98	0.45
1:A:1081:LEU:HB3	1:A:1086:ILE:HD12	1.99	0.45
1:A:1163:TYR:HB2	1:A:1164:PRO:CD	2.47	0.45
1:F:4163:TYR:HB2	1:F:4164:PRO:CD	2.47	0.44
1:D:3092:ILE:HD11	1:D:3241:ALA:HB1	1.99	0.44
1:B:6038:MET:HG2	1:B:6057:LEU:HD13	2.00	0.44
1:B:6108:LEU:HD22	1:B:6152:HIS:HB2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:5147:ILE:HG12	1:E:5244:ILE:CD1	2.48	0.44
1:F:4102:ILE:O	1:F:4222:ASN:ND2	2.49	0.43
1:F:4027:ASP:HB3	1:F:4030:ARG:HG2	1.99	0.43
1:A:1163:TYR:HB2	1:A:1164:PRO:HD3	2.01	0.43
1:C:2107:VAL:HG23	1:C:2151:THR:HG23	2.00	0.42
1:C:2234:MET:HB3	1:C:2235:LYS:H	1.66	0.42
1:E:5091:ARG:HB3	1:E:5215:MET:HG2	2.00	0.42
1:E:5044:LEU:HD11	1:E:5054:ARG:HB2	2.01	0.42
1:A:1230:ASN:O	1:A:1232:GLU:N	2.53	0.41
1:A:1104:VAL:HG11	1:A:1229:PRO:HG3	2.02	0.41
1:D:3163:TYR:HB2	1:D:3164:PRO:CD	2.50	0.41
1:F:4016:LEU:HG	1:F:4063:ILE:CD1	2.50	0.41
1:B:6089:PHE:O	1:B:6213:ALA:HA	2.21	0.41
1:D:3027:ASP:HA	1:D:3028:PRO:HD2	1.86	0.41
1:C:2163:TYR:HB2	1:C:2164:PRO:HD3	2.03	0.41
1:C:2094:THR:HB	1:C:2220:ILE:HG12	2.02	0.41
1:D:3063:ILE:CD1	1:D:3086:ILE:HD11	2.51	0.40
1:F:4049:GLU:HG3	1:F:4068:GLY:HA3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/253 (99%)	242 (96%)	5 (2%)	4 (2%)	11	5
1	B	239/253 (94%)	236 (99%)	2 (1%)	1 (0%)	38	35
1	C	239/253 (94%)	235 (98%)	3 (1%)	1 (0%)	38	35
1	D	243/253 (96%)	240 (99%)	2 (1%)	1 (0%)	38	35
1	E	243/253 (96%)	239 (98%)	3 (1%)	1 (0%)	38	35
1	F	238/253 (94%)	233 (98%)	4 (2%)	1 (0%)	38	35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1453/1518 (96%)	1425 (98%)	19 (1%)	9 (1%)	28	23

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1163	TYR
1	A	1231	ALA
1	A	1235	LYS
1	E	5163	TYR
1	F	4163	TYR
1	B	6163	TYR
1	C	2163	TYR
1	D	3163	TYR
1	A	1232	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/202 (100%)	197 (98%)	5 (2%)	53	56
1	B	192/202 (95%)	191 (100%)	1 (0%)	91	94
1	C	193/202 (96%)	189 (98%)	4 (2%)	59	63
1	D	197/202 (98%)	195 (99%)	2 (1%)	80	84
1	E	197/202 (98%)	193 (98%)	4 (2%)	60	65
1	F	192/202 (95%)	191 (100%)	1 (0%)	91	94
All	All	1173/1212 (97%)	1156 (99%)	17 (1%)	71	77

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1170	ASP
1	A	1196	GLU
1	A	1230	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1232	GLU
1	A	1236	GLN
1	B	6196	GLU
1	C	2084	LEU
1	C	2092	ILE
1	C	2185	GLU
1	C	2196	GLU
1	D	3147	ILE
1	D	3196	GLU
1	E	5029	GLU
1	E	5030	ARG
1	E	5196	GLU
1	E	5215	MET
1	F	4196	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1103	ASN
1	E	5188	GLN
1	E	5225	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	1254	-	4,4,4	0.73	0	6,6,6	0.37	0
3	TDR	A	2001	-	7,9,9	0.90	0	7,12,12	5.74	5 (71%)
3	TDR	B	2002	-	7,9,9	0.93	0	7,12,12	5.73	5 (71%)
3	TDR	D	2004	-	7,9,9	0.91	0	7,12,12	5.71	5 (71%)
3	TDR	F	2003	-	7,9,9	0.90	0	7,12,12	5.72	5 (71%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PO4	A	1254	-	-	0/0/0/0	0/0/0/0
3	TDR	A	2001	-	-	0/0/0/0	0/1/1/1
3	TDR	B	2002	-	-	0/0/0/0	0/1/1/1
3	TDR	D	2004	-	-	0/0/0/0	0/1/1/1
3	TDR	F	2003	-	-	0/0/0/0	0/1/1/1

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2004	TDR	N1-C2-N3	-9.25	121.75	128.40
3	F	2003	TDR	N1-C2-N3	-9.16	121.81	128.40
3	B	2002	TDR	N1-C2-N3	-9.15	121.82	128.40
3	A	2001	TDR	N1-C2-N3	-8.89	122.00	128.40
3	A	2001	TDR	C5-C6-N1	-7.46	118.95	125.26
3	D	2004	TDR	C5-C6-N1	-7.14	119.22	125.26
3	F	2003	TDR	C5-C6-N1	-7.03	119.32	125.26
3	B	2002	TDR	C5-C6-N1	-6.94	119.40	125.26
3	A	2001	TDR	C5-C4-N3	-6.30	118.29	125.24
3	B	2002	TDR	C5-C4-N3	-6.16	118.45	125.24
3	F	2003	TDR	C5-C4-N3	-6.05	118.57	125.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2004	TDR	C5-C4-N3	-5.81	118.83	125.24
3	B	2002	TDR	C6-N1-C2	4.15	122.00	115.36
3	A	2001	TDR	C6-N1-C2	4.19	122.07	115.36
3	F	2003	TDR	C6-N1-C2	4.21	122.10	115.36
3	D	2004	TDR	C6-N1-C2	4.29	122.22	115.36
3	D	2004	TDR	C4-N3-C2	6.16	120.55	115.16
3	A	2001	TDR	C4-N3-C2	6.17	120.56	115.16
3	F	2003	TDR	C4-N3-C2	6.29	120.66	115.16
3	B	2002	TDR	C4-N3-C2	6.43	120.79	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2001	TDR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/253 (100%)	-0.07	12 (4%) 32 38	3, 8, 28, 42	0
1	B	243/253 (96%)	-0.27	5 (2%) 64 68	2, 7, 17, 22	1 (0%)
1	C	243/253 (96%)	0.04	9 (3%) 42 49	3, 13, 27, 44	0
1	D	247/253 (97%)	0.00	14 (5%) 24 30	3, 10, 25, 54	0
1	E	247/253 (97%)	-0.09	10 (4%) 39 45	3, 8, 22, 49	0
1	F	242/253 (95%)	-0.33	3 (1%) 79 82	2, 7, 18, 21	0
All	All	1475/1518 (97%)	-0.12	53 (3%) 43 50	2, 8, 24, 54	1 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1234	MET	9.8
1	E	5235	LYS	9.1
1	D	3237	THR	8.5
1	E	5237	THR	7.9
1	D	3234	MET	7.7
1	E	5236	GLN	7.0
1	A	1231	ALA	5.9
1	B	6231	ALA	5.7
1	A	1233	THR	5.6
1	D	3236	GLN	5.4
1	D	3228	ILE	5.1
1	E	5230	ASN	4.8
1	D	3235	LYS	4.6
1	A	1001	MET	4.5
1	D	3230	ASN	4.5
1	C	2236	GLN	4.4
1	A	1235	LYS	4.1
1	E	5238	GLU	4.0
1	F	4007	PHE	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	3238	GLU	3.7
1	D	3005	ASP	3.6
1	C	2235	LYS	3.6
1	A	1002	SER	3.5
1	A	1230	ASN	3.4
1	A	1101	HIS	3.1
1	E	5003	LYS	3.1
1	B	6101	HIS	2.9
1	A	1236	GLN	2.9
1	C	2145	LYS	2.9
1	C	2148	GLY	2.8
1	C	2234	MET	2.7
1	B	6239	SER	2.7
1	E	5007	PHE	2.6
1	B	6145	LYS	2.6
1	D	3226	GLN	2.6
1	C	2104	VAL	2.5
1	A	1039	ASP	2.5
1	C	2094	THR	2.5
1	E	5239	SER	2.4
1	D	3225	GLN	2.4
1	A	1232	GLU	2.3
1	C	2003	LYS	2.2
1	D	3004	SER	2.2
1	D	3170	ASP	2.2
1	D	3101	HIS	2.1
1	E	5243	LYS	2.1
1	F	4033	LYS	2.1
1	F	4030	ARG	2.1
1	B	6253	LEU	2.1
1	D	3224	THR	2.1
1	A	1240	HIS	2.1
1	E	5253	LEU	2.0
1	C	2240	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	PO4	A	1254	5/5	0.94	0.23	3.93	27,27,27,27	0
3	TDR	F	2003	9/9	0.88	0.16	2.90	17,17,17,17	0
3	TDR	D	2004	9/9	0.89	0.13	0.72	18,18,18,18	0
3	TDR	B	2002	9/9	0.87	0.12	0.61	12,12,13,13	0
3	TDR	A	2001	9/9	0.94	0.11	-0.09	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.