



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 08:56 pm GMT

PDB ID : 3HN2  
Title : Crystal structure of 2-dehydropantoate 2-reductase FROM *Geobacter metal-*  
lireducens GS-15  
Authors : Patskovsky, Y.; Toro, R.; Morano, C.; Rutter, M.; Chang, S.; Sauder, J.M.;  
Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Ge-  
nomics (NYSGXRC)  
Deposited on : 2009-05-29  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	 5% 82% 13% . .
1	B	312	 12% 84% 11% . .
1	C	312	 6% 79% 15% . . .
1	D	312	 20% 81% 14% . . .

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-dehydropantoate 2-reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	302	Total	C	N	O	S	0	0	0
			2327	1476	409	431	11			
1	B	302	Total	C	N	O	S	0	1	0
			2333	1482	411	429	11			
1	C	302	Total	C	N	O	S	0	0	0
			2326	1477	409	429	11			
1	D	301	Total	C	N	O	S	0	0	0
			2322	1475	408	428	11			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q39SB2
A	0	SER	-	EXPRESSION TAG	UNP Q39SB2
A	1	LEU	-	EXPRESSION TAG	UNP Q39SB2
A	303	GLU	-	EXPRESSION TAG	UNP Q39SB2
A	304	GLY	-	EXPRESSION TAG	UNP Q39SB2
A	305	HIS	-	EXPRESSION TAG	UNP Q39SB2
A	306	HIS	-	EXPRESSION TAG	UNP Q39SB2
A	307	HIS	-	EXPRESSION TAG	UNP Q39SB2
A	308	HIS	-	EXPRESSION TAG	UNP Q39SB2
A	309	HIS	-	EXPRESSION TAG	UNP Q39SB2
A	310	HIS	-	EXPRESSION TAG	UNP Q39SB2
B	-1	MET	-	EXPRESSION TAG	UNP Q39SB2
B	0	SER	-	EXPRESSION TAG	UNP Q39SB2
B	1	LEU	-	EXPRESSION TAG	UNP Q39SB2
B	303	GLU	-	EXPRESSION TAG	UNP Q39SB2
B	304	GLY	-	EXPRESSION TAG	UNP Q39SB2
B	305	HIS	-	EXPRESSION TAG	UNP Q39SB2
B	306	HIS	-	EXPRESSION TAG	UNP Q39SB2
B	307	HIS	-	EXPRESSION TAG	UNP Q39SB2
B	308	HIS	-	EXPRESSION TAG	UNP Q39SB2
B	309	HIS	-	EXPRESSION TAG	UNP Q39SB2

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	310	HIS	-	EXPRESSION TAG	UNP Q39SB2
C	-1	MET	-	EXPRESSION TAG	UNP Q39SB2
C	0	SER	-	EXPRESSION TAG	UNP Q39SB2
C	1	LEU	-	EXPRESSION TAG	UNP Q39SB2
C	303	GLU	-	EXPRESSION TAG	UNP Q39SB2
C	304	GLY	-	EXPRESSION TAG	UNP Q39SB2
C	305	HIS	-	EXPRESSION TAG	UNP Q39SB2
C	306	HIS	-	EXPRESSION TAG	UNP Q39SB2
C	307	HIS	-	EXPRESSION TAG	UNP Q39SB2
C	308	HIS	-	EXPRESSION TAG	UNP Q39SB2
C	309	HIS	-	EXPRESSION TAG	UNP Q39SB2
C	310	HIS	-	EXPRESSION TAG	UNP Q39SB2
D	-1	MET	-	EXPRESSION TAG	UNP Q39SB2
D	0	SER	-	EXPRESSION TAG	UNP Q39SB2
D	1	LEU	-	EXPRESSION TAG	UNP Q39SB2
D	303	GLU	-	EXPRESSION TAG	UNP Q39SB2
D	304	GLY	-	EXPRESSION TAG	UNP Q39SB2
D	305	HIS	-	EXPRESSION TAG	UNP Q39SB2
D	306	HIS	-	EXPRESSION TAG	UNP Q39SB2
D	307	HIS	-	EXPRESSION TAG	UNP Q39SB2
D	308	HIS	-	EXPRESSION TAG	UNP Q39SB2
D	309	HIS	-	EXPRESSION TAG	UNP Q39SB2
D	310	HIS	-	EXPRESSION TAG	UNP Q39SB2

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total O 3 3	0	0
2	C	4	Total O 4 4	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.32Å 82.18Å 201.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 38.04 – 2.50	Depositor EDS
% Data completeness (in resolution range)	98.0 (20.00-2.50) 97.8 (38.04-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.55 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, $R_{free}$	0.243 , 0.292 0.243 , 0.292	Depositor DCC
$R_{free}$ test set	1377 reflections (3.21%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.4	Xtriage
Anisotropy	0.382	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 60.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9315	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/2366	0.64	1/3198 (0.0%)
1	B	0.35	0/2376	0.59	0/3212
1	C	0.36	0/2365	0.59	0/3197
1	D	0.34	0/2361	0.59	0/3192
All	All	0.36	0/9468	0.60	1/12799 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	4
1	C	0	3
1	D	0	2
All	All	0	9

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	152	ARG	O-C-N	-5.34	114.16	122.70

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	130	ARG	Peptide
1	B	131	GLY	Peptide
1	B	150	LEU	Peptide
1	B	152	ARG	Peptide
1	C	150	LEU	Peptide



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2327	0	2345	15	0
1	B	2333	0	2360	7	0
1	C	2326	0	2353	21	0
1	D	2322	0	2350	15	0
2	A	3	0	0	0	0
2	C	4	0	0	0	0
All	All	9315	0	9408	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 55 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:154:THR:HG23	1:C:155:GLY:H	1.40	0.86
1:A:152:ARG:O	1:A:153:ASP:HB2	1.86	0.75
1:C:155:GLY:HA2	1:C:158:GLU:HB2	1.72	0.69
1:D:150:LEU:HD22	1:D:151:PRO:HD3	1.75	0.68
1:C:148:GLU:HG3	1:C:151:PRO:HA	1.80	0.64

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	300/312 (96%)	277 (92%)	20 (7%)	3 (1%)	18 32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	301/312 (96%)	285 (95%)	15 (5%)	1 (0%)	44	66
1	C	300/312 (96%)	281 (94%)	16 (5%)	3 (1%)	18	32
1	D	299/312 (96%)	280 (94%)	15 (5%)	4 (1%)	14	25
All	All	1200/1248 (96%)	1123 (94%)	66 (6%)	11 (1%)	20	36

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	133	PRO
1	A	153	ASP
1	B	133	PRO
1	C	133	PRO
1	D	133	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/245 (96%)	212 (90%)	24 (10%)	8	16
1	B	237/245 (97%)	210 (89%)	27 (11%)	7	12
1	C	236/245 (96%)	206 (87%)	30 (13%)	5	9
1	D	236/245 (96%)	212 (90%)	24 (10%)	8	16
All	All	945/980 (96%)	840 (89%)	105 (11%)	7	13

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	269	ILE
1	C	63	GLU
1	D	206	LEU
1	B	291	GLU
1	C	20	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	229	GLN
1	D	40	ASN
1	C	40	ASN
1	B	40	ASN
1	C	192	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	302/312 (96%)	0.45	15 (4%) 30 31	16, 56, 86, 117	0
1	B	302/312 (96%)	0.75	38 (12%) 4 3	26, 64, 97, 109	0
1	C	302/312 (96%)	0.54	18 (5%) 23 23	31, 61, 100, 120	0
1	D	301/312 (96%)	1.08	61 (20%) 1 1	34, 68, 113, 133	0
All	All	1207/1248 (96%)	0.71	132 (10%) 6 6	16, 62, 103, 133	0

The worst 5 of 132 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	131	GLY	9.9
1	D	35	GLU	9.2
1	D	42	LEU	9.0
1	D	152	ARG	8.0
1	B	155	GLY	6.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.