



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:02 am GMT

PDB ID : 5HNP  
Title : The structure of the kdo-capped saccharide binding subunit of the O-12 specific ABC transporter, Wzt  
Authors : Mallette, E.; Mann, E.; Whitfield, C.; Kimber, M.S.  
Deposited on : 2016-01-18  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 2%; height: 10px; background-color: red;"></div> <div style="width: 78%; height: 10px; background-color: green;"></div> <div style="width: 15%; height: 10px; background-color: yellow;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div> <div style="width: 2%; background-color: red;"></div> <div style="width: 80%; background-color: green;"></div> <div style="width: 5%; background-color: yellow;"></div> <div style="width: 13%; background-color: grey;"></div> </div> </div>
1	B	190	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 2%; height: 10px; background-color: red;"></div> <div style="width: 79%; height: 10px; background-color: green;"></div> <div style="width: 13%; height: 10px; background-color: yellow;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%;"></div> </div> <div> <div style="width: 2%; background-color: red;"></div> <div style="width: 81%; background-color: green;"></div> <div style="width: 6%; background-color: yellow;"></div> <div style="width: 13%; background-color: grey;"></div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CL	A	501	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5293 atoms, of which 2578 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	161	Total	C	H	N	O	S	0	1	0
			2524	784	1281	219	236	4			
1	B	166	Total	C	H	N	O	S	0	0	0
			2567	797	1297	223	246	4			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	254	MET	-	initiating methionine	UNP A0A0B5INH8
A	255	HIS	-	expression tag	UNP A0A0B5INH8
A	256	HIS	-	expression tag	UNP A0A0B5INH8
A	257	HIS	-	expression tag	UNP A0A0B5INH8
A	258	HIS	-	expression tag	UNP A0A0B5INH8
A	259	HIS	-	expression tag	UNP A0A0B5INH8
A	260	HIS	-	expression tag	UNP A0A0B5INH8
A	261	GLU	-	expression tag	UNP A0A0B5INH8
A	262	ASN	-	expression tag	UNP A0A0B5INH8
A	263	LEU	-	expression tag	UNP A0A0B5INH8
A	264	TYR	-	expression tag	UNP A0A0B5INH8
A	265	PHE	-	expression tag	UNP A0A0B5INH8
A	266	GLN	-	expression tag	UNP A0A0B5INH8
A	267	GLY	-	expression tag	UNP A0A0B5INH8
A	443	SER	-	expression tag	UNP A0A0B5INH8
B	254	MET	-	initiating methionine	UNP A0A0B5INH8
B	255	HIS	-	expression tag	UNP A0A0B5INH8
B	256	HIS	-	expression tag	UNP A0A0B5INH8
B	257	HIS	-	expression tag	UNP A0A0B5INH8
B	258	HIS	-	expression tag	UNP A0A0B5INH8
B	259	HIS	-	expression tag	UNP A0A0B5INH8
B	260	HIS	-	expression tag	UNP A0A0B5INH8
B	261	GLU	-	expression tag	UNP A0A0B5INH8
B	262	ASN	-	expression tag	UNP A0A0B5INH8
B	263	LEU	-	expression tag	UNP A0A0B5INH8

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	264	TYR	-	expression tag	UNP A0A0B5INH8
B	265	PHE	-	expression tag	UNP A0A0B5INH8
B	266	GLN	-	expression tag	UNP A0A0B5INH8
B	267	GLY	-	expression tag	UNP A0A0B5INH8
B	443	SER	-	expression tag	UNP A0A0B5INH8

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0



- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	103	Total O 103 103	0	0
3	B	98	Total O 98 98	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ABC transporter



- Molecule 1: ABC transporter



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	47.78Å 104.07Å 106.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.74 – 2.20 46.74 – 1.86	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.74-2.20) 92.9 (46.74-1.86)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.11 (at 1.86Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.193 , 0.219 0.193 , 0.219	Depositor DCC
$R_{free}$ test set	1380 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 48.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.010 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5293	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.24	0/1262	0.48	0/1697
1	B	0.24	0/1287	0.49	0/1732
All	All	0.24	0/2549	0.49	0/3429

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1243	1281	1283	6	0
1	B	1270	1297	1297	13	0
2	A	1	0	0	0	0
3	A	103	0	0	3	1
3	B	98	0	0	5	1
All	All	2715	2578	2580	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:316:THR:O	3:B:501:HOH:O	1.87	0.93
1:A:340:GLU:OE2	3:A:601:HOH:O	1.99	0.80
1:A:431:ILE:HD13	1:B:431:ILE:HD13	1.90	0.54
1:B:379:ARG:NH2	3:B:504:HOH:O	2.42	0.53
1:B:321:LYS:HE3	1:B:420:LYS:HB2	1.95	0.48
1:A:425:ARG:HG2	3:A:652:HOH:O	2.14	0.48
1:B:316:THR:C	3:B:501:HOH:O	2.46	0.45
1:B:312:LYS:HB3	1:B:318:LEU:HD21	2.00	0.44
1:B:316:THR:HB	3:B:501:HOH:O	2.19	0.43
1:A:435:LEU:HG	1:B:362:ASP:HB3	2.01	0.43
1:A:360:TRP:HB2	1:B:431:ILE:HB	2.01	0.43
1:B:312:LYS:H	1:B:318:LEU:CD2	2.33	0.42
1:B:318:LEU:HD12	3:B:570:HOH:O	2.19	0.41
1:B:312:LYS:N	1:B:318:LEU:HD21	2.37	0.40
1:A:425:ARG:NH2	3:A:602:HOH:O	2.21	0.40
1:B:310:ILE:O	1:B:318:LEU:HD23	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:647:HOH:O	3:B:581:HOH:O[2_654]	1.91	0.29

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/190 (83%)	155 (98%)	3 (2%)	0	100	100
1	B	164/190 (86%)	160 (98%)	4 (2%)	0	100	100
All	All	322/380 (85%)	315 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/154 (86%)	128 (97%)	4 (3%)	46	58
1	B	135/154 (88%)	131 (97%)	4 (3%)	46	58
All	All	267/308 (87%)	259 (97%)	8 (3%)	46	58

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	291	GLU
1	A	312	LYS
1	A	378	GLN
1	A	406	ASP
1	B	312	LYS
1	B	314	ASP
1	B	378	GLN
1	B	404	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	161/190 (84%)	-0.38	1 (0%) 89 88	14, 32, 66, 86	0
1	B	166/190 (87%)	-0.28	3 (1%) 69 66	12, 35, 71, 106	0
All	All	327/380 (86%)	-0.33	4 (1%) 79 77	12, 34, 69, 106	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	404	ASN	4.0
1	A	286	LEU	2.5
1	B	405	GLY	2.2
1	B	316	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CL	A	501	1/1	0.98	0.12	3.81	29,29,29,29	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.