



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 04:33 am GMT

PDB ID : 5HPY
Title : Crystal Structure of RhoA.GDP.MgF3-in complex with human Myosin 9b RhoGAP domain
Authors : Yi, F.S.; Ren, J.Q.; Feng, W.
Deposited on : 2016-01-21
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3166 (2.40-2.40)
Clashscore	112137	3674 (2.40-2.40)
Ramachandran outliers	110173	3616 (2.40-2.40)
Sidechain outliers	110143	3617 (2.40-2.40)
RSRZ outliers	101464	3195 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	<div> <div>9%</div> <div> <div></div> <div>70%</div> <div>11%</div> <div>•</div> <div>17%</div> </div> </div>
1	D	232	<div> <div>11%</div> <div> <div></div> <div>68%</div> <div>14%</div> <div>18%</div> </div> </div>
2	B	185	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>17%</div> <div>• •</div> </div> </div>
2	F	185	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>14%</div> <div>• •</div> </div> </div>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5968 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-IXb.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	192	Total	C	N	O	S	0	0	0
			1501	966	257	269	9			
1	D	190	Total	C	N	O	S	0	0	0
			1496	964	254	267	11			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	185	GLY	-	expression tag	UNP Q13459
A	186	PRO	-	expression tag	UNP Q13459
A	187	GLY	-	expression tag	UNP Q13459
A	188	SER	-	expression tag	UNP Q13459
A	189	GLU	-	expression tag	UNP Q13459
A	190	PHE	-	expression tag	UNP Q13459
D	185	GLY	-	expression tag	UNP Q13459
D	186	PRO	-	expression tag	UNP Q13459
D	187	GLY	-	expression tag	UNP Q13459
D	188	SER	-	expression tag	UNP Q13459
D	189	GLU	-	expression tag	UNP Q13459
D	190	PHE	-	expression tag	UNP Q13459


- Molecule 2 is a protein called Transforming protein RhoA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	178	Total	C	N	O	S	0	1	0
			1398	884	234	270	10			
2	F	178	Total	C	N	O	S	0	0	0
			1370	866	229	265	10			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P61586
B	-2	PRO	-	expression tag	UNP P61586
B	-1	GLY	-	expression tag	UNP P61586
B	0	SER	-	expression tag	UNP P61586
B	1	GLU	-	expression tag	UNP P61586
B	2	PHE	-	expression tag	UNP P61586
B	25	ASN	PHE	engineered mutation	UNP P61586
F	-3	GLY	-	expression tag	UNP P61586
F	-2	PRO	-	expression tag	UNP P61586
F	-1	GLY	-	expression tag	UNP P61586
F	0	SER	-	expression tag	UNP P61586
F	1	GLU	-	expression tag	UNP P61586
F	2	PHE	-	expression tag	UNP P61586
F	25	ASN	PHE	engineered mutation	UNP P61586

- Molecule 3 is TRIFLUOROMAGNESATE (three-letter code: MGF) (formula: F_3Mg).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	F	Mg	0	0
			4	3	1		
3	F	1	Total	F	Mg	0	0
			4	3	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Mg 1 1	0	0
4	F	2	Total Mg 2 2	0	0

- Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C N O P 28 10 5 11 2	0	0
5	F	1	Total C N O P 28 10 5 11 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	28	Total O 28 28	0	0
6	B	45	Total O 45 45	0	0
6	D	20	Total O 20 20	0	0
6	F	43	Total O 43 43	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Unconventional myosin-IXb




• Molecule 1: Unconventional myosin-IXb




• Molecule 2: Transforming protein RhoA



• Molecule 2: Transforming protein RhoA





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.06Å 90.76Å 93.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.70 – 2.40 48.70 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.70-2.40) 99.8 (48.70-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.83 (at 2.39Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.211 , 0.246 0.211 , 0.246	Depositor DCC
R_{free} test set	1867 reflections (4.84%)	DCC
Wilson B-factor (Å ²)	51.5	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 57.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5968	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.37 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.7810e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG, MGF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/1527	0.57	0/2067
1	D	0.41	0/1523	0.55	0/2061
2	B	0.50	0/1428	0.61	0/1936
2	F	0.48	0/1397	0.59	0/1898
All	All	0.45	0/5875	0.58	0/7962

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1501	0	1526	20	0
1	D	1496	0	1535	18	0
2	B	1398	0	1374	20	0
2	F	1370	0	1325	18	0
3	B	4	0	0	0	0
3	F	4	0	0	0	0
4	B	1	0	0	0	0
4	F	2	0	0	0	0
5	B	28	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	28	0	12	0	0
6	A	28	0	0	1	0
6	B	45	0	0	0	0
6	D	20	0	0	1	0
6	F	43	0	0	3	0
All	All	5968	0	5784	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:ALA:HB1	1:A:373:ILE:HG13	1.56	0.87
1:D:351:ALA:HB1	1:D:373:ILE:HG13	1.54	0.86
2:B:118:LYS:HB3	2:B:121:LEU:HD12	1.76	0.67
1:D:284:THR:N	6:D:501:HOH:O	2.29	0.66
1:D:383:GLU:OE1	1:D:386:ARG:NH1	2.28	0.65
1:A:362:ASP:HB3	1:A:364:LEU:H	1.66	0.61
1:A:293:ARG:HD2	2:F:140:LYS:HZ1	1.67	0.59
1:A:362:ASP:HB3	1:A:364:LEU:N	2.17	0.58
1:A:311:LEU:HD11	1:A:380:LEU:HD21	1.87	0.56
2:F:85:SER:OG	2:F:87:ASP:OD1	2.13	0.56
1:A:196:GLY:HA2	1:A:197:HIS:CB	2.37	0.55
2:B:113:ILE:HG12	2:B:173:MET:HG2	1.89	0.55
1:D:284:THR:HG22	1:D:318:ASN:HD22	1.72	0.55
2:B:123:ASN:O	2:B:128:ARG:NH1	2.40	0.54
1:D:311:LEU:HD11	1:D:380:LEU:HD21	1.89	0.54
2:B:23[B]:ILE:HD12	2:B:31:PRO:HG3	1.91	0.52
1:D:363:PRO:HB2	2:F:72:LEU:HD21	1.91	0.52
2:B:46:ILE:HG22	2:B:53:VAL:HB	1.93	0.51
1:A:284:THR:OG1	1:A:287:GLN:HG2	2.10	0.51
2:F:142:GLU:HA	2:F:145:ARG:NH1	2.26	0.51
2:F:162:LYS:NZ	6:F:306:HOH:O	2.41	0.51
1:A:293:ARG:HD2	2:F:140:LYS:NZ	2.25	0.51
2:F:4:ILE:HD12	2:F:52:GLN:O	2.11	0.50
2:B:86:ILE:HG12	2:B:117:ASN:O	2.11	0.50
2:F:169:GLU:OE1	2:F:169:GLU:N	2.44	0.50
1:D:367:MET:O	1:D:370:VAL:HG12	2.11	0.50
1:D:362:ASP:OD1	1:D:365:THR:OG1	2.30	0.50
2:B:119:LYS:HB2	2:B:159:CYS:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:ASN:O	1:A:245:GLU:HG3	2.13	0.48
1:A:363:PRO:O	1:A:366:SER:HB2	2.14	0.48
1:A:288:TYR:HD2	1:A:355:LEU:HD22	1.79	0.47
1:A:386:ARG:HG3	1:A:387:LYS:N	2.29	0.47
1:A:387:LYS:O	1:A:391:LYS:HG2	2.13	0.47
2:B:19:THR:OG1	2:B:59:ASP:OD2	2.32	0.47
1:D:306:ALA:O	1:D:310:VAL:HG23	2.15	0.47
2:F:163:THR:OG1	2:F:165:ASP:HB2	2.15	0.47
1:D:370:VAL:HG13	1:D:371:LEU:HG	1.96	0.47
2:F:164:LYS:HD2	2:F:164:LYS:N	2.30	0.47
2:B:19:THR:O	2:B:23[A]:ILE:HG12	2.15	0.46
2:B:86:ILE:HD12	2:B:122:ARG:NH2	2.31	0.46
1:D:220:LEU:HD22	1:D:328:HIS:CD2	2.51	0.46
2:B:168:ARG:O	2:B:172:GLU:OE1	2.34	0.46
2:F:168:ARG:O	2:F:172:GLU:HG3	2.16	0.45
1:D:370:VAL:O	1:D:373:ILE:HG12	2.16	0.45
2:F:177:ALA:O	2:F:180:GLN:HG2	2.16	0.45
2:B:168:ARG:HE	2:B:168:ARG:HB3	1.46	0.45
2:B:24:VAL:HG12	2:B:167:VAL:HG11	1.99	0.45
1:D:248:GLN:O	1:D:252:THR:HG23	2.17	0.45
2:F:157:MET:HE1	2:F:169:GLU:HB3	1.98	0.45
1:A:289:GLY:O	1:A:293:ARG:HG3	2.17	0.44
1:A:391:LYS:O	1:A:395:ILE:HG13	2.17	0.44
2:B:28:ASP:OD1	2:B:164:LYS:NZ	2.31	0.44
1:A:367:MET:O	1:A:370:VAL:HG12	2.17	0.44
2:F:123:ASN:HA	6:F:318:HOH:O	2.17	0.43
2:F:71:PRO:HA	2:F:74:TYR:CD2	2.52	0.43
2:F:24:VAL:O	2:F:28:ASP:HA	2.18	0.43
2:F:93:GLU:O	2:F:96:PRO:HD2	2.19	0.43
1:D:284:THR:HG22	1:D:318:ASN:ND2	2.31	0.43
1:A:298:PRO:HG2	1:A:302:GLU:OE2	2.19	0.42
2:B:85:SER:OG	2:B:118:LYS:HD2	2.20	0.42
1:D:345:ALA:O	1:D:349:ILE:HG12	2.19	0.42
2:B:24:VAL:O	2:B:28:ASP:HA	2.20	0.42
2:B:8:LEU:HD11	2:B:81:LEU:HG	2.02	0.42
2:F:100:THR:HB	2:F:101:PRO:HD3	2.02	0.41
2:B:142:GLU:H	2:B:142:GLU:HG3	1.35	0.41
1:A:386:ARG:O	1:A:390:VAL:HG13	2.21	0.41
1:D:250:LEU:HD23	1:D:257:VAL:HG11	2.03	0.41
2:B:71:PRO:HA	2:B:74:TYR:CD2	2.56	0.41
2:B:166:GLY:O	2:B:169:GLU:HG2	2.21	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:241:ASN:O	1:D:245:GLU:HG3	2.21	0.40
1:D:262:PHE:HA	1:D:263:PRO:HD3	1.97	0.40
1:A:298:PRO:HG3	6:F:318:HOH:O	2.21	0.40
1:A:247:ARG:HD2	6:A:504:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/232 (80%)	179 (96%)	6 (3%)	1 (0%)	32	46
1	D	184/232 (79%)	179 (97%)	4 (2%)	1 (0%)	32	46
2	B	177/185 (96%)	173 (98%)	4 (2%)	0	100	100
2	F	176/185 (95%)	172 (98%)	4 (2%)	0	100	100
All	All	723/834 (87%)	703 (97%)	18 (2%)	2 (0%)	44	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	197	HIS
1	D	298	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	159/200 (80%)	154 (97%)	5 (3%)	45	66
1	D	162/200 (81%)	157 (97%)	5 (3%)	45	66
2	B	153/160 (96%)	151 (99%)	2 (1%)	73	87
2	F	147/160 (92%)	145 (99%)	2 (1%)	71	86
All	All	621/720 (86%)	607 (98%)	14 (2%)	56	75

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	251	GLN
1	A	299	GLU
1	A	362	ASP
1	A	370	VAL
1	A	387	LYS
2	B	142	GLU
2	B	145	ARG
1	D	212	VAL
1	D	283	MET
1	D	323	GLU
1	D	331	LYS
1	D	371	LEU
2	F	142	GLU
2	F	145	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	MGF	B	201	5,6	0,3,3	0.00	-	0,3,3	0.00	-
5	GDP	B	203	3,4	25,30,30	1.36	2 (8%)	26,47,47	2.07	6 (23%)
3	MGF	F	201	5,6	0,3,3	0.00	-	0,3,3	0.00	-
5	GDP	F	202	3,4	25,30,30	1.23	2 (8%)	26,47,47	1.99	6 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MGF	B	201	5,6	-	0/0/0/0	0/0/0/0
5	GDP	B	203	3,4	-	0/12/32/32	0/3/3/3
3	MGF	F	201	5,6	-	0/0/0/0	0/0/0/0
5	GDP	F	202	3,4	-	0/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	202	GDP	C5-C4	3.31	1.48	1.40
5	B	203	GDP	C5-C4	3.39	1.48	1.40
5	F	202	GDP	C6-C5	4.11	1.49	1.41
5	B	203	GDP	C6-C5	4.17	1.49	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	203	GDP	C5-C6-N1	-4.50	117.08	123.48
5	F	202	GDP	C5-C6-N1	-3.99	117.81	123.48
5	B	203	GDP	C6-C5-C4	-3.87	117.00	120.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	202	GDP	C6-C5-C4	-3.49	117.37	120.84
5	F	202	GDP	C4-C5-N7	-3.11	106.41	109.41
5	B	203	GDP	N3-C2-N1	-3.09	122.94	127.46
5	F	202	GDP	N3-C2-N1	-2.93	123.17	127.46
5	B	203	GDP	C4-C5-N7	-2.79	106.72	109.41
5	B	203	GDP	C2-N3-C4	3.83	119.64	115.16
5	F	202	GDP	C6-N1-C2	4.29	122.22	116.06
5	F	202	GDP	C2-N3-C4	5.04	121.04	115.16
5	B	203	GDP	C6-N1-C2	5.46	123.91	116.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/232 (82%)	0.56	21 (10%) 6 5	43, 74, 97, 109	0
1	D	190/232 (81%)	0.71	26 (13%) 3 3	45, 76, 100, 122	0
2	B	178/185 (96%)	0.32	4 (2%) 62 59	34, 59, 89, 104	0
2	F	178/185 (96%)	0.43	5 (2%) 53 51	39, 62, 100, 111	0
All	All	738/834 (88%)	0.51	56 (7%) 15 13	34, 67, 97, 122	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	252	THR	4.8
1	D	199	GLY	4.6
1	D	313	HIS	4.4
1	A	361	SER	4.2
1	A	252	THR	4.0
1	D	225	MET	4.0
1	D	212	VAL	3.8
1	D	211	SER	3.8
1	A	201	CYS	3.7
1	D	357	CYS	3.6
2	B	132	ALA	3.6
1	A	199	GLY	3.5
1	D	293	ARG	3.5
1	D	297	LEU	3.5
1	D	198	PHE	3.4
2	F	134	MET	3.2
1	D	200	VAL	3.2
1	D	255	ALA	3.2
1	D	316	GLU	3.1
1	D	253	ASP	3.0
2	F	125	GLU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	302	GLU	2.9
1	A	255	ALA	2.9
2	B	145	ARG	2.9
1	A	196	GLY	2.9
2	B	172	GLU	2.9
2	B	125	GLU	2.7
1	A	281	PRO	2.7
1	A	313	HIS	2.6
1	A	198	PHE	2.6
1	A	285	PHE	2.5
1	D	250	LEU	2.5
1	A	356	ARG	2.5
1	D	356	ARG	2.5
1	A	226	HIS	2.4
1	D	299	GLU	2.4
1	A	256	ALA	2.4
1	D	314	LEU	2.4
1	D	312	GLU	2.4
1	A	357	CYS	2.4
1	A	211	SER	2.3
1	A	257	VAL	2.3
2	F	47	GLU	2.3
1	A	395	ILE	2.3
1	D	358	PRO	2.2
1	D	302	GLU	2.2
1	A	316	GLU	2.2
2	F	132	ALA	2.2
1	A	251	GLN	2.2
1	D	364	LEU	2.1
1	A	197	HIS	2.1
1	D	249	ALA	2.1
1	D	251	GLN	2.1
1	D	391	LYS	2.1
1	D	390	VAL	2.0
2	F	44	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	GDP	F	202	28/28	0.95	0.18	0.57	39,56,62,68	0
5	GDP	B	203	28/28	0.98	0.18	0.38	34,50,58,67	0
3	MGF	F	201	4/4	0.98	0.16	-0.54	36,40,41,47	0
4	MG	F	204	1/1	0.95	0.13	-1.49	49,49,49,49	0
4	MG	B	202	1/1	0.96	0.14	-1.81	37,37,37,37	0
3	MGF	B	201	4/4	0.99	0.13	-2.12	37,38,39,40	0
4	MG	F	203	1/1	0.96	0.09	-3.40	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.