



# Full wwPDB X-ray Structure Validation Report i

Feb 14, 2017 – 03:46 am GMT

PDB ID : 4HWP  
Title : Crystal structure of E. coli Threonyl-tRNA synthetase bound to a novel inhibitor  
Authors : Hilgers, M.T.  
Deposited on : 2012-11-08  
Resolution : 1.81 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	X16	B	702	-	-	-	X

## 2 Entry composition [\(i\)](#)

There are 4 unique types of molecules in this entry. The entry contains 7245 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Threonine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	409	3354	2116	596	619	23	0	0	0
1	B	398	3251	2053	573	602	23	0	0	0


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	240	MET	-	EXPRESSION TAG	UNP P0A8M3
A	241	ALA	-	EXPRESSION TAG	UNP P0A8M3
A	643	LEU	-	EXPRESSION TAG	UNP P0A8M3
A	644	GLU	-	EXPRESSION TAG	UNP P0A8M3
A	645	HIS	-	EXPRESSION TAG	UNP P0A8M3
A	646	HIS	-	EXPRESSION TAG	UNP P0A8M3
A	647	HIS	-	EXPRESSION TAG	UNP P0A8M3
A	648	HIS	-	EXPRESSION TAG	UNP P0A8M3
A	649	HIS	-	EXPRESSION TAG	UNP P0A8M3
A	650	HIS	-	EXPRESSION TAG	UNP P0A8M3
B	240	MET	-	EXPRESSION TAG	UNP P0A8M3
B	241	ALA	-	EXPRESSION TAG	UNP P0A8M3
B	643	LEU	-	EXPRESSION TAG	UNP P0A8M3
B	644	GLU	-	EXPRESSION TAG	UNP P0A8M3
B	645	HIS	-	EXPRESSION TAG	UNP P0A8M3
B	646	HIS	-	EXPRESSION TAG	UNP P0A8M3
B	647	HIS	-	EXPRESSION TAG	UNP P0A8M3
B	648	HIS	-	EXPRESSION TAG	UNP P0A8M3
B	649	HIS	-	EXPRESSION TAG	UNP P0A8M3
B	650	HIS	-	EXPRESSION TAG	UNP P0A8M3

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0

- Molecule 3 is N-[3-(4-AMINO-2-METHYLQUINAZOLIN-7-YL)PHENYL]SULFONYL]-L-THREONINAMIDE (three-letter code: X16) (formula: C<sub>19</sub>H<sub>21</sub>N<sub>5</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O S 29 19 5 4 1	0	0
3	B	1	Total C N O S 29 19 5 4 1	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	287	Total O 287 287	0	0
4	B	293	Total O 293 293	0	0



### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Threonine-tRNA ligase



- Molecule 1: Threonine-tRNA ligase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.16 Å    110.14 Å    115.45 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	43.08 – 1.81 43.08 – 1.81	Depositor EDS
% Data completeness (in resolution range)	98.7 (43.08-1.81) 98.7 (43.08-1.81)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle^1$	3.04 (at 1.81 Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
$R$ , $R_{free}$	0.194 , 0.213 0.193 , 0.212	Depositor DCC
$R_{free}$ test set	4988 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.0	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.014 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7245	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, X16

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.74	4/3431 (0.1%)	0.81	7/4621 (0.2%)
1	B	0.74	4/3322 (0.1%)	0.80	5/4473 (0.1%)
All	All	0.74	8/6753 (0.1%)	0.80	12/9094 (0.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	434	TRP	CD2-CE2	5.93	1.48	1.41
1	A	266	TRP	CD2-CE2	5.92	1.48	1.41
1	B	478	TRP	CD2-CE2	5.80	1.48	1.41
1	A	310	TRP	CD2-CE2	5.59	1.48	1.41
1	B	310	TRP	CD2-CE2	5.48	1.48	1.41
1	B	304	TRP	CD2-CE2	5.43	1.47	1.41
1	A	304	TRP	CD2-CE2	5.32	1.47	1.41
1	A	536	TRP	CD2-CE2	5.29	1.47	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	611	ARG	NE-CZ-NH1	9.45	125.02	120.30
1	A	611	ARG	NE-CZ-NH2	-8.64	115.98	120.30
1	A	325	ARG	NE-CZ-NH2	-7.24	116.68	120.30
1	B	518	MET	CG-SD-CE	-6.35	90.05	100.20
1	A	325	ARG	NE-CZ-NH1	6.03	123.31	120.30
1	B	486	ASP	CB-CG-OD1	5.89	123.60	118.30
1	A	375	ARG	NE-CZ-NH2	-5.88	117.36	120.30
1	B	263	MET	O-C-N	-5.79	113.44	122.70
1	A	486	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	512	ARG	NE-CZ-NH1	5.22	122.91	120.30
1	B	354	ARG	NE-CZ-NH1	-5.18	117.71	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	243	ASP	CB-CG-OD2	5.16	122.94	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3354	0	3267	43	0
1	B	3251	0	3185	53	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	29	0	20	5	0
3	B	29	0	20	5	0
4	A	287	0	0	12	1
4	B	293	0	0	12	1
All	All	7245	0	6492	92	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:490:PRO:HG3	1:B:509:MET:CE	1.79	1.13
1:A:263:MET:HE3	4:B:802:HOH:O	1.57	1.05
1:A:242:ARG:HH22	1:A:250:GLN:HE22	1.04	1.01
1:B:490:PRO:HG3	1:B:509:MET:HE2	1.43	0.98
1:A:255:HIS:HD1	1:A:267:HIS:HE1	0.98	0.93
1:A:255:HIS:HD1	1:A:267:HIS:CE1	1.86	0.92
1:A:528:GLU:OE1	1:A:649:HIS:HD2	1.59	0.86
1:B:376:VAL:H	3:B:702:X16:HN2A	1.21	0.86
1:A:376:VAL:H	3:A:702:X16:HN2A	1.24	0.85
1:B:473:CYS:SG	4:B:1068:HOH:O	2.34	0.85
1:B:638:LYS:HB2	4:B:1085:HOH:O	1.77	0.83

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:609:ARG:HD2	4:A:1005:HOH:O	1.77	0.83
1:B:255:HIS:ND1	1:B:267:HIS:HE1	1.76	0.83
1:B:473:CYS:HB3	4:B:1068:HOH:O	1.79	0.82
1:B:453:GLU:CD	4:B:1076:HOH:O	2.19	0.79
1:B:419:LYS:HE2	4:B:1076:HOH:O	1.84	0.77
1:B:490:PRO:HG3	1:B:509:MET:HE3	1.67	0.76
1:B:473:CYS:CB	4:B:1068:HOH:O	2.32	0.76
1:A:329:ILE:HD11	1:B:329:ILE:HD11	1.71	0.72
1:B:484:GLN:HE22	3:B:702:X16:HN5	1.38	0.69
1:A:572:ASP:OD2	1:A:585:HIS:HE1	1.76	0.67
1:A:403:LEU:HG	1:A:407:MET:HE3	1.77	0.67
1:A:403:LEU:HD11	1:A:407:MET:HE2	1.75	0.67
1:B:572:ASP:OD2	1:B:585:HIS:HE1	1.78	0.66
1:A:268:ASN:HD22	1:B:292:GLU:H	1.42	0.66
1:B:585:HIS:HD2	4:B:815:HOH:O	1.79	0.64
1:B:594:LEU:HD22	1:B:608:VAL:HG22	1.80	0.64
1:A:252:ASP:OD1	1:A:267:HIS:HD2	1.79	0.64
1:A:585:HIS:HD2	4:A:922:HOH:O	1.82	0.63
1:A:292:GLU:H	1:B:268:ASN:HD22	1.45	0.63
1:B:620:ASP:OD2	1:B:623:GLU:HB2	1.98	0.63
1:B:255:HIS:ND1	1:B:267:HIS:CE1	2.65	0.62
1:A:455:GLN:NE2	4:A:993:HOH:O	2.18	0.61
1:A:528:GLU:OE1	1:A:649:HIS:CD2	2.48	0.61
1:A:242:ARG:HH22	1:A:250:GLN:NE2	1.87	0.61
1:B:419:LYS:HD3	1:B:453:GLU:HG3	1.83	0.60
1:B:382:ASP:HB3	1:B:518:MET:CE	2.31	0.60
1:B:252:ASP:OD1	1:B:267:HIS:HD2	1.84	0.59
1:B:556:ASN:HD21	1:B:573:LEU:HD11	1.68	0.58
1:B:310:TRP:O	1:B:314:LYS:HB2	2.04	0.58
1:A:405:TYR:CE1	1:A:416:ILE:HD12	2.39	0.57
1:B:325:ARG:HD3	4:B:1093:HOH:O	2.03	0.57
1:A:475:ASP:OD1	1:A:649:HIS:HE1	1.88	0.56
1:A:419:LYS:HE2	4:A:972:HOH:O	2.08	0.54
1:A:403:LEU:HD11	1:A:407:MET:CE	2.37	0.54
1:B:598:ASP:O	1:B:602:GLU:HG2	2.07	0.53
1:B:490:PRO:CG	1:B:509:MET:HE3	2.37	0.53
1:A:257:GLN:HG3	1:A:265:PHE:CE2	2.44	0.52
1:B:403:LEU:HD11	1:B:407:MET:HE2	1.90	0.51
1:A:255:HIS:ND1	1:A:267:HIS:HE1	1.84	0.51
1:B:403:LEU:HG	1:B:407:MET:HE3	1.92	0.51
1:B:638:LYS:O	1:B:639:GLN:HB2	2.11	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:490:PRO:CG	1:B:509:MET:CE	2.71	0.50
1:A:404:VAL:HA	1:A:514:ILE:HG23	1.94	0.49
1:A:637:LEU:HD13	1:A:645:HIS:CE1	2.48	0.49
1:B:412:GLY:HA3	1:B:637:LEU:HD11	1.95	0.48
1:A:325:ARG:HD3	4:A:1080:HOH:O	2.13	0.48
1:B:342:ASN:HD21	1:B:495:ALA:HA	1.79	0.47
1:B:638:LYS:CB	4:B:1085:HOH:O	2.48	0.47
1:B:620:ASP:OD2	1:B:623:GLU:CB	2.63	0.47
1:B:637:LEU:HB2	4:B:958:HOH:O	2.13	0.47
1:A:363:ARG:HB2	3:A:702:X16:H29B	1.97	0.47
1:B:242:ARG:HB3	1:B:247:ILE:HD11	1.97	0.47
1:B:301:ARG:O	1:B:305:GLU:HG3	2.15	0.46
1:A:292:GLU:H	1:B:268:ASN:ND2	2.12	0.46
1:B:382:ASP:HB3	1:B:518:MET:HE1	1.96	0.46
1:B:503:GLU:OE1	1:B:505:LYS:NZ	2.39	0.45
1:A:376:VAL:N	3:A:702:X16:HN2A	2.04	0.45
1:A:298:MET:HA	4:A:1079:HOH:O	2.16	0.45
1:A:543:VAL:HG23	1:A:590:VAL:HG11	1.97	0.45
1:B:375:ARG:HA	3:B:702:X16:N28	2.32	0.44
1:A:403:LEU:CG	1:A:407:MET:HE3	2.46	0.44
3:A:702:X16:H13	3:A:702:X16:H1B	2.00	0.44
1:B:285:LEU:HB3	1:B:290:TYR:HB2	1.99	0.44
1:A:375:ARG:NH1	4:A:1070:HOH:O	2.30	0.44
1:A:544:ILE:HD13	1:A:558:LEU:HB3	2.00	0.44
1:A:242:ARG:N	4:A:1037:HOH:O	2.51	0.43
1:A:638:LYS:HD3	4:A:982:HOH:O	2.17	0.43
1:A:339:GLN:HB3	1:B:257:GLN:OE1	2.18	0.43
1:B:242:ARG:HH21	1:B:246:LYS:HD3	1.82	0.43
1:A:325:ARG:CD	4:A:1080:HOH:O	2.66	0.43
1:B:288:TYR:CE1	1:B:403:LEU:HD13	2.55	0.42
1:B:376:VAL:N	3:B:702:X16:HN2A	2.00	0.42
1:A:590:VAL:O	1:A:611:ARG:HB3	2.20	0.42
1:A:375:ARG:HA	3:A:702:X16:N28	2.34	0.42
1:B:363:ARG:HB2	3:B:702:X16:H29B	2.02	0.41
1:A:642:GLU:N	4:A:1028:HOH:O	2.47	0.41
1:B:543:VAL:HG11	1:B:585:HIS:CE1	2.55	0.41
1:B:325:ARG:CD	4:B:1093:HOH:O	2.66	0.41
1:B:288:TYR:CZ	1:B:403:LEU:HD13	2.56	0.41
1:A:257:GLN:NE2	4:A:859:HOH:O	2.53	0.40
1:B:335:PRO:O	1:B:339:GLN:HG2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1074:HOH:O	4:B:925:HOH:O[3_545]	1.32	0.88

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	407/411 (99%)	401 (98%)	6 (2%)	0	100 100
1	B	396/411 (96%)	391 (99%)	5 (1%)	0	100 100
All	All	803/822 (98%)	792 (99%)	11 (1%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	364/365 (100%)	353 (97%)	11 (3%)	46 30
1	B	353/365 (97%)	342 (97%)	11 (3%)	45 29
All	All	717/730 (98%)	695 (97%)	22 (3%)	45 29

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	313	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	334	CYS
1	A	347	SER
1	A	502	ASN
1	A	512	ARG
1	A	549	ASP
1	A	558	LEU
1	A	578	ILE
1	A	599	LYS
1	A	633	ARG
1	A	640	LEU
1	B	259	GLU
1	B	267	HIS
1	B	276	GLU
1	B	287	GLU
1	B	334	CYS
1	B	426	LYS
1	B	504	ARG
1	B	512	ARG
1	B	599	LYS
1	B	618	SER
1	B	620	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	250	GLN
1	A	257	GLN
1	A	267	HIS
1	A	268	ASN
1	A	291	GLN
1	A	342	ASN
1	A	556	ASN
1	A	585	HIS
1	A	649	HIS
1	B	267	HIS
1	B	268	ASN
1	B	291	GLN
1	B	312	ASN
1	B	342	ASN
1	B	556	ASN
1	B	585	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	X16	A	702	2	29,31,31	2.70	7 (24%)	40,46,46	2.02	14 (35%)
3	X16	B	702	2	29,31,31	2.42	5 (17%)	40,46,46	1.82	9 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	X16	A	702	2	-	0/23/23/23	0/3/3/3
3	X16	B	702	2	-	0/23/23/23	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	X16	C12-S9	-10.56	1.61	1.76

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	702	X16	C12-S9	-9.44	1.62	1.76
3	A	702	X16	O11-S9	-6.58	1.36	1.43
3	A	702	X16	O10-S9	-5.11	1.38	1.43
3	B	702	X16	O11-S9	-2.79	1.40	1.43
3	B	702	X16	S9-N5	-2.57	1.59	1.64
3	A	702	X16	S9-N5	-2.34	1.59	1.64
3	A	702	X16	C23-C22	-2.01	1.38	1.41
3	A	702	X16	C21-C22	2.89	1.47	1.42
3	B	702	X16	C21-C22	3.39	1.47	1.42
3	B	702	X16	C27-C21	4.27	1.49	1.45
3	A	702	X16	C27-C21	4.69	1.49	1.45

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	702	X16	O10-S9-O11	-5.24	112.84	119.55
3	A	702	X16	O10-S9-O11	-4.36	113.97	119.55
3	B	702	X16	C21-C27-N26	-4.33	117.50	121.94
3	A	702	X16	C21-C22-N24	-4.20	118.50	122.84
3	A	702	X16	C4-N5-S9	-3.17	120.85	124.21
3	A	702	X16	C13-C12-C17	-3.01	116.84	120.61
3	B	702	X16	C21-C22-N24	-2.99	119.75	122.84
3	A	702	X16	C21-C27-N26	-2.72	119.14	121.94
3	B	702	X16	O10-S9-C12	-2.35	105.04	107.95
3	A	702	X16	C20-C21-C22	2.00	120.43	118.34
3	B	702	X16	C19-C18-C23	2.03	121.23	118.19
3	B	702	X16	C20-C21-C22	2.06	120.48	118.34
3	A	702	X16	C13-C12-S9	2.14	122.13	119.77
3	A	702	X16	C25-N26-C27	2.23	122.00	118.16
3	B	702	X16	C23-C22-N24	2.24	122.17	118.75
3	A	702	X16	C16-C17-C12	2.29	121.75	119.89
3	A	702	X16	O6-C2-C3	2.52	114.78	109.04
3	A	702	X16	O11-S9-C12	2.64	111.22	107.95
3	B	702	X16	C25-N26-C27	3.11	123.53	118.16
3	A	702	X16	C23-C22-N24	3.23	123.66	118.75
3	A	702	X16	C12-S9-N5	3.47	110.61	105.91
3	A	702	X16	C14-C13-C12	3.62	122.77	118.96
3	B	702	X16	C12-S9-N5	3.81	111.07	105.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	702	X16	5	0
3	B	702	X16	5	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	409/411 (99%)	0.22	31 (7%) 15 12	17, 25, 57, 88	0
1	B	398/411 (96%)	-0.07	11 (2%) 53 48	17, 25, 45, 88	0
All	All	807/822 (98%)	0.08	42 (5%) 28 23	17, 25, 54, 88	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	601	VAL	4.4
1	A	575	ASN	4.2
1	A	602	GLU	3.9
1	B	503	GLU	3.8
1	B	498	VAL	3.8
1	A	548	THR	3.6
1	A	502	ASN	3.5
1	A	310	TRP	3.5
1	B	473	CYS	3.5
1	B	501	ASP	3.0
1	B	500	GLU	2.9
1	A	549	ASP	2.9
1	A	599	LYS	2.8
1	B	345	LEU	2.8
1	A	553	GLU	2.8
1	A	554	TYR	2.8
1	A	501	ASP	2.8
1	A	613	GLY	2.7
1	A	578	ILE	2.7
1	A	547	ILE	2.7
1	A	611	ARG	2.7
1	A	621	VAL	2.6
1	A	415	LYS	2.6
1	A	612	ARG	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	500	GLU	2.5
1	A	550	SER	2.5
1	A	604	GLY	2.5
1	A	610	THR	2.5
1	A	598	ASP	2.5
1	B	549	ASP	2.5
1	A	597	GLY	2.4
1	A	358	PHE	2.4
1	A	552	SER	2.3
1	A	313	TYR	2.3
1	A	277	LEU	2.2
1	A	273	ILE	2.1
1	B	271	TRP	2.1
1	B	288	TYR	2.1
1	A	643	LEU	2.1
1	B	637	LEU	2.1
1	B	639	GLN	2.1
1	A	546	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	X16	B	702	29/29	0.96	0.15	2.04	25,27,29,32	0
3	X16	A	702	29/29	0.94	0.17	1.73	25,27,28,31	0
2	ZN	B	701	1/1	1.00	0.10	0.08	21,21,21,21	0
2	ZN	A	701	1/1	1.00	0.10	-1.41	22,22,22,22	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.